



UNIVERSIDAD
DE GRANADA



IMAG
INSTITUTO DE MATEMÁTICAS
Universidad de Granada



EXCELENCIA
MARÍA
DE MAEZTU

MULTIPLE ORTHOGONAL POLYNOMIALS IN TWO VARIABLES

Juan Antonio Villegas-Recio
in partnership with Lidia Fernández

Departamento de Matemática Aplicada and Instituto de Matemáticas IMAG
Universidad de Granada, Spain.

D2PO 2023
14th December 2023

MULTIPLE ORTHOGONALITY

Polynomials satisfy orthogonality relations with respect to r measures: μ_1, \dots, μ_r .

Let $\langle \cdot, \cdot \rangle_j$ be the respective integral inner product ($j = 1, \dots, r$) and

$\vec{n} = (n_1, \dots, n_r) \in \mathbb{N}^r$, $|\vec{n}| = n_1 + \dots + n_r$.

Type II Multiple Orthogonal Polynomials

$P_{\vec{n}}(x)$ is a monic polynomial with $\deg(P_{\vec{n}}) = |\vec{n}|$ and

$$\left\langle P_{\vec{n}}, x^k \right\rangle_j = 0, \quad k = 0, \dots, n_j - 1, \quad j = 1, \dots, r \quad (1)$$

Type I Multiple Orthogonal Polynomials

$(A_{\vec{n},1}(x), \dots, A_{\vec{n},r}(x))$, vector of polynomials, $\deg(A_{\vec{n},j}) \leq n_j - 1$, ($j = 1, \dots, r$) and

$$\sum_{j=1}^r \left\langle A_{\vec{n},j}, x^k \right\rangle_j = \begin{cases} 0 & \text{if } k = 0, \dots, |\vec{n}| - 2 \\ 1 & \text{if } k = |\vec{n}| - 1. \end{cases} \quad (2)$$

BIVARIATE OP

$$\mathbb{X}_j = (x^j, x^{j-1}y, \dots, y^j)^t.$$

A column polynomial vector of degree n can be represented as

$$\mathbb{P}_n = G_{n,n}\mathbb{X}_n + G_{n,n-1}\mathbb{X}_{n-1} + \dots + G_{n,1}\mathbb{X}_1 + G_{n,0}\mathbb{X}_0,$$

where $G_{n,j}$ are matrices of size $(n+1) \times (j+1)$.

Given a bidimensional measure $\mu(x, y)$, $\text{supp}(\mu) = \Omega \subseteq \mathbb{R}^2$, we can extend the definition of inner product $\langle f, g \rangle_\mu$ to column vectors.

$F = (f_1, f_2, \dots, f_n)^t$, $G = (g_1, g_2, \dots, g_m)^t$ column vectors of functions. Then we define

$$\langle F, G \rangle_\mu := \mathcal{L}_\mu[F \cdot G^T] = \int_\Omega F \cdot G^T d\mu = \left(\int_\Omega f_i \cdot g_j d\mu \right)_{i,j=1}^{n,m}. \quad (3)$$

BIVARIATE MULTIPLE ORTHOGONAL POLYNOMIALS

Type II Multiple Orthogonal Polynomials

$$\mathbb{P}_{\vec{n}}^n = \mathbb{X}_n + \sum_{k=0}^{n-1} G_{n,k} \mathbb{X}_k$$

which satisfies

$$\langle \mathbb{P}_{\vec{n}}, \mathbb{X}_k \rangle_j = 0_{(n+1) \times (k+1)}, \quad k = 0, \dots, n_j - 1, j = 1, \dots, r. \quad (4)$$

Type I Multiple Orthogonal Polynomials

$$\mathbb{A}_{\vec{n},j} = (A_{\vec{n},j}^{(1)}(x, y), \dots, A_{\vec{n},j}^{(n)}(x, y))^t \quad (\deg A_{\vec{n},j}^{(i)} \leq n_j - 1) \quad (j = 1, \dots, r)$$

satisfying

$$\sum_{j=1}^r \langle \mathbb{X}_k, \mathbb{A}_{\vec{n},j} \rangle_j = \begin{cases} 0_{(k+1) \times n} & \text{if } k = 0, \dots, n-2 \\ I_n & \text{if } k = n-1 \end{cases} \quad (5)$$


Type I and Type II MOP are equivalent: $\exists \mathbb{P}_{\vec{n}}^n \Leftrightarrow \exists \mathbb{A}_{\vec{n},j} \quad (j = 1, \dots, r).$

ADMISSIBLE MULTI-INDICES

A multi-index \vec{n} is **admissible** if there exist a number $n \in \mathbb{N}_0$ such that:

$$n(n+1) = \sum_{j=1}^r n_j(n_j+1). \quad (6)$$

This number n is the degree of type II polynomials $\mathbb{P}_{\vec{n}}^n$ and the size of type I polynomial vectors $\mathbb{A}_{\vec{n},j}$.



BIORTHOGONALITY

There is a relation between type I and type II MOP:

Theorem 1

Let μ_1, \dots, μ_r be a perfect system of r 2-dimensional measures. Given \vec{n}, \vec{m} multi-indices satisfying (6) for n and m , respectively, the following biorthogonality holds for type I and type II MOP:

$$\langle \mathbb{P}_{\vec{n}}^n, \mathbb{Q}_{\vec{m}} \rangle = \begin{cases} 0_{(n+1) \times m} & \text{if } \vec{m} \leq \vec{n} \\ 0_{(n+1) \times m} & \text{if } n \leq m - 2 \\ I_{n+1} & \text{if } n = m - 1. \end{cases} \quad (7)$$

NEAREST NEIGHBOUR RELATION

Generalisation of the TTRR satisfied by any OPS.

Theorem 2

Let μ_1, \dots, μ_r be a perfect system of 2-dimensional measures. Let $\vec{n} \in \mathbb{N}^r$ be a multi-index satisfying (6) for $n \in \mathbb{N}$ and let us consider a path $\{\vec{m}_k : k = 0, \dots, n+1\}$ where $\vec{m}_0 = \vec{0}$, $\vec{m}_n = \vec{n}$, each \vec{m}_k satisfies (6) for k and $\vec{m}_k \leq \vec{m}_{k+1}$ for $k = 0, \dots, n$. Then, there exist matrices A_0, \dots, A_r of sizes $(n+1) \times (n-j+1)$, $(j = 0, \dots, r)$ such that

$$x\mathbb{P}_{\vec{n}}^n = L_{n+1,1}\mathbb{P}_{\vec{m}_{n+1}}^{n+1} + A_0\mathbb{P}_{\vec{n}}^n + \sum_{j=1}^r A_j\mathbb{P}_{\vec{m}_{n-j}}^{n-j}, \quad (8)$$

where $L_{n+1,1} = (I_{n+1}|0_{(n+1) \times 1})$. In addition, $A_j = \langle x\mathbb{P}_{\vec{n}}^n, \mathbb{Q}_{\vec{m}_{n-j+1}} \rangle$.

FUTURE RESEARCH TOPICS

- Find a Nearest Neighbour Relation for more general paths and other types of Nearest Neighbour Relations.
- Christoffel-Darboux formula.
- Look for applications to Hermite-Padé rational approximation and random matrices.
- Generalise existing examples to the bivariate case: Jacobi-Piñeiro, Jacobi-Angelesco, OP defined in the Simplex, etc.
- ...

ACKNOWLEDGEMENTS

Universidad de Granada; IMAG-María de Maeztu, grant "IMAG CEX2020-001105-M"; GOYA: "Grupo de Ortogonalidad Y Aplicaciones" and "Departamento de Matemática Aplicada".

Universidad de Almería; TAPO: "Teoría de Aproximación y Polinomios Ortogonales" and "Departamento de Matemáticas".



UNIVERSIDAD
DE GRANADA



IMAG
INSTITUTO DE MATEMÁTICAS
Universidad de Granada



UNIVERSIDAD
DE ALMERÍA



EXCELENCIA
MARÍA
DE MAEZTU

