





Engineering the Next Generation of Cardiometabolic Care


A Structural and Clinical Analysis of MariTide (maridebart cafraglutide)

Phase 2 Efficacy, Receptology, and the MARITIME Phase 3 Roadmap




The current standard of care faces twin barriers of dosing fatigue and weight plateaus

Current Landscape



- Market dominated by single and dual agonists (GLP-1/GIP)
- Requires high-frequency weekly dosing
- Patients often hit an efficacy plateau within 52 to 72 weeks

The MariTide Imperative



- A shift to peptide-antibody conjugates
- Enables monthly (or less frequent) dosing
- Explores novel GIP antagonism to bypass historical weight plateaus

Human genetics reveal GIP inhibition as a master switch for weight regulation

1. The Genetic Insight




deCODE Genetics identified that natural genetic inhibition of the glucose-dependent insulintropic polypeptide (GIP) receptor strongly correlates with reduced body mass.

2. The Biological Paradox




While competitors focused on agonizing (activating) GIP alongside GLP-1, Amgen's genetic data suggested the exact opposite approach: blocking GIP pathways.

3. The Synergistic Solution



Pre-clinical models confirmed that simultaneously activating GLP-1 and inhibiting GIP yields a substantially stronger weight-loss effect than targeting either receptor alone.

A pioneering peptide-antibody conjugate built for durability




The structural durability of this conjugate molecule transforms the treatment paradigm from weekly injections to true once-monthly (or less frequent) administration.

MariTide diverges entirely from the standard incretin pharmacology playbook

	GLP-1 Action	GIP Action	Glucagon Action	Dosing Frequency
Semaglutide	● Agonist	● Neutral	● Neutral	Weekly
Tirzepatide	● Agonist	● Agonist	● Neutral	Weekly
Retatrutide	● Agonist	● Agonist	● Agonist	Weekly
MariTide	● Agonist	● Antagonist	● Neutral	Monthly

Key Insight: MariTide is the only late-stage candidate utilizing GIP antagonism combined with an ultra-long-acting monthly profile.

Phase 2 data demonstrates profound weight reduction without a 52-week plateau



Phase 2 Weight Loss at 52 Weeks

Efficacy Estimand:

- Up to 20% (Cohort A)
- Up to 17% (Cohort B)

Treatment Policy Estimand (ITT):

- 12.3% – 16.2% (Cohort A)
- 8.4% – 12.3% (Cohort B)

Trajectory indicates high potential for further weight reduction beyond 52 weeks.

Weight reduction drives compounding systemic cardiometabolic benefits



Glycemic Control

Up to 2.2% reduction in HbA1c for patients in the Type 2 Diabetes cohort.



Inflammation

Significant reductions in high-sensitivity C-reactive protein (hs-CRP).



Cardiovascular

Meaningful improvements in systolic blood pressure.




Adiposity

Sustained reductions in waist circumference and select lipid parameters.

The Phase 2 data confirms that MariTide addresses the holistic cardiometabolic burden, validating its progression into expansive multi-indication trials.

Optimized dose escalation preserves efficacy while neutralizing GI tolerability barriers



Tolerability Outcomes

- **Measurement:** Tracked using the rigorous MINVR (modified index of nausea/vomiting/retching) daily patient reporting tool.
- **Phase 1 PK-LDI:** 0% discontinuation due to GI AEs at any time during the study using the optimized low-dose initiation.
- **Phase 2:** Discontinuation rates due to GI AEs dropped to 7.8% (or lower) in dose-escalation arms.


An uncompromising internal efficacy bar consolidates Amgen's pipeline

AMG 786

(Phase 1, dropped early 2024)

AMG 513

(Phase 1, dropped August 2026)




Amgen discontinued these early-stage assets—despite active clinical holds being lifted—because **they could not meet the unprecedented internal efficacy and durability bar** established by MariTide.

MariTide Phase 3 Scale-Up

Resources and capital are now aggressively concentrated on accelerating the MARITIME multi-indication clinical program.

The MARITIME Phase 3 Program: Mapping a comprehensive metabolic platform



The blueprint for a market disruption

The Next Era of Metabolic Care

Structural Innovation

The pioneering peptide-antibody conjugate scaffold shatters the weekly dosing barrier, introducing true once-monthly administration.

Pharmacological Divergence

Leveraging genetic insights to antagonize GIP while agonizing GLP-1 breaks the 52-week efficacy plateau seen in current standards of care.

Systemic Scale

The aggressive expansion of the MARITIME program across CV, HF, and OSA positions MariTide not just as a weight-loss intervention, but as a foundational cardiometabolic therapy.