

T4: Nonparametric and distance-based learning

Fundamentos del Aprendizaje Automático

Curso 2025/2026

Structure

- ① Introduction
 - Contextualization
- ② Density estimation
 - Histogram approach
 - Parzen windows
 - k_n -Nearest Neighbor estimator
 - Final remarks
- ③ The Nearest Neighbor rule
 - Formulation
 - Metrics
 - The k -Nearest-Neighbor rule
- ④ Other models
 - Decision tree
 - Support Vector Machine

Outline

1 Introduction

Contextualization

2 Density estimation

Histogram approach

Parzen windows

k_n -Nearest Neighbor estimator

Final remarks

3 The Nearest Neighbor rule

Formulation

Metrics

The k -Nearest-Neighbor rule

4 Other models

Decision tree

Support Vector Machine

Bayesian statistical learning

$$P(\omega|\mathbf{x}) = \frac{p(\mathbf{x}|\omega) \cdot P(\omega)}{p(\mathbf{x})}$$

Bayesian statistical learning

$$P(\omega|\mathbf{x}) = \frac{p(\mathbf{x}|\omega) \cdot P(\omega)}{p(\mathbf{x})}$$

- **Ideal** case: $p(\mathbf{x}|\omega)$ and $P(\omega)$ are known and well defined

Bayesian statistical learning

$$P(\omega|\mathbf{x}) = \frac{p(\mathbf{x}|\omega) \cdot P(\omega)}{p(\mathbf{x})}$$

- **Ideal** case: $p(\mathbf{x}|\omega)$ and $P(\omega)$ are known and well defined
- **Practical** case: $p(\mathbf{x}|\omega)$ and $P(\omega)$ are estimated from assortment \mathcal{D}

Bayesian statistical learning

$$P(\omega|\mathbf{x}) = \frac{p(\mathbf{x}|\omega) \cdot P(\omega)}{p(\mathbf{x})}$$

- **Ideal** case: $p(\mathbf{x}|\omega)$ and $P(\omega)$ are known and well defined
- **Practical** case: $p(\mathbf{x}|\omega)$ and $P(\omega)$ are estimated from assortment \mathcal{D}

How is it estimated?

Approach	Module	Prior $P(\omega)$	Likelihood $p(\mathbf{x} \omega)$	
			Assumption	Estimation
<i>Parametric</i>	T2	Frequentist	Multivariate distribution Statistical independence and univariate distributions	Maximum Likelihood Estimation
<i>Nonparametric</i>	T4	Frequentist	No distribution is assumed	Histogram, Parzen windows, Nearest Neighbor

Nonparametric learning

Two scenarios:

Nonparametric learning

Two scenarios:

1. Estimate *likelihoods* $p(\mathbf{x}|\omega)$ and *priors* $P(\omega)$

Nonparametric learning

Two scenarios:

1. Estimate *likelihoods* $p(\mathbf{x}|\omega)$ and *priors* $P(\omega)$

2. Estimate *posteriors* $P(\omega|\mathbf{x})$

Nonparametric learning

Two scenarios:

1. Estimate *likelihoods* $p(\mathbf{x}|\omega)$ and *priors* $P(\omega)$
 - **Key procedure:** (probability) density estimation
2. Estimate *posteriors* $P(\omega|\mathbf{x})$

Nonparametric learning

Two scenarios:

1. Estimate *likelihoods* $p(\mathbf{x}|\omega)$ and *priors* $P(\omega)$
 - **Key procedure:** (probability) density estimation
 - Reconstructing the probability **density function** of a random variable
 - **Independent** of the true distribution
2. Estimate *posteriors* $P(\omega|\mathbf{x})$

Nonparametric learning

Two scenarios:

1. Estimate *likelihoods* $p(\mathbf{x}|\omega)$ and *priors* $P(\omega)$
 - **Key procedure:** (probability) density estimation
 - Reconstructing the probability **density function** of a random variable
 - **Independent** of the true distribution
 - The function **approximates** the **true likelihood** $\Rightarrow \hat{p}(\mathbf{x}|\omega)$
2. Estimate *posteriors* $P(\omega|\mathbf{x})$

Nonparametric learning

Two scenarios:

1. Estimate *likelihoods* $p(\mathbf{x}|\omega)$ and *priors* $P(\omega)$
 - **Key procedure:** (probability) density estimation
 - Reconstructing the probability **density function** of a random variable
 - **Independent** of the true distribution
 - The function **approximates** the **true likelihood** $\Rightarrow \hat{p}(\mathbf{x}|\omega)$
2. Estimate *posteriors* $P(\omega|\mathbf{x})$
 - Avoids *likelihood* and *prior* estimation

Outline

① Introduction

Contextualization

② Density estimation

Histogram approach

Parzen windows

k_n -Nearest Neighbor estimator

Final remarks

③ The Nearest Neighbor rule

Formulation

Metrics

The k -Nearest-Neighbor rule

④ Other models

Decision tree

Support Vector Machine

Formulation


- Data assortment $\mathcal{D} = \{(\mathbf{x}_i, \omega_i)\}_{i=1}^{|\mathcal{D}|} \subset \mathbb{R}^d \times \mathcal{W}$

Formulation

- Data **assortment** $\mathcal{D} = \{(\mathbf{x}_i, \omega_i)\}_{i=1}^{|\mathcal{D}|} \subset \mathbb{R}^d \times \mathcal{W}$
 - **Goal:** **estimate** the unknown $p(\mathbf{x}|\omega)$ for each label



Formulation

- Data assortment $\mathcal{D} = \{(\mathbf{x}_i, \omega_i)\}_{i=1}^{|\mathcal{D}|} \subset \mathbb{R}^d \times \mathcal{W}$
 → **Goal:** estimate the unknown $p(\mathbf{x}|\omega)$ for each label





Formulation

- Data assortment $\mathcal{D} = \{(\mathbf{x}_i, \omega_i)\}_{i=1}^{|\mathcal{D}|} \subset \mathbb{R}^d \times \mathcal{W}$
 → **Goal:** estimate the unknown $p(\mathbf{x}|\omega)$ for each label



Formulation



- Data assortment $\mathcal{D} = \{(\mathbf{x}_i, \omega_i)\}_{i=1}^{|\mathcal{D}|} \subset \mathbb{R}^d \times \mathcal{W}$
 → **Goal:** estimate the unknown $p(\mathbf{x}|\omega)$ for each label




- Histogram estimator:
 → Model $p(\mathbf{x}|\omega)$ as a histogram
 → **Process:**
 1. Divide data space into regions
 2. Count samples per region
 3. Density as a ratio

Formulation

- Data assortment $\mathcal{D} = \{(\mathbf{x}_i, \omega_i)\}_{i=1}^{|\mathcal{D}|} \subset \mathbb{R}^d \times \mathcal{W}$
 → **Goal:** estimate the unknown $p(\mathbf{x}|\omega)$ for each label



- Histogram estimator:
 → Model $p(\mathbf{x}|\omega)$ as a histogram
 → **Process:**
 1. Divide data space into regions
 2. Count samples per region
 3. Density as a ratio



Formulation

- Assume a set of regions $\{\mathcal{R}_1, \dots, \mathcal{R}_N\}$ that divide the data space

Formulation

- Assume a set of regions $\{\mathcal{R}_1, \dots, \mathcal{R}_N\}$ that divide the data space
- The probability of $\mathbf{x} \in \mathcal{D}_\omega$ being in region \mathcal{R}_j :

$$P(\mathbf{x} \in \mathcal{R}_j) = \int_{\mathcal{R}_j} p(\mathbf{u}) d\mathbf{u} \approx p(\mathbf{x}|\omega) \cdot V \text{ where } V \in \mathbb{R}^d$$

Formulation

- Assume a set of regions $\{\mathcal{R}_1, \dots, \mathcal{R}_N\}$ that divide the data space
- The probability of $\mathbf{x} \in \mathcal{D}_\omega$ being in region \mathcal{R}_j :

$$P(\mathbf{x} \in \mathcal{R}_j) = \int_{\mathcal{R}_j} p(\mathbf{u}) d\mathbf{u} \approx p(\mathbf{x}|\omega) \cdot V \text{ where } V \in \mathbb{R}^d$$

- Assuming regular sample distribution in data space:

$$P(\mathbf{x} \in \mathcal{R}_j) = \frac{k_j}{|\mathcal{D}|} \text{ where } k_j = |\{i : \mathbf{x}_i \in \mathcal{R}_j\}|$$

Formulation

- Assume a set of regions $\{\mathcal{R}_1, \dots, \mathcal{R}_N\}$ that divide the data space
- The probability of $\mathbf{x} \in \mathcal{D}_\omega$ being in region \mathcal{R}_j :

$$P(\mathbf{x} \in \mathcal{R}_j) = \int_{\mathcal{R}_j} p(\mathbf{u}) d\mathbf{u} \approx p(\mathbf{x}|\omega) \cdot V \text{ where } V \in \mathbb{R}^d$$

- Assuming regular sample distribution in data space:

$$P(\mathbf{x} \in \mathcal{R}_j) = \frac{k_j}{|\mathcal{D}|} \text{ where } k_j = |\{i : \mathbf{x}_i \in \mathcal{R}_j\}|$$

- The density function is approximated as:

$$\hat{p}(\mathbf{x}|\omega) = \frac{k_j}{|\mathcal{D}| \cdot V}$$

Weight feature example

Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$

Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$k_{[90,95]} = 23$

Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$$k_{[90,95]} = 23$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$

Weight feature example


- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$$k_{[90,95]} = 23$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$



Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$


Case #1

Case #2

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$k_{[90,95]} = 23$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$



Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1


Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$k_{[90,95]} = 23$

$$\hat{p}(\mathbf{x} = 91 | \omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$

Case #2

Volume range $[90, 92.5]\text{kg} \Rightarrow V = 2.5\text{kg}$



Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$


$$k_{[90,95]} = 23$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$

Case #2

Volume range $[90, 92.5]\text{kg} \Rightarrow V = 2.5\text{kg}$

$$k_{[90,92.5]} = 14$$



Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$$k_{[90,95]} = 23$$


$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$

Case #2

Volume range $[90, 92.5]\text{kg} \Rightarrow V = 2.5\text{kg}$

$$k_{[90,92.5]} = 14$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,92.5]}}{|\mathcal{D}| \cdot V} \approx 0.011$$



Weight feature example


- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$

Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$$k_{[90,95]} = 23$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$




Case #2

Volume range $[90, 92.5]\text{kg} \Rightarrow V = 2.5\text{kg}$

$$k_{[90,92.5]} = 14$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,92.5]}}{|\mathcal{D}| \cdot V} \approx 0.011$$



Weight feature example

- Unaccessible male **weight distribution**: $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
- Our **real scenario**:
 - Assortment $\mathcal{D}_{\text{male}}$ with $|\mathcal{D}_{\text{male}}| = 500$ samples
 - Approximate $p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male})$


Case #1

Volume range $[90, 95]\text{kg} \Rightarrow V = 5\text{kg}$

$$k_{[90,95]} = 23$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,95]}}{|\mathcal{D}| \cdot V} \approx 0.009$$

$$p(\mathbf{x} = 91 \text{ kg} | \omega = \text{male}) = 0.014$$




Case #2

Volume range $[90, 92.5]\text{kg} \Rightarrow V = 2.5\text{kg}$

$$k_{[90,92.5]} = 14$$

$$\hat{p}(\mathbf{x} = 91|\omega) = \frac{k_{[90,92.5]}}{|\mathcal{D}| \cdot V} \approx 0.011$$



Classification example

- Classification task: *male VS female*:

Classification example

- **Classification** task: *male VS female*:
 - $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
 - $p(\mathbf{x}|\omega = \text{female}) = \mathcal{N}(60, \sqrt{20})$

Classification example

- **Classification** task: *male* VS *female*:
 - $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
 - $p(\mathbf{x}|\omega = \text{female}) = \mathcal{N}(60, \sqrt{20})$
- What **we have**:
 - $\mathcal{D}_{\text{male}}$ assortment with $|\mathcal{D}_{\text{male}}| = 450$ samples
 - $\mathcal{D}_{\text{female}}$ assortment with $|\mathcal{D}_{\text{female}}| = 550$ samples

Classification example

- **Classification** task: *male* VS *female*:
 - $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
 - $p(\mathbf{x}|\omega = \text{female}) = \mathcal{N}(60, \sqrt{20})$
- What **we have**:
 - $\mathcal{D}_{\text{male}}$ assortment with $|\mathcal{D}_{\text{male}}| = 450$ samples
 - $\mathcal{D}_{\text{female}}$ assortment with $|\mathcal{D}_{\text{female}}| = 550$ samples
- What **we need**: Most likely **gender** when $\mathbf{x} = 72\text{kg}$

Classification example

- **Classification** task: *male VS female*:
 - $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
 - $p(\mathbf{x}|\omega = \text{female}) = \mathcal{N}(60, \sqrt{20})$
- What **we have**:
 - $\mathcal{D}_{\text{male}}$ assortment with $|\mathcal{D}_{\text{male}}| = 450$ samples
 - $\mathcal{D}_{\text{female}}$ assortment with $|\mathcal{D}_{\text{female}}| = 550$ samples
- What **we need**: Most likely **gender** when **$\mathbf{x} = 72\text{kg}$**

Male

Volume $V = 5\text{kg}$

$$k_{[70,75]} = 76$$

$$\hat{p}(72\text{ kg}|\text{male}) = 0.1689$$

$$P(\text{male}) = 450/450+550 = 0.45$$

Classification example

- **Classification** task: *male VS female*:
 - $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
 - $p(\mathbf{x}|\omega = \text{female}) = \mathcal{N}(60, \sqrt{20})$
- What **we have**:
 - $\mathcal{D}_{\text{male}}$ assortment with $|\mathcal{D}_{\text{male}}| = 450$ samples
 - $\mathcal{D}_{\text{female}}$ assortment with $|\mathcal{D}_{\text{female}}| = 550$ samples
- What **we need**: Most likely **gender** when **x = 72kg**

Male

Volume $V = 5\text{kg}$

$$k_{[70,75]} = 76$$

$$\hat{p}(72\text{ kg}|\text{male}) = 0.1689$$

$$P(\text{male}) = 450/450+550 = 0.45$$

Female

Volume $V = 5\text{kg}$

$$k_{[70,75]} = 3$$

$$\hat{p}(72\text{ kg}|\text{female}) = 0.001$$

$$P(\text{female}) = 550/450+550 = 0.55$$

Classification example

- **Classification** task: *male* VS *female*:
 - $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
 - $p(\mathbf{x}|\omega = \text{female}) = \mathcal{N}(60, \sqrt{20})$
- What **we have**:
 - $\mathcal{D}_{\text{male}}$ assortment with $|\mathcal{D}_{\text{male}}| = 450$ samples
 - $\mathcal{D}_{\text{female}}$ assortment with $|\mathcal{D}_{\text{female}}| = 550$ samples
- What **we need**: Most likely **gender** when **$\mathbf{x} = 72\text{kg}$**

Male

Volume $V = 5\text{kg}$

$k_{[70,75]} = 76$

$\hat{p}(72\text{ kg}|\text{male}) = 0.1689$

$P(\text{male}) = 450/450+550 = 0.45$

Female

Volume $V = 5\text{kg}$

$k_{[70,75]} = 3$

$\hat{p}(72\text{ kg}|\text{female}) = 0.001$

$P(\text{female}) = 550/450+550 = 0.55$

$$\hat{\omega} = \arg \max_{\omega \in \{\text{male}, \text{female}\}} \hat{p}(\mathbf{x} = 72|\omega) \cdot P(\omega)$$

Classification example

- **Classification** task: *male VS female*:
 - $p(\mathbf{x}|\omega = \text{male}) = \mathcal{N}(80, \sqrt{40})$
 - $p(\mathbf{x}|\omega = \text{female}) = \mathcal{N}(60, \sqrt{20})$
- What **we have**:
 - $\mathcal{D}_{\text{male}}$ assortment with $|\mathcal{D}_{\text{male}}| = 450$ samples
 - $\mathcal{D}_{\text{female}}$ assortment with $|\mathcal{D}_{\text{female}}| = 550$ samples
- What **we need**: Most likely **gender** when $\mathbf{x} = 72\text{kg}$

Male

Volume $V = 5\text{kg}$

$k_{[70,75]} = 76$

$\hat{p}(72\text{ kg}|\text{male}) = 0.1689$

$P(\text{male}) = 450/450+550 = 0.45$

Female

Volume $V = 5\text{kg}$

$k_{[70,75]} = 3$

$\hat{p}(72\text{ kg}|\text{female}) = 0.001$

$P(\text{female}) = 550/450+550 = 0.55$

$$\hat{\omega} = \arg \max_{\omega \in \{\text{male}, \text{female}\}} \hat{p}(\mathbf{x} = 72|\omega) \cdot P(\omega) \Rightarrow \text{male}$$

Limitations

- Does reducing the volume always improve the estimation?

Limitations

- Does **reducing the volume** always **improve** the estimation?
- Remarkably **senstitive** to **resolution**:
 - **Large** $V \Rightarrow$ oversmoothing / high bias
 - **Small** $V \Rightarrow$ noisy estimate / high variance

Limitations

- Does **reducing the volume** always **improve** the estimation?
- Remarkably **sensitive** to **resolution**:
 - **Large V** \Rightarrow oversmoothing / high bias
 - **Small V** \Rightarrow noisy estimate / high variance
- Also **discontinuous**, sensible to **bin alignment**, and **coarse**

Formulation

Parzen window estimator: Generalization of the histogram estimator

Formulation

Parzen window estimator: Generalization of the histogram estimator

- Query \mathbf{x} contributes locally to the density around its own position:

$$\hat{p}(\mathbf{x}|\omega) = \frac{k_j}{|\mathcal{D}| \cdot V} \Rightarrow \frac{k_j(\mathbf{x})}{|\mathcal{D}| \cdot V}$$

Formulation

Parzen window estimator: Generalization of the histogram estimator

- Query \mathbf{x} contributes locally to the density around its own position:

$$\hat{p}(\mathbf{x}|\omega) = \frac{k_j}{|\mathcal{D}| \cdot V} \Rightarrow \frac{k_j(\mathbf{x})}{|\mathcal{D}| \cdot V}$$


- Instead of rigid bins, **kernel function** around \mathbf{x} :

$$\hat{p}(\mathbf{x}|\omega) = \frac{1}{|\mathcal{D}| \cdot h^d} \sum_{i=1}^{|\mathcal{D}|} K\left(\frac{\mathbf{x} - \mathbf{x}_i}{h}\right)$$

$h \rightarrow$ window width

(FAA)

Window shapes



$$K_{\text{rec}}(u) = \begin{cases} 1/2 & |u| \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$K_{\text{epa}}(u) = \begin{cases} \frac{3}{4}(1 - u^2) & |u| \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$K_{\text{tri}}(u) = \begin{cases} 1 - |u| & |u| \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

$$K_{\text{gau}}(u) = \frac{1}{\sqrt{2\pi}} e^{-u^2/2}$$

Formulation

Issues

- **Main limitation:** fixed window width h regardless of x

Issues

- **Main limitation:** fixed window width h regardless of x
- Same neighborhood size regardless of (local) data density/sparsity:
 - High density \Rightarrow too many samples \Rightarrow over-smoothed estimation
 - Low density \Rightarrow too few samples \Rightarrow under-smoothed estimation

Formulation

k_n -**Nearest Neighbor estimator**: Addresses the limitations in [Parzen](#)

Formulation

k_n -Nearest Neighbor estimator: Addresses the limitations in Parzen

- ✗ Does not fix window width h
- ✓ Fixes number of samples k and lets the window volume vary

$$\hat{p}(\mathbf{x}|\omega) = \frac{k_j}{|\mathcal{D}| \cdot V} \Rightarrow \frac{k}{|\mathcal{D}| \cdot V(\mathbf{x})}$$

Formulation

Posterior probability

Posterior probability

- Assortment \mathcal{D} from an unknown distribution
 - Subset $\mathcal{D}_\omega = \{(\mathbf{x}_i, y_i) \in \mathcal{D} : y_i = \omega\}_{i=1}^{|\mathcal{D}|}$ with class ω

Posterior probability

- Assortment \mathcal{D} from an unknown distribution
 - Subset $\mathcal{D}_\omega = \{(\mathbf{x}_i, y_i) \in \mathcal{D} : y_i = \omega\}_{i=1}^{|\mathcal{D}|}$ with class ω
- Estimation using k neighbors
 - k_ω : number of neighbors with class ω

Posterior probability

- Assortment \mathcal{D} from an unknown distribution
 - Subset $\mathcal{D}_\omega = \{(\mathbf{x}_i, y_i) \in \mathcal{D} : y_i = \omega\}_{i=1}^{|\mathcal{D}|}$ with class ω
- Estimation using k neighbors
 - k_ω : number of neighbors with class ω

$$\hat{P}(\omega|\mathbf{x}) = \frac{\hat{p}(\mathbf{x}|\omega) \cdot P(\omega)}{\hat{p}(\mathbf{x})}$$

Posterior probability

- Assortment \mathcal{D} from an unknown distribution
 - Subset $\mathcal{D}_\omega = \{(\mathbf{x}_i, y_i) \in \mathcal{D} : y_i = \omega\}_{i=1}^{|\mathcal{D}|}$ with class ω
- Estimation using k neighbors
 - k_ω : number of neighbors with class ω

$$\hat{P}(\omega|\mathbf{x}) = \frac{\hat{p}(\mathbf{x}|\omega) \cdot P(\omega)}{\hat{p}(\mathbf{x})} \approx \frac{\frac{k_\omega/|\mathcal{D}_\omega|}{V(\mathbf{x})} \cdot \frac{|\mathcal{D}_\omega|}{|\mathcal{D}|}}{\frac{k/|\mathcal{D}|}{V(\mathbf{x})}}$$

Posterior probability

- Assortment \mathcal{D} from an unknown distribution
 - Subset $\mathcal{D}_\omega = \{(\mathbf{x}_i, y_i) \in \mathcal{D} : y_i = \omega\}_{i=1}^{|\mathcal{D}|}$ with class ω
- Estimation using k neighbors
 - k_ω : number of neighbors with class ω

$$\hat{P}(\omega|\mathbf{x}) = \frac{\hat{p}(\mathbf{x}|\omega) \cdot P(\omega)}{\hat{p}(\mathbf{x})} \approx \frac{\frac{k_\omega/|\mathcal{D}_\omega|}{V(\mathbf{x})} \cdot \frac{|\mathcal{D}_\omega|}{|\mathcal{D}|}}{\frac{k/|\mathcal{D}|}{V(\mathbf{x})}} = \frac{k_\omega}{k}$$

Issues

- Hyperparameter k remarkably influences the estimation:
 - $k \downarrow$: noisy estimate
 - $k \uparrow$: oversmoothed estimate
- Computational inefficiency
- Lack of smoothness / discontinuity
 - Piecewise estimate

Comparative summary of the density estimators

Estimator	Volume size	Number of points	Advantages	Disadvantages
Histogram	Fixed (bins)	Variable	Simple	Discontinuous
Parzen	Fixed (width h)	Variable	Smooth	Not adaptive
k_n -NN	Variable	Fixed	Adaptive	Not smooth, costly