Please check the examination details below before entering your candidate information				
Candidate surname			Other name	s
Pearson Edexcel Level 3 GCE	Centre	Number		Candidate Number
Thursday 6 June 2019				
Afternoon (Time: 1 hour 30 minute	es)	Paper Re	eference 9	FM0/02
Further Mathematics Advanced Paper 2: Core Pure Mathematics 2				
You must have: Mathematical Formulae and Statistical Tables (Green), calculator Total Marks				

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear.
 Answers without working may not gain full credit.
- Answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶





Answer ALL questions. Write your answers in the spaces provided.

1. (a) Prove that

$$\tanh^{-1}(x) = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right) \qquad -k < x < k$$

stating the value of the constant k.

(5)

(b) Hence, or otherwise, solve the equation

$$2x = \tanh\left(\ln\sqrt{2 - 3x}\right)$$

(5)

I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	
I	

Question 1 continued	
(Total fo	r Question 1 is 10 marks)



2. The roots of the equation

$$x^3 - 2x^2 + 4x - 5 = 0$$

are p, q and r.

Without solving the equation, find the value of

(i)
$$\frac{2}{p} + \frac{2}{q} + \frac{2}{r}$$

(ii)
$$(p-4)(q-4)(r-4)$$

(iii)
$$p^3 + q^3 + r^3$$

(8)

Question 2 continued				



Question 2 continued

Question 2 continued	
	Tradition Order 2 2 9 1
	Total for Question 2 is 8 marks)



3.

$$f(x) = \frac{1}{\sqrt{4x^2 + 9}}$$

(a) Using a substitution, that should be stated clearly, show that

$$\int f(x)dx = A \sinh^{-1}(Bx) + c$$

where c is an arbitrary constant and A and B are constants to be found.

(4)

(b) Hence find, in exact form in terms of natural logarithms, the mean value of f(x) over the interval [0, 3].

1	2)
Ų.	4)

(Total for Ouestion 3 is 6 marks)	Question 3 continued
(Total for Ouestion 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Ouestion 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
(Total for Question 3 is 6 marks)	
	(Total for Question 3 is 6 marks)



4. The infinite series C and S are defined by

$$C = \cos\theta + \frac{1}{2}\cos 5\theta + \frac{1}{4}\cos 9\theta + \frac{1}{8}\cos 13\theta + \dots$$

$$S = \sin\theta + \frac{1}{2}\sin 5\theta + \frac{1}{4}\sin 9\theta + \frac{1}{8}\sin 13\theta + \dots$$

Given that the series C and S are both convergent,

(a) show that

$$C + iS = \frac{2e^{i\theta}}{2 - e^{4i\theta}}$$
 (4)

(b) Hence show that

$$S = \frac{4\sin\theta + 2\sin 3\theta}{5 - 4\cos 4\theta}$$

(4)



Question 4 continued	
	(Total for Question 4 is 8 marks)



5. An engineer is investigating the motion of a sprung diving board at a swimming pool. Let *E* be the position of the end of the diving board when it is at rest in its equilibrium position and when there is no diver standing on the diving board.

A diver jumps from the diving board.

The vertical displacement, h cm, of the end of the diving board above E is modelled by the differential equation

$$4\frac{d^2h}{dt^2} + 4\frac{dh}{dt} + 37h = 0$$

where *t* seconds is the time after the diver jumps.

(a) Find a general solution of the differential equation.

(2)

When t = 0, the end of the diving board is 20 cm below E and is moving upwards with a speed of 55 cm s⁻¹.

(b) Find, according to the model, the maximum vertical displacement of the end of the diving board above E.

(8)

(c) Comment on the suitability of the model for large values of t.

(2)

Question 5 continued				



Question 5 continued

14

v			
Χ.			
	Ē	m	
\sim 1		ж	
	N A	9	
: 24		61	ä
×Ι	м	m	
$^{\sim}$	â	a	ú
XI	м	z	
:)			
		×	á
V/I	ø		۰
	9		'n
		X	۰
\sim	ľ٧	я	h
V		ĸ.	
×Ι		-	•
×	\rightarrow	_	2
X.	~	•	7
v.	sal	L	_
O.	_	=	9
ΧI	K	×	
O	7		,
XJ	ĸ.		
V	~		
$^{\prime}$	_		
	Sá	æ	
: 24			
X.	X	Z	Ē
1	_	•	Ì
×ι	Kı	iΧ	
()	ы	ш	
Х.	_	_	7
V1	L.	S	
\sim	,	•	ņ
×Ι	ĸ.		
(N			
Χ.	ă.	Ä	á
V	2	r	•
C)			
×ι	×		
	1		
\sim 1		z	
	\sim		
	-	_	
V1	Ľ		
\sim			
V	ĸ		
			b
ΧI	K		
1	•	ш	,
$^{\sim}$		z	s
V	~		
		~	
ΧI	×	ø	þ
			2
	24	ŝ	s
X	ĸ.	×	
(X	ĸ.	G	
	74	ø	۲
	a	ь	ć
M	\mathbb{P}	S	
X	2	2	
X	D X	X	

Question 5 continued	
(Total for Question 5 is 12 marks)	



6.	In an Argand diagram, the points A , B and C are the vertices of an equilateral triangle with its centre at the origin. The point A represents the complex number $6 + 2i$.			
(a) Find the complex numbers represented by the points B and C , giving your answers in the form $x + iy$, where x and y are real and exact.				
		(6)		
	The points D , E and F are the midpoints of the sides of triangle ABC .			
	(b) Find the exact area of triangle <i>DEF</i> .			
		(3)		

Question 6 continued



Question 6 continued

Question 6 continued	
	(Total for Question 6 is 9 marks)



7.

$$\mathbf{M} = \begin{pmatrix} 2 & -1 & 1 \\ 3 & k & 4 \\ 3 & 2 & -1 \end{pmatrix} \quad \text{where } k \text{ is a constant}$$

(a) Find the values of k for which the matrix M has an inverse.

(2)

(b) Find, in terms of p, the coordinates of the point where the following planes intersect

$$2x - y + z = p$$

$$3x - 6y + 4z = 1$$

$$3x + 2y - z = 0$$

(5)

(c) (i) Find the value of q for which the set of simultaneous equations

$$2x - y + z = 1$$

$$3x - 5y + 4z = q$$

$$3x + 2y - z = 0$$

can be solved.

(ii) For this value of q, interpret the solution of the set of simultaneous equations geometrically.

(4)

			į	

Question 7 continued




Question 7 continued

Question 7 continued
(Total for Question 7 is 11 marks)
1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2



8.



e 1 Figure 2

Figure 1 shows the central vertical cross section *ABCD* of a paddling pool that has a circular horizontal cross section. Measurements of the diameters of the top and bottom of the paddling pool have been taken in order to estimate the volume of water that the paddling pool can contain.

Using these measurements, the curve BD is modelled by the equation

$$y = \ln(3.6x - k)$$
 $1 \le x \le 1.18$

as shown in Figure 2.

(a) Find the value of k.

(1)

(b) Find the depth of the paddling pool according to this model.

(2)

The pool is being filled with water from a tap.

(c) Find, in terms of h, the volume of water in the pool when the pool is filled to a depth of h m.

(5)

Given that the pool is being filled at a constant rate of 15 litres every minute,

(d) find, in cm h⁻¹, the rate at which the water level is rising in the pool when the depth of the water is 0.2 m.

(3)

Question 8 continued



Question 8 continued		

Question 8 continued		



Question 8 continued	
	(Total for Question 8 is 11 marks)
	TOTAL FOR PAPER IS 75 MARKS
	TO THE FOR THE ENTRY TO THINK