

AQUÍ VA EL TÍTULO DEL ARTÍCULO

ABSTRACT. Aquí va el resumen

CONTENTS

1. Introduction	1
2. Background	1
3. Compact quotients	1
4. Tidy frames	6
5. Stacked and strongly stacked frames	10
6. Separation properties in frames	13
7. Quotients of stacked frames	18
8. Relationship to the property (H)	21
9. Marcos KC	21
10. Examples	24
10.1. The frame $[0, 1]$	24
References	27

1. INTRODUCTION

[PP21, SS06a, Hoc69]

2. BACKGROUND

[Sim06a, Sex03, Joh82]

3. COMPACT QUOTIENTS

Let $f: A \rightarrow B$ be a frame morphism with right adjoint $f_*: B \rightarrow A$. If $F \subseteq A$ and $G \subseteq B$ are filters, we define:

$$(1) \quad b \in f^*F \iff f_*(b) \in F \quad a \in f_*G \iff f(a) \in G,$$

where f^*F and f_*G are filters in B and A , respectively.

Proposition 3.1. *For $f = f^*: A \rightarrow B$ a frame morphism and $G \in B^\wedge$, then $f_*G \in A^\wedge$.*

Proof. By (1), f_*G is a filter in A . We need that f_*G satisfy the open filter condition. Let $X \subseteq A$ be such that $\bigvee X \in f_*G$, with X directed. Then

$$Y = \{f(x) \mid x \in X\}$$

is directed and $f(\bigvee X) = \bigvee f[X] = \bigvee Y \in G$. Since G is an open filter and $f(\bigvee X) = \bigvee f[X] \in G$, there exists $x \in X$ such that $f(x) \in G$. Therefore, $x \in f_*G$. $\square$

Simmons shows in [Sim04, Lemma 8.9 and Corollary 8.10] that the following diagram

$$(2) \quad \begin{array}{ccc} A & \xrightarrow{f^\infty} & A \\ U_A \downarrow & & \downarrow U_A \\ \mathcal{O}S & \xrightarrow{F^\infty} & \mathcal{O}S \end{array}$$

commutes laxly, that is, $U_A \circ f^\infty \leq F^\infty \circ U_A$. In this diagram f^∞ and F^∞ represent the fitted nuclei associated with the filters $F \in A^\wedge$ and $\nabla \in \mathcal{O}S^\wedge$, respectively, that is,

$$f^\infty = v_F \quad \text{and} \quad F^\infty = v_\nabla.$$

Here we prove something more general, since we consider $j \in NA$, $F \in A_j^\wedge$ and $j_*F \in A^\wedge$ to produce the diagram

$$\begin{array}{ccc} A & \xrightarrow{f^\infty} & A \\ j^* \downarrow & & \downarrow j^* \\ A_j & \xrightarrow{\hat{f}^\infty} & A_j \end{array}$$

where $\hat{f}^\infty$ and f^∞ are the fitted nuclei associated with the filters j_*F and F , respectively.

The following lemma is part of the analysis developed in this work and constitutes a key tool in the construction of the commutative diagrams that follow.

Lemma 3.2. *For $j = j^*$, f and $\hat{f}$ as before, it holds that $j \circ \hat{f} \leq f \circ j$.*

Proof. Recall that, by (1)

$$\hat{f} = \bigvee \{v_y \mid y \in j_*F\} \quad \text{y} \quad f = \bigvee \{v_{j(y)} \mid j(y) \in F\}.$$

Let $a \in A$ be. For every $y \in j_*F$, we have

$$j(v_y(a)) = j(y \succ a) \leq (j(y) \succ j(a)) = v_{j(y)}(j(a)) \leq f(j(a)),$$

where the last inequality follows from $j(y) \in F$. Therefore, since j preserves arbitrary joins,

$$j(\hat{f}(a)) = j\left(\bigvee \{v_y(a) \mid y \in j_*F\}\right) = \bigvee \{j(v_y(a)) \mid y \in j_*F\} \leq f(j(a)).$$

Hence, $j \circ \hat{f} \leq f \circ j$. $\square$

We now prove the above for an arbitrary ordinal.

Corollary 3.3. *For j , f and $\hat{f}$ as before, it holds that $j \circ \hat{f}^\alpha \leq f^\alpha \circ j$ where $\alpha \in \text{Ord}$.*

Proof. We proceed by transfinite induction on α . If $\alpha = 0$, the result is immediate.

For the induction step, suppose that the inequality holds for α . Then

$$j \circ \hat{f}^{\alpha+1} = j \circ \hat{f} \circ \hat{f}^\alpha \leq f \circ j \circ \hat{f}^\alpha \leq f \circ f^\alpha \circ j = f^{\alpha+1} \circ j,$$

where the first inequality follows from Lemma 3.2 and the second from the induction hypothesis.

Finally, let λ be a limit ordinal. For every $a \in A$,

$$j(\hat{f}(a)) = j\left(\bigvee_{\alpha < \lambda} \hat{f}^\alpha\right) = \bigvee_{\alpha < \lambda} j(\hat{f}^\alpha(a)) \leq \bigvee_{\alpha < \lambda} f^\alpha(j(a)) = f^\lambda(j(a)).$$

where the inequality follows from the induction hypothesis. Hence

$$j \circ \hat{f}^\lambda \leq f^\lambda \circ j.$$

This completes the proof. $\square$

Recalling that $v_F = f^\infty$ and $v_{j_*F} = \hat{f}^\infty$, Corollary 3.3 yields

$$j \circ v_{j_*F} \leq v_F \circ j$$

We now compare the compact quotients induced by F and j_*F , and construct a frame morphism between them making the corresponding diagram commute.

If we consider j , v_F and v_{j_*F} we have the following diagram

$$(3) \quad \begin{array}{ccc} A & \xrightleftharpoons[(v_{j_*F})_*]{(v_{j_*F})^*} & A_{j_*F} \\ \downarrow j & \searrow H & \\ A_j & \xrightleftharpoons[(v_F)^*]{(v_F)_*} & A_F \end{array}$$

where A_F and A_{j_*F} are the compact quotients produced by v_F and v_{j_*F} , respectively.

Define

$$H: A \rightarrow A_F, \quad H = v_F \circ j.$$

Since v_F is idempotent, $H(A) \subseteq A_F$. Hence, by restricting H to A_{j_*F} , we obtain a map

$$h: A_{j_*F} \rightarrow A_F, \quad h = H|_{A_{j_*F}}.$$

As finite meets in quotient frames are inherited from the ambient frame, h preserves finite meets. It remains to verify that h preserves arbitrary joins.

Lemma 3.4. *The map $h: A_{j_*F} \rightarrow A_F$ preserves arbitrary joins.*

Proof. Let $X \subseteq A_{j_*F}$. Since joins in A_{j_*F} are given by

$$\bigvee_{v_{j_*F}} X = v_{j_*F}(\bigvee X)$$

we have

$$\begin{aligned} h\left(\bigvee_{v_{j_*F}} X\right) &= v_F\left(j\left(v_{j_*F}\left(\bigvee X\right)\right)\right) \\ &\leq v_F\left(v_F\left(j\left(\bigvee X\right)\right)\right) \\ &= v_F\left(j\left(\bigvee X\right)\right), \end{aligned}$$

where the inequality follows from Corollary 3.3 and the equality from the idempotency of v_F .

Since H is a frame morphism,

$$v_F\left(j\left(\bigvee X\right)\right) = \bigvee_{v_F}\{H(x) \mid x \in X\} = \bigvee_{v_F}\{h(x) \mid x \in X\}.$$

Therefore,

$$h\left(\bigvee_{v_{j_*F}} X\right) \leq \bigvee_{v_F} h[X].$$

Conversely, for every $x \in X$,

$$x \leq \bigvee_{v_{j_*F}} X,$$

and hence, by monotonicity of h ,

$$h(x) \leq h\left(\bigvee_{v_{j_*F}} X\right).$$

Since this holds for every $x \in X$, we have

$$\bigvee_{v_F} h[X] \leq h\left(\bigvee_{v_{j_*F}} X\right).$$

Thus,

$$h\left(\bigvee_{v_{j_*F}} X\right) = \bigvee_{v_F} h[X].$$

Therefore h preserves arbitrary joins. $\square$

The previous construction yields the following commutative diagram induced by the compact quotients associated with the fitted nuclei.

Proposition 3.5. *The diagram*

$$(4) \quad \begin{array}{ccc} A & \xrightarrow{v_{j^*F}} & A_{j^*F} \\ j \downarrow & & \downarrow h \\ A_j & \xrightarrow{v_F} & A_F \end{array}$$

is commutative.

Proof. By Lemma 3.4, h is a frame morphism. The commutativity of the diagram follows from the definition of h . $\square$

A particular instance of diagram (4) is obtained by taking $j^* = \mathcal{U}_A$, where $S = \text{pt } A$. Thus, we obtain the following commutative diagram:

$$\begin{array}{ccc} A & \xrightarrow{v_F} & A_F \\ \mathcal{U}_A \downarrow & & \downarrow h \\ \mathcal{O}S & \xrightarrow{v_\nabla} & \mathcal{O}S_\nabla \end{array}$$

Recall that $\text{pt } A_F = Q$. Hence, the spatial reflection of A_F is given by

$$\mathcal{U}_{A_F}: A_F \rightarrow \mathcal{O} \text{pt } A_F = \mathcal{O}Q.$$

Similarly, since $\text{pt}((\mathcal{O}S)_\nabla) = Q$, we have the spatial reflection

$$\mathcal{U}_{\mathcal{O}S_\nabla}: \mathcal{O}S_\nabla \rightarrow \mathcal{O} \text{pt}(\mathcal{O}S)_\nabla = \mathcal{O}Q.$$

Consequently, the two induced morphisms from A to $\mathcal{O}Q$ are both given by

$$x \mapsto \mathcal{U}_A(x) \cap Q.$$

Therefore, the preceding constructions fit into the following commutative diagram:

$$(5) \quad \begin{array}{ccccc} A & \xrightarrow{v_F} & A_F & & \\ \mathcal{U}_A \downarrow & & \downarrow h & \searrow \mathcal{U}_{A_F} & \\ & & & & \mathcal{O}Q \\ \mathcal{O}S & \xrightarrow{v_\nabla} & \mathcal{O}S_\nabla & \nearrow \mathcal{U}_{\mathcal{O}S_\nabla} & \end{array}$$

Since the above diagram is determined by $F \in A^\wedge$ and the associated compact saturated subset $Q \in \mathcal{Q}S$, we refer to diagram (5) as the Q -square.

Theorem 3.6. *Under the hypotheses of the Q -square, if A satisfies property (H), then*

$$\mathcal{O}Q \cong \uparrow Q'.$$

Equivalently, the frame of open sets of the point space of A_F is isomorphic to a compact closed quotient of a Hausdorff space.

Proof. Suppose that A satisfies **(H)**. Since **(H)** is conservative, $S = \text{pt } A$ is a T_2 space. Hence, for every $Q \in \mathcal{Q}S$, the set Q is closed, and therefore $Q' \in \mathcal{O}S$

Moreover, since every T_2 space is strongly stacked, we have

$$v_{\nabla} = [Q'].$$

As $Q' \in \mathcal{O}S$, it follows that

$$[Q'] = u_{Q'}$$

Consequently,

$$(\mathcal{O}S)_{v_{\nabla}} = (\mathcal{O}S)_{u_{Q'}} \cong \uparrow Q'.$$

Now, $u_{Q'}$ is a complemented nucleus in $\mathcal{O}S$, by [PP12, Proposition VI.3.3], implies that $\mathcal{O}S_{v_{\nabla}}$ is spatial. Therefore,

$$(\mathcal{O}S)_{v_{\nabla}} \cong \mathcal{O} \text{pt}((\mathcal{O}S)_{v_{\nabla}}).$$

Since

$$\text{pt}((\mathcal{O}S)_{v_{\nabla}}) = Q,$$

we obtain

$$\mathcal{O}Q \cong (\mathcal{O}S)_{v_{\nabla}} \cong \uparrow Q'.$$

□

4. TIDY FRAMES

Proposition 4.1. *If A is a tidy frame, then A_j is a tidy frame.*

Proof. Let A be α -tidy and let $F \in A^{\wedge}$. By Proposition 3.1, $j_*F \in A^{\wedge}$. Let

$$\hat{f} = \bigvee_{y \in j_*F} v_y \quad \text{and} \quad f = \bigvee_{x \in F} v_x.$$

be the prenuclei associated with j_*F and F , respectively.

Since A is tidy, there exists an ordinal α such that, for every $y \in j_*F$,

$$y \vee \hat{d} = 1, \quad \text{with } \hat{d} = \hat{f}^{\alpha}(0).$$

Let

$$d = f^{\alpha}(j(0)),$$

by Corollary 3.3,

$$j(\hat{d}) = j(\hat{f}^{\alpha}(0)) \leq f^{\alpha}(j(0)) = d.$$

Since j is inflationary, $\hat{d} \leq j(\hat{d})$, and hence

$$\hat{d} \leq d.$$

Now, let $x \in F$. Thus, $x \in A_j$, we have $j(x) = x$, and therefore $x \in j_*F$. Then,

$$x \vee \hat{d} = 1.$$

As $\hat{d} \leq d$, it follows that

$$1 = x \vee \hat{d} \leq x \vee d \leq 1.$$

Hence $x \vee d = 1$. Therefore, A_j is α -tidy. $\square$

The following theorem establishes that tidiness is preserved under binary coproducts.

Proposition 4.2. *Let A_1 and A_2 be α -tidy frames. Then $A_1 \oplus A_2$ is α -tidy.*

Proof. By construction of the coproduct $A_1 \oplus A_2$, consider the canonical frame morphism

$$\iota_i: A_i \rightarrow A_1 \oplus A_2, \quad i = 1, 2$$

together with their right adjoints $(\iota_i)_*: A_1 \oplus A_2 \rightarrow A_i$.

Then, we have the following diagram

$$\begin{array}{ccccccc} & & & \iota_i & & & \\ & & & \curvearrowright & & & \\ A_i & \xrightarrow{\alpha_i} & A_1 \times A_2 & \xrightarrow{\downarrow(-)} & \mathcal{L}(A_1 \times A_2) & \xrightarrow{\sim} & A_1 \oplus A_2 \\ & & & \curvearrowleft & & & \\ & & & (\iota_i)_* & & & \end{array}$$

By Proposition 3.1,

$$(\iota_i)_* F \in A_i^\wedge.$$

Since A_i is α -tidy, for every $x_i \in (\iota_i)_* F$ there exists $d_i \in A_i$ such that

$$x_i \vee d_i = 1,$$

where

$$d_i = f_i^\alpha(0) \quad \text{and} \quad f_i = \bigvee_{y \in (\iota_i)_* F} v_y.$$

By the definition of $(\iota_i)_* F$,

$$x_i \in (\iota_i)_* F \iff \iota_i(x_i) \in F.$$

Thus,

$$x_1 \oplus 1, 1 \oplus x_2 \in F.$$

Since F is a filter,

$$(x_1 \oplus 1) \wedge (1 \oplus x_2) = x_1 \oplus x_2 \in F.$$

Now let

$$f_F = \bigvee_{z \in F} v_z$$

be the prenucleus associated with F . We first show that, for $i = 1, 2$,

$$\iota_i \circ f_i \leq f_F \circ \iota_i.$$

Indeed, let $a \in A_i$ and $y \in (\iota_i)_* F$. Since $\iota_i(y) \in F$,

$$\begin{aligned} \iota_i(v_y(a)) &= \iota_i(y \succ a) \\ &\leq (\iota_i(y) \succ \iota_i(a)) \\ &= v_{\iota_i(y)}(\iota_i(a)) \\ &\leq f_F(\iota_i(a)). \end{aligned}$$

Therefore, since the join defining f_i is pointwise and ι_i preserves arbitrary joins,

$$\iota_i(f_i(a)) = \iota_i\left(\bigvee_{y \in (\iota_i)_* F} v_y(a)\right) = \bigvee_{y \in (\iota_i)_* F} \iota_i(v_y(a)) \leq f_F(\iota_i(a)).$$

Hence,

$$\iota_i \circ f_i \leq f_F \circ \iota_i.$$

We now prove by transfinite induction that

$$\iota_i \circ f_i^\beta \leq f_F^\beta \circ \iota_i$$

for every ordinal β .

If $\beta = 0$, the result is immediate. Suppose that it holds for β . Then

$$\begin{aligned} \iota_i \circ f_i^{\beta+1} &= \iota_i \circ f_i \circ f_i^\beta \\ &\leq f_F \circ \iota_i \circ f_i^\beta \\ &\leq f_F \circ f_F^\beta \circ \iota_i \\ &= f_F^{\beta+1} \circ \iota_i. \end{aligned}$$

Now let λ be a limit ordinal. For every $a \in A_i$,

$$\begin{aligned} \iota_i(f_i^\lambda(a)) &= \iota_i\left(\bigvee_{\beta < \lambda} f_i^\beta(a)\right) \\ &= \bigvee_{\beta < \lambda} \iota_i(f_i^\beta(a)) \\ &\leq \bigvee_{\beta < \lambda} f_F^\beta(\iota_i(a)) \\ &= f_F^\lambda(\iota_i(a)). \end{aligned}$$

Thus,

$$\iota_i \circ f_i^\beta \leq f_F^\beta \circ \iota_i$$

for every ordinal β .

Evaluating at $0 \in A_i$ for $\beta = \alpha$, we obtain

$$\iota_1(d_1) \leq f_F^\alpha(0 \oplus 1) \quad \text{and} \quad \iota_2(d_2) \leq f_F^\alpha(1 \oplus 0).$$

Therefore,

$$\begin{aligned} d_1 \oplus d_2 &= (d_1 \oplus 1) \wedge (1 \oplus d_2) \\ &\leq f_F^\alpha(0 \oplus 1) \wedge f_F^\alpha(1 \oplus 0) \\ &= f_F^\alpha(0 \oplus 1 \wedge 1 \oplus 0). \end{aligned}$$

Set $d = f_F^\alpha(0 \oplus 0)$, then $d_1 \oplus d_2 \leq d$. Finally,

$$\begin{aligned} 1 \oplus 1 &= (x_1 \vee d_1) \oplus (x_2 \vee d_2) \\ &= (x_1 \oplus x_2) \vee (d_1 \oplus d_2) \\ &\leq (x_1 \oplus x_2) \vee d \\ &\leq 1 \oplus 1. \end{aligned}$$

Therefore,

$$(x_q \oplus x_2) \vee d = 1 \oplus 1$$

and $A_1 \oplus A_2$ is α -tidy. $\square$

Corollary 4.3. *Let A_1 and A_2 be α -tidy and β -tidy, respectively. Then $A_1 \oplus A_2$ is γ -tidy, where*

$$\gamma = \max\{\alpha, \beta\}.$$

Proof. Let $\gamma = \max\{\alpha, \beta\}$. Then A_1 and A_2 are both γ -tidy. Hence, by Proposition 4.2, $A_1 \oplus A_2$ is γ -tidy. $\square$

We extend the previous result to arbitrary coproducts.

Proposition 4.4. *Tidiness is preserved under arbitrary coproducts.*

Proof. Let $\{A_i\}_{i \in I}$ be a family of tidy frames, and for each $i \in I$, let α_i be a tidiness degree of A_i . Consider

$$A = \bigoplus_{i \in I} A_i$$

and let

$$\iota_i: A_i \rightarrow A$$

denote the canonical coproduct morphisms. Since ι_i is a frame morphism, it has a right adjoint $(\iota_i)_*$.

Let $F \in A^\wedge$. By Proposition 3.1,

$$(\iota_i)_* F \in A_i^\wedge$$

for every $i \in I$. Moreover,

$$x_i \in (\iota_i)_* F \iff \iota_i(x_i) \in F.$$

Set $\alpha = \sup\{\alpha_i \mid i \in I\}$. Then every A_i is α -tidy. Hence, for every $x_i \in (\iota_i)_* F$,

$$x_i \vee d_i = 1,$$

where

$$d_i = f_i^\alpha(0) \quad \text{and} \quad f_i = \bigvee_{y \in (\iota_i)_* F} v_y.$$

Let

$$f_F = \bigvee_x \cdot \in Fv_x$$

be the prenucleus associated with F . As in the proof of Proposition 4.2, for every $i \in I$ we have

$$\iota_i \circ f_i \leq f_F \circ \iota_i,$$

and, by transfinite induction,

$$\iota_i \circ f_i^\beta \leq f_F^\beta \circ \iota_i$$

for every ordinal β .

Evaluating at $0 \in A_i$ for $\beta = \alpha$, we obtain

$$\iota_i(d_i) = \iota_i(f_i^\alpha(0)) \leq f_F^\alpha(\iota_i(0)) = f_F^\alpha(\langle 0 \rangle).$$

Set

$$d = f_F^\alpha(\langle 0 \rangle).$$

Therefore,

$$\iota_i(d_i) \leq d$$

for every $i \in I$. Taking arbitrary joins yields

$$\langle d_i \rangle \bigvee_{i \in I} \iota_i(d_i) \leq d.$$

Now let

$$\langle x_i \rangle \in F.$$

Since $x_i \vee d_i = 1$ for every $i \in I$, we have

$$\langle x_i \rangle \vee \langle d_i \rangle = \langle x_i \vee d_i \rangle = \langle 1 \rangle.$$

As $\langle d_i \rangle \leq d$,

$$\langle 1 \rangle = \langle x_i \rangle \vee \langle d_i \rangle \leq \langle x_i \rangle \vee d \leq \langle 1 \rangle.$$

Thus,

$$\langle x_i \rangle \vee d = \langle 1 \rangle.$$

Therefore, $A = \bigoplus_{i \in I} A_i$ is α -tidy, and hence tidy. $\square$

Corollary 4.5. *The subcategory of tidy frames is a coreflective subcategory of $\mathbf{Frm}$.*

5. STACKED AND STRONGLY STACKED FRAMES

Lemma 5.1. *If $\alpha \in \mathbf{Ord}$ and $X \in CS$, then*

$$(6) \quad \partial_F^\alpha(X) = (f_F^\alpha(X'))'.$$

Proof. By transfinite induction on $\alpha \in \mathbf{Ord}$:

- If $\alpha = 0$, then $\partial_F^0 = \text{id}$, that is, $\partial_F^0(X) = X$ and $f_F^0 = \text{id}$. Thus

$$\partial_F^0(X) = X = (X'') = (f_F^0(X'))'.$$

- Assume that $\partial_F^\alpha(X) = (f_F^\alpha(X'))'$ holds. By definition,

$$\partial_F^{\alpha+1}(X) = \partial_F(\partial_F^\alpha(X)).$$

Furthermore,

$$\partial_F^{\alpha+1}(X) = (f_F((\partial_F^\alpha(X))'))'.$$

By induction hypothesis, $\partial_F^\alpha(X) = (f_F^\alpha(X'))'$ and substituting we obtain

$$\partial_F^{\alpha+1}(X) = (f_F(f_F^\alpha(X')))' = (f_F^{\alpha+1}(X'))'$$

which is what we wanted.

- Let λ be a limit ordinal and assume that the statement holds for all $\alpha < \lambda$. By definition,

$$\partial_F^\lambda(X) = \bigcap_{\alpha < \lambda} \partial_F^\alpha(X).$$

By De Morgan laws, we have that

$$(\partial_F^\lambda(X))' = \left(\bigcap_{\alpha < \lambda} \partial_F^\alpha(X) \right)' = \bigcup_{\alpha < \lambda} (\partial_F^\alpha(X))'.$$

Using the induction hypothesis, for all $\alpha < \lambda$,

$$(\partial_F^\alpha(X))' = f_F^\alpha(X'),$$

Then,

$$(\partial_F^\lambda(X))' = \bigcup_{\alpha < \lambda} f_F^\alpha(X') = f_F^\lambda(X').$$

Therefore,

$$(\partial_F^\lambda(X))' = f_F^\lambda(X'),$$

that is,

$$\partial_F^\lambda(X) = (f_F^\lambda(X'))'.$$

□

As a particular case, we have the following.

Corollary 5.2. *If $X \in \mathcal{CS}$, $U \in \mathcal{OS}$ and $\alpha = \infty$, then*

$$(7) \quad \begin{aligned} \partial_F^\infty(X) = (v_F(X'))' &\iff \partial_F^\infty(U')' = (v_F(U)) \\ &\iff \partial_F^\infty(X)' = (v_F(X')). \end{aligned}$$

Lemma 5.3. *If $A = \mathcal{OS}$ and $E \subseteq S$, then*

$$\mathcal{U}_*[E]\mathcal{U} = [E].$$

Proof. Since A is spacial, then $\mathcal{U}$ is an isomorphism. Thus, if $U \in \mathcal{O} \text{ pt } A$ exists $V \in \text{pt } A$ such that $U = \mathcal{U}(V)$ and as $U \in \text{pt } A$, then $U = V$. Next,

$$\begin{aligned} (\mathcal{U}_*[E]\mathcal{U})(U) &= \mathcal{U}_*[E](\mathcal{U}(U)) = \mathcal{U}_*([E](U)) \\ &= \mathcal{U}_*(E \cup U)^\circ = \bigcup \{W \in \mathcal{OS} \mid W \subseteq (E \cup U)^\circ\} \\ &= (E \cup U)^\circ = [E](U). \end{aligned}$$

Therefore $\mathcal{U}_*[E]\mathcal{U} = [E]$.

□

Proposition 5.4. . For $F \in A^\wedge$, $Q \in \mathcal{QS}$ and $\nabla = \nabla(Q)$, if $k \in [v_\nabla, w_\nabla]$, then $\mathcal{U}_* \circ k \circ \mathcal{U} \in [v_F, w_F]$.

Proof. If $k \in [v_\nabla, w_\nabla]$, then $\mathcal{U}_* \circ k \circ \mathcal{U} \in NA$. Additionally, for $F \in A^\wedge$ if $j \in [v_F, w_F]$, then $F = \nabla(j)$. In this way we must prove that $F = \nabla(\mathcal{U}_* \circ k \circ \mathcal{U})$.

By Hofmann-Mislove Theorem, we have that

$$x \in F \iff Q \subseteq \mathcal{U}(x) \iff \mathcal{U}(x) \in \nabla = \nabla(Q).$$

Furthermore, for all $x \in A$

$$(\mathcal{U}_* \circ k \circ \mathcal{U})(x) = \mathcal{U}_*(k(\mathcal{U}(x))) = \bigwedge (S \setminus k(\mathcal{U}(x))).$$

Since $k \in [v_\nabla, w_\nabla]$, then $\nabla = \nabla(k)$, that is, for $V \in \nabla$ it holds that $k(V) = S$. Therefore,

$$\begin{aligned} x \in F &\iff \mathcal{U}(x) \in \nabla \iff S \setminus k(\mathcal{U}(x)) = \emptyset \\ &\iff \bigwedge (S \setminus k(\mathcal{U}(x))) = 1 \\ &\iff x \in \nabla(\mathcal{U}_* \circ k \circ \mathcal{U}) \end{aligned}$$

□

We have a morphism $\mathcal{U}: [v_\nabla, w_\nabla] \rightarrow [v_F, w_F]$. Note that this occurs at both ends of the interval on NA .

Take $k \in [v_\nabla, w_\nabla]$. Since v_F is a fitted nuclei, then it holds

$$(8) \quad v_F \leq \mathcal{U}_* \circ k \circ \mathcal{U}.$$

Similarly, as w_F is the largest element in its block,

$$(9) \quad \mathcal{U}_* \circ k \circ \mathcal{U} \leq w_F.$$

Proposition 5.5. If $F \in A^\wedge$ and $\nabla \in \mathcal{OS}^\wedge$, then

$$w_F = \mathcal{U}_* \circ w_\nabla \circ \mathcal{U}.$$

Proof. By (9), we must verify that $w_F \leq \mathcal{U}_* \circ w_\nabla \circ \mathcal{U}$. Equivalently, we must prove that

$$\mathcal{U} \circ w_F \leq w_\nabla \circ \mathcal{U},$$

that is, for all $x \in A$

$$\mathcal{U}(w_F(x)) \subseteq [M'](\mathcal{U}(x)) = (M' \cup \mathcal{U}(x))^\circ.$$

Since $\mathcal{U}(w_F(x)) \in \mathcal{OS}$, it is enough that $\mathcal{U}(w_F(x)) \subseteq M' \cup \mathcal{U}(x)$. As a contradiction, let $y \in \mathcal{U}(w_F(x))$ be such that $y \notin M'$ and $y \notin \mathcal{U}(x)$. Then

$$y \in M \quad \text{and} \quad x \leq y.$$

By hypothesis, $y \in \mathcal{U}(w_F(x))$, then

$$w_F(x) \not\leq y \quad \text{and} \quad w_F(x) = \bigwedge \{m \in M \mid x \leq m\},$$

that is, $w_F(x) \leq y$ and it is a contradiction.

Therefore, $\mathcal{U}(w_F(x)) \subseteq M' \cup \mathcal{U}(x)$. $\square$

Definition 5.6. Let A be a frame.

- (1) A is *strongly stacked* if $\mathcal{U}_*[Q']\mathcal{U} = v_F$.
- (2) A is *stacked* if $(\mathcal{U}_*([Q'])\mathcal{U})(0) = v_F(0)$.

for any (F, Q) HM pair.

Note that strongly stacked implies stacked. Our aim is to prove that the above properties are conservative, that is, S is a stacked (strongly stacked) space if and only if its frame of open sets $\mathcal{O}S$ is stacked (strongly stacked).

Proposition 5.7. *The notions of strongly stacked and stacked are conservative.*

Proof. Consider $S \in \text{Top}$, $A = \mathcal{O}S$, $F \in A^\wedge$ and $Q \in \mathcal{Q}S$.

First, suppose that S is strongly stacked, then $v_F = [Q']$ and, by Lemma 5.3, $\mathcal{U}_*[Q']\mathcal{U} = [Q'] = v_F$. Therefore $\mathcal{O}S$ is strongly stacked.

Now, suppose that S is stacked, then

$$\overline{Q} = \partial_F^\infty(S) \iff \overline{Q}' = (\partial_F^\infty(S))'.$$

Furthermore,

$$\begin{aligned} (\mathcal{U}_*[Q']\mathcal{U})(\emptyset) &= \mathcal{U}_*([Q'](\{V \in \text{pt } \mathcal{O}S \mid \emptyset \not\subseteq V\})) \\ &= \mathcal{U}_*([Q'](\emptyset)) = \mathcal{U}_*(Q')^\circ \\ &= \bigcup \{W \in \mathcal{O}S \mid W \subseteq (Q')^\circ\} \\ &= (Q')^\circ = \overline{Q}'. \end{aligned}$$

By Corollary 5.2, $v_F(\emptyset) = (\partial_F^\infty(\emptyset))'$. Therefore,

$$v_F(\emptyset) = (\mathcal{U}_*[Q']\mathcal{U})(\emptyset),$$

and $\mathcal{O}S$ is stacked. $\square$

6. SEPARATION PROPERTIES IN FRAMES

If $S \in \text{Top}$ is an T_1 and strongly stacked, then it holds that $w_\nabla = v_\nabla$. Additionally, if S is stacked and T_1 , then it is sober.

Thus, we could ask ourselves if the same occurs in frames. In this section we will study the above question.

Take $p \in \text{pt } A$ and $(F_p, \uparrow p)$ a HM pair, then for

$$M = \{m \in A \setminus F_p \mid (\forall a \in A \setminus F_p)[a \leq m]\},$$

the nuclei $w_{F_p}(x) = \bigwedge \{m \in M \mid x \leq m\}$ is $w_p(x)$.

To prove this, note that $A \setminus F_p = \{x \in A \mid x \leq p\} = \downarrow p$. As,

$$M = \{m \in \downarrow p \mid [\forall a \in \downarrow p](a \leq m)\} = \{p\}.$$

Hence, $w_{F_p}(x) = \bigwedge \{p \mid x \leq p\}$, that is,

$$(10) \quad w_{F_p}(x) = \begin{cases} 1, & \text{if } x \not\leq p \\ p, & \text{if } x \leq p \end{cases} = w_p(x)$$

for all $x \in A$. Therefore $w_{F_p} = w_p$.

Proposition 6.1. *Let $A \in \text{Frm}$ and (F, Q) be any HM pair. If $v_F(0) = w_F(0)$, then for all $p \in \text{pt } A$, p is maximal in $\text{pt } A$.*

Proof. Consider $p, q \in \text{pt } A$ such that $p \leq q$. Note that for $q \in \text{pt } A$, we have $(F_q, \uparrow q)$ a HM pair, with

$$F_q = \{x \in A \mid \uparrow q \subseteq \mathcal{U}(x)\} = \{x \in A \mid (\forall r \in \text{pt } A)[r \leq q \Rightarrow x \not\leq r]\}.$$

By hypothesis, $v_{F_q}(0) = w_{F_q}(0)$ and by (10)

$$q = w_q(0) = w_{F_q}(0) = v_{F_q}(0) \leq v_{F_q}(p) = p.$$

Therefore, $p = q$ and every point is maximal in $\text{pt } A$. $\square$

As we show later on (2), for all (F, Q, ∇) HM tupla we have

$$\mathcal{U} \circ v_F \leq v_\nabla \circ \mathcal{U}.$$

Additionally, if A is strongly stacked, then $v_F = \mathcal{U}_*[Q']\mathcal{U}$ and

$$\mathcal{U}_* \circ v_\nabla \circ \mathcal{U} \leq \mathcal{U}_* \circ [Q']\mathcal{U}$$

Thus, $v_F = \mathcal{U}_* \circ v_\nabla \circ \mathcal{U}$, equivalently $\mathcal{U} \circ v_F = v_\nabla \circ \mathcal{U}$.

Therefore, if A is strongly stacked, then the diagram (2) is commutative. The following result tells us under which conditions we have the converse implication.

Proposition 6.2. *If $A \in \text{Frm}$ is tidy and the diagram*

$$\begin{array}{ccc} A & \xrightarrow{v_F} & A \\ \mathcal{U} \downarrow & & \downarrow \mathcal{U} \\ \mathcal{OS} & \xrightarrow{v_\nabla} & \mathcal{OS} \end{array}$$

commutes, then A is strongly stacked.

Proof. Let $A \in \text{Frm}$, $S = \text{pt } A$ and (F, Q) be any HM pair. By hypothesis, A is tidy and by previous results we have the implications

$$A \text{ tidy} \Rightarrow \mathcal{OS} \text{ tidy} \Rightarrow S \text{ stacked and packed}.$$

Since S is packed, if $Q \in \mathcal{QS}$, then $Q \in \mathcal{CS}$ and $Q' \in \mathcal{OS}$. Then, as S is stacked, we have that $\overline{Q} = \partial_\nabla^\infty(S)$, that is, $Q' = v_\nabla(\emptyset)$ and as $\mathcal{OS}$ is tidy, $v_\nabla = u_{Q'} = [Q']$.

Additionally, if $v_\nabla \circ \mathcal{U} = \mathcal{U} \circ v_F$, then

$$v_\nabla \circ \mathcal{U} \leq \mathcal{U} \circ v_F \quad \text{and} \quad \mathcal{U} \circ v_F \leq v_\nabla \circ \mathcal{U},$$

that is,

$$[Q'] \circ \mathcal{U} \leq \mathcal{U} \circ v_F \quad \text{and} \quad \mathcal{U} \circ v_F \leq [Q'] \circ \mathcal{U}.$$

By properties of the right adjoint

$$\mathcal{U}_* \circ [Q'] \circ \mathcal{U} \leq v_F \quad \text{and} \quad v_F \leq \mathcal{U}_* \circ [Q'] \circ \mathcal{U}.$$

Therefore $v_F = \mathcal{U}_* \circ [Q'] \circ \mathcal{U}$ and A is strongly stacked. $\square$

Consequently, we have the following result.

Corollary 6.3. *If A is tidy, then $S = \text{pt } A$ is strongly stacked.*

Proof. Since S is strongly stacked if and only if $v_\nabla = [Q']$, the result follows from the fact that

$$\mathcal{O}S \text{ tidy} \Rightarrow S \text{ strongly stacked}$$

$\square$

With the morphism $\mathcal{U}_A$, we have the nuclei sp_A defined by

$$\text{sp}_A(x) = \bigwedge \{p \in \text{pt } A \mid x \leq p\}$$

for $x \in A$. Combining this fact with the Proposition 6.1 we know information about the collapse on admisibility interval.

Proposition 6.4. *For $A \in \text{Frm}$ and (F, Q) any HM pair, the following statements hold:*

- (1) *If A is T_1 and stacked, then $v_F(0) = w_F(0)$.*
- (2) *If $v_F(0) = w_F(0)$, then A is stacked.*
- (3) *If $v_F(0) = w_F(0)$ and the set $Y = \{q \in \text{pt } A \mid a \leq q\}$ is non-empty for all $a \in A$, then A is T_1 .*

Proof.

- (1) If A is T_1 , then

$$\mathcal{O}S \text{ is } T_1 \Rightarrow S \text{ is } T_1 \Rightarrow Q = M.$$

Since A is stacked,

$$v_F(0) = (\mathcal{U}_* \circ [Q'] \circ \mathcal{U})(0) = (\mathcal{U}_* \circ [M'] \circ \mathcal{U})(0) = w_F(0).$$

- (2) Recall that for every HM pair, the following inequality holds:

$$(11) \quad v_F \leq \mathcal{U}_* \circ [Q'] \circ \mathcal{U} \leq w_F$$

and by hypothesis $v_F(0) = w_F(0)$, that is, $v_F(0) = (\mathcal{U}_* \circ [Q'] \circ \mathcal{U})(0) = w_F(0)$. Therefore, A is stacked.

- (3) Consider $p \in \text{pt } A$ and $a \in A$ such that $p \leq a$. By hypothesis $\{q \in \text{pt } A \mid a \leq q\} \neq \emptyset$. Then

$$p \leq a \leq \text{sp}(a) = \bigwedge \{q \in \text{pt } A \mid a \leq q\}$$

for some $q \in Y$. Thus, $p \leq a \leq q$ and by Proposición 6.1, $p = q$. Therefore, $p = a$ and every point is maximal in A , that is, A is T_1 . $\square$

Corollary 6.5. *For $A \in \text{Frm}$ and (F, Q) any HM pair, the following statements hold:*

- (1) *If A is T_1 and strongly stacked, then $v_F = w_F$.*
- (2) *If $v_F = w_F$, then A is strongly stacked.*
- (3) *If $v_F = w_F$ and the set $Y = \{q \in \text{pt } A \mid a \leq q\}$ is non-empty for all $a \in A$, then A is T_1 .*

Proof.

- (1) If A is strongly stacked, then $v_F = \mathcal{U}_* \circ [Q'] \circ \mathcal{U}$ and by T_1 we have that $Q = M$. Therefore $v_F = w_F$.
- (2) This follows from (11).
- (3) This follows from Proposition 6.4.

□

Note that, by 6.1 and 6.4, if $v_F(0) = w_F(0)$, then A is stacked and every point is maximal in $\text{pt } A$. Particularly, for each $p \in \text{pt } A$ and $(F_p, \uparrow p)$ it holds $v_{F_p}(0) = p$. Furthermore,

$$v_{F_p}(p) = v_{F_p}(0) = 0_{A_{F_p}}.$$

If A is T_1 , p is maximal in A , then p is maximal in A_{F_p} and $A_{F_p} = \{q, 1\}$.

Now, $F_{F_p} = \nabla(v_{F_p})$ and $x \in F_p$ if and only if $v_{F_p}(x) = 1$. In this way, under the previous hypotheses, it holds that

$$v_{F_p}(x) = \begin{cases} 1, & \text{si } x \not\leq p \\ p, & \text{si } x \leq p \end{cases} = \begin{cases} 1, & \text{si } x \in F_p \\ p, & \text{si } x \notin F_p \end{cases} = w_{F_p}(x).$$

Therefore, if A is T_1 and stacked, then $v_{F_p} = w_{F_p}$ for each $p \in \text{pt } A$. With this in mind, we could ask ourselves if there exists some way to recover the notion of strongly stacked by means of T_1 and stacked.

Now, we need the following information:

- (1) Let $A^\wedge$ be a topological space where the basic open sets of $\mathcal{O}(A^\wedge)$ are given by

$$D(x) = \{F \in A^\wedge \mid x \in F\}$$

where $x \in A$.

- (2) For $A \in \text{Frm}$, we can construct a frame VA as in [Sim04]. Each element in $\text{pt}(VA)$ is of the form

$$(F, a), \quad \text{where } F \in A^\wedge \text{ y } a \in [v_F(0), w_F(0)].$$

- (3) Take a continuous surjective map $\pi: \text{pt}(VA) \rightarrow A^\wedge$, defined by

$$(F, a) \mapsto F.$$

- (4) By the functor $\mathcal{O}(-)$ we have

$$\mathcal{O}(\pi) = \pi^{-1}: \mathcal{O}(A^\wedge) \rightarrow \mathcal{O}(\text{pt}(VA)).$$

(5) Open sets in $\text{pt}(VA)$ are given by the following:

$$(12) \quad (F, a) \in [-](x) \iff x \in F \quad \text{and} \quad (F, a) \in \langle - \rangle(x) \iff x \not\leq a.$$

Particularly,

$$(13) \quad a \in [F](x) \iff x \in F \quad \text{and} \quad a \in \langle F \rangle(x) \iff x \not\leq a.$$

Proposition 6.6. *Let $A, VA \in \text{Frm}$ and $A^\wedge$ be as above. The following statements hold:*

- (1) *If A is T_1 and stacked, then $\mathcal{O}(A^\wedge) \cong \mathcal{O}(\text{pt}(VA))$ under the continuous function π .*
- (2) *If the morphism $\mathcal{O}(\pi) = \pi^{-1}$ induces that $\mathcal{O}(A^\wedge) \cong \mathcal{O}(\text{pt}(VA))$, then A is stacked.*
- (3) *If the morphism $\mathcal{O}(\pi) = \pi^{-1}$ induces that $\mathcal{O}(A^\wedge) \cong \mathcal{O}(\text{pt}(VA))$ and*

$$\mathcal{V}(a) = \{q \in \text{pt } A \mid a \leq q\} \neq \emptyset,$$

then A is T_1 .

Proof.

- (1) If A is T_1 and stacked, then for each $F \in A^\wedge$ it holds that

$$v_F(0) = w_F(0).$$

Take $(F, a), (G, b) \in \text{pt}(VA)$ such that $\pi(F, a) = F = \pi(G, a) = G$. By T_1 and stacked we have

$$a = v_F(0) = v_G(0) = b.$$

Therefore $(F, a) = (G, b)$, that is, π is injective. Since it is also surjective, π is bijective.

We must show that $\pi^{-1}: A^\wedge \rightarrow \text{pt}(VA)$ is continuous. For this, recall that by (12)

$$\begin{aligned} (F, a) \in [-](x) &\iff x \in F \\ &\iff (F, a) \in \pi^{-1}(D(x)) \\ &\iff \pi([-](x)) = D(x). \end{aligned}$$

For the other basic open sets, by (13) we have

$$a \notin \langle F \rangle(x) \iff x \leq a \iff v_F(x) \leq a.$$

By T_1 and stacked we have that $a = v_F(0)$ and thus

$$v_F(0) \notin \langle F \rangle(x) \iff x \leq v_F(0) \iff v_F(x) = v_F(0).$$

As $v_F(0) \neq 1$,

$$v_F(0) \notin \langle F \rangle(x) \iff x \notin F$$

or equivalently

$$a \in \langle F \rangle(x) \iff (F, a) \in \langle - \rangle(x) \iff x \in F.$$

Therefore, $\pi(\langle - \rangle(x)) = D(x)$ y π^{-1} is continuous. Hence, π is a homeomorphism and

$$\mathcal{O}(A^\wedge) \cong \mathcal{O}(\text{pt}(VA)).$$

- (2) Suppose that $\mathcal{O}(A^\wedge) \cong \mathcal{O}(\text{pt}(VA))$ under $\mathcal{O}(\pi) = \pi^{-1}$. Since π^{-1} is bijective, then π is also bijective. Thus, if $F \in A^\wedge$, then

$$(F, v_F(0)), (F, w_F(0)) \in \text{pt}(VA)$$

and

$$\pi(F, v_F(0)) = F = \pi(F, w_F(0)).$$

By the injectivity of π we have that $v_F(0) = w_F(0)$ and therefore A is stacked.

- (3) Suppose that $\mathcal{O}(A^\wedge) \cong \mathcal{O}(\text{pt}(VA))$ and A is stacked. Consider $a \in A$ and $p \in \text{pt } A$ such that $p \leq a$. Moreover, by the previous item we have that

$$\mathcal{O}(A^\wedge) \cong \mathcal{O}(\text{pt}(VA)) \Rightarrow v_F(0) = w_F(0)$$

and every point is maximal in $\text{pt } A$. Since $\mathcal{V}(a) \neq \emptyset$, there exists $q \in \text{pt } A$ such that $a \leq q$. By the maximality of p and q , we have

$$p \leq a \leq q = p.$$

Therefore, $p = a$ and A is T_1 .

□

Note that item (3) holds whenever A is spatial. In the spatial setting, this corresponds to the topological fact that

$$S \text{ is } T_1 \text{ and stacked} \Rightarrow S \text{ is sober.}$$

However, the point-free setting allows for a potentially broader phenomenon, since the conditions of being T_1 and stacked do not a priori force a frame to be spatial. This leads naturally to the following question:

Do there exist T_1 stacked frames that are not spatial?

7. QUOTIENTS OF STACKED FRAMES

Remember that for $(f: A \rightarrow B) \in \text{Frm}$, $j \in NA$ and $k \in NB$, we have the following definitions:

$$N(f)(j) = \bigvee \{u_{f(j(a))} \wedge v_{f(a)} \mid a \in A\} \in NA_j$$

and

$$N(f)_*(k) = f_* \circ k \circ f \in NA,$$

where f_* is the right adjoint of f .

Proposition 7.1. *Let A be a frame and $j \in NA$. If A is strongly stacked, then A_j is strongly stacked.*

Proof. Consider $S = \text{pt } A$, $S_j = \text{pt } A_j$ and let $F \in A_j^\wedge$ be. We define:

$$G = j_*(F), \quad Q = S_j \setminus F, \quad R = S \setminus G,$$

where j_* is the right adjoint associated with the frame morphism $j^*: A \rightarrow A_j$ and $x \in G$ if and only if $j(x) \in F$. Considering the spatial reflections

$$\mathcal{U}: A \rightarrow \mathcal{O}S \quad \text{and} \quad \mathcal{U}_j: A_j \rightarrow \mathcal{O}S_j$$

we have the following diagram

$$\begin{array}{ccccc}
 & & A & \xrightarrow{\mathcal{U}} & \mathcal{O}S \\
 & \nearrow v_G & \downarrow j^* & \nearrow [R'] & \downarrow sj \\
 A & \xrightarrow{\mathcal{U}} & \mathcal{O}S & & A_j \xrightarrow{\mathcal{U}_j} \mathcal{O}S_j \\
 \downarrow j^* & & \downarrow sj & \nearrow v_F & \downarrow [Q'] \\
 A_j & \xrightarrow{\mathcal{U}_j} & \mathcal{O}S_j & &
 \end{array}$$

Using the right adjoints of $\mathcal{U}$ and $\mathcal{U}_j$ we obtain

$$v_F, \quad l = N(\mathcal{U}_j)_*([Q']), \quad m = N(j^*)(N(\mathcal{U})_*([R'])), \quad n = N(j^*)(N(\mathcal{U}_j \circ j^*)_*)([Q'])$$

are nuclei in NA_j . Let us see what is the relationship between these nuclei.

By construction, we have $v_F \leq N(\mathcal{U}_j)_*([Q'])$ and we must verify the other inequality for A_j to be strongly stacked.

By hypothesis, we have $v_G = N(\mathcal{U})_*([R'])$. Moreover, for $x \in A$, we have $j(v_x) \leq v_{j(x)}$ and, by definition of the adjusted nuclei associated with F and G ,

$$(14) \quad j \circ v_G \leq v_F.$$

Furthermore, for $(j: A \rightarrow A_j) \in \text{Frm}$

$$N(j^*)(v_G) = \bigvee \{u_{j(v_G(a))} \wedge v_{j(a)} \mid a \in A\} \quad \text{y} \quad v_F = \bigvee \{u_{v_F(a)} \wedge v_a \mid a \in A_j\}.$$

Note, by (14), $u_{j(v_G)} \leq u_{v_F}$ and therefore

$$(15) \quad m \leq v_F.$$

To l and n , we have

$$n = \bigvee \{u_{j^*(n(x))} \wedge v_{j^*(x)} \mid x \in A\} \quad \text{and} \quad l = \bigvee \{u_{l(x)} \wedge v_x \mid x \in A_j\},$$

where

$$j^*(n(x)) = j^*((j_* \circ \mathcal{U}_{j*} \circ [Q'] \circ \mathcal{U}_j \circ j^*)(x)) \quad \text{y} \quad l(x) = N(\mathcal{U}_j)_*([Q'])(x).$$

Then,

$$\begin{aligned} j^*(n(x)) &= j^*(j_*(\bigvee\{a \in A \mid \mathcal{U}_j(a) \subseteq Q' \cup \mathcal{U}_j(j(x))\})) \\ &= \bigvee\{a \in A \mid \mathcal{U}_j(a) \subseteq Q' \cup \mathcal{U}_j(j(x))\}. \end{aligned}$$

Since $x \in A_j$,

$$l(x) = \bigvee\{a \in A \mid \mathcal{U}_j(a) \subseteq Q' \cup \mathcal{U}_j(j(x))\}.$$

Therefore $j(n(x)) = l(x)$.

To $m(x)$ we have

$$m(x) = \bigvee\{u_{j^*}(\mathcal{U}_* \circ [R'] \circ \mathcal{U})(x) \wedge v_{j^*}(x) \mid x \in A\}.$$

Since $A_j \subseteq A$, then $Q = R \cap A_j$ and

$$m(x) = \bigvee\{u_{j^*}(\mathcal{U}_* \circ [Q'] \circ \mathcal{U})(x) \wedge v_x \mid x \in A_j\}.$$

Note

$$\begin{aligned} j^*(\mathcal{U}_* \circ [Q'] \circ \mathcal{U})(x) &= j^*(\bigvee\{a \in A \mid \mathcal{U}(a) \subseteq Q' \cup \mathcal{U}(x)\}) \\ &= \bigvee\{j^*(a) \in A_j \mid \mathcal{U}(j^*(a)) \subseteq Q' \cup \mathcal{U}(x)\}. \end{aligned}$$

To $x \in A_j$

$$\begin{aligned} \mathcal{U}(j^*(a)) \subseteq Q' \cup \mathcal{U}(j^*(x)) &\iff \mathcal{U}(j^*(a)) \cap S_j \subseteq (Q' \cup \mathcal{U}(j^*(x))) \cap S_j \\ &\iff \mathcal{U}_j(j^*(a)) \subseteq Q' \cup \mathcal{U}_j(j^*(x)), \end{aligned}$$

that is,

$$\bigvee\{j^*(a) \in A_j \mid \mathcal{U}(j^*(a)) \subseteq Q' \cup \mathcal{U}(x)\} = j^*(N(\mathcal{U}_j \circ j)_*)([Q']).$$

Thus, we have

$$\bigvee\{u_{j^*}(N(\mathcal{U}_j \circ j)_*)([Q'])(x) \wedge x \mid x \in A\} = m(x).$$

Since $(N(\mathcal{U}_j \circ j)_*)([Q']) \leq j(N(\mathcal{U}_j \circ j)_*)([Q'])$ and by (15)

$$l \leq m \leq v_F \leq l.$$

Therefore $v_F = \mathcal{U}_{j^*} \circ [Q'] \circ \mathcal{U}_j$, that is, A_j is strongly stacked. $\square$

Since every strongly stacked frame is stacked, we obtain the following result.

Corollary 7.2. *Let A be a frame and $j \in NA$. If A is strongly stacked, then A_j is stacked.*

Proof. By the previous proposition, if A is strongly stacked, then A_j is strongly stacked and this implies that A_j is stacked. $\square$

8. RELATIONSHIP TO THE PROPERTY (H)

If A is a Hausdorff frame, then A is T_1 . In this way, to be able to use the example of Paseka and Smarda it is enough that (H) implies stacked.

Proposition 8.1. *If $A \in \text{Frm}$ satisfies (H), then A is stacked.*

Proof. If A satisfies (H), then A is T_1 , that is, for every pair of points $x, y \in A$ with $x \neq y$, there exist neighborhoods U, V of x, y respectively such that $U \cap V = \emptyset$. This implies that

$$\mathcal{U}_* \circ [Q'] \circ \mathcal{U} = w_F,$$

that is, $\mathcal{U}_* \circ [Q'] \circ \mathcal{U}(0) = w_F(0)$ and since $w_F \neq \text{tp}$, we have $w_F(0) \neq 1$.

By construction, we have that $v_F(0) \leq w_F(0)$ and thus, we need to verify that the reverse inequality holds to conclude that A is stacked. For a contradiction, suppose that

$$w_F(0) \not\leq v_F(0).$$

Since $1 \neq w_F(0) \not\leq v_F(0)$, by (H), exists $c \in A$ such that

$$c \not\leq w_F(0) \quad \text{y} \quad \neg c \not\leq v_F(0).$$

If $\neg c \not\leq v_F(0)$, then $c \notin F$ □

9. MARCOS KC

Definition 9.1. Let A be a frame. We say that A is:

- (1) *KC* if each compact quotient is closed.
- (2) *Hausdorff closed compact* (or *KCH* for short) if each compact quotient of A is closed and Hausdorff.

La Definición 9.1 nos dice que si $j \in NA$, entonces $\nabla(j) \in A^\wedge$ y $j = u_a$ para algún $a \in A$. De manera adicional, para 2) pedimos que A_j cumpla la propiedad (H).

Proposition 9.2. *If $A \in \text{Frm}$ satisfies KC, then A_j satisfies KC for each $j \in N(A)$.*

Proof. Consider $k \in NA_j$ such that $(A_j)_k$ is compact. If $(A_j)_k$ is compact, then $\nabla(k) \in A_j^\wedge$ and, by Proposition 3.1, $j_* \nabla(K) \in A^\wedge$.

Let $l = j_* \circ k \circ j^* \in NA$ be, then A_l is a compact quotient of A and there exists $a \in A$ such that $l = u_a$. Thus

$$\begin{array}{ccccccc} & & & l & & & \\ & & & \curvearrowright & & & \\ A & \xrightarrow{j^*} & A_j & \xrightarrow{k} & (A_j)_k & \xrightarrow{j_*} & A_j \subseteq A \end{array}$$

and $a \vee x = k(j(x))$. Therefore, if $x = a$, $k(j(x)) = a$.

We need that $k = u_b$ for some $b \in A_j$. If $x \in A_j$ and $b = j(a)$

$$\begin{aligned} u_b(x) &= b \vee x = b \vee j(x) = j(j(a) \vee j(x)) \\ &= j(k(j(a)) \vee x) \\ &= j(u_a(x)) \\ &= j(k(x)) \\ &= k(x). \end{aligned}$$

Hence $u_b = k$. □

Proposition 9.3. *If A is a KC frame, then A is a T_1 frame.*

Proof. Let $p \in \text{pt } A$ and $a \in A$ such that $p \leq a \leq 1$. Consider

$$w_p(x) = \begin{cases} 1 & \text{si } x \not\leq p \\ p & \text{si } x \leq p \end{cases}$$

to $x \in A$. $P = \nabla(w_p) = \{x \in A \mid x \not\leq p\}$ is a completely prime filter (in particular, $P \in A^\wedge$). Because A is KC , then A_{w_p} is a compact closed quotient. Thus $u_p = w_p$ and

$$u_p(a) = a \quad \text{y} \quad w_p(a) = 1.$$

that is, $a = 1$. Therefore p is maximal. □

One of our aims is study nuclei such that produce compact quotient, particularly, nuclei that produce compact closed quotient.

Define the following subset of A

$$uA = \{a \in A \mid A_{u_a} \text{ es un cociente compacto}\}.$$

Note that, to $a \in uA$ we have $\nabla(u_a) \in A^\wedge$, that is, define the map

$$\alpha: uA \rightarrow A^\wedge.$$

Let us see the properties of this map.

Lemma 9.4. *For $uA \subseteq A$ as before, the following statements hold:*

- (1) *If A is tidy, then α is surjective.*
- (2) *If u_a is fitted for all $a \in uA$, then α is injective.*
- (3) *If A is tidy and $a \in uA$, then α is bijective if and only if u_a is fitted.*

Proof.

- (1) Consider $F = \nabla(j) \in A^\wedge$ and v_F the fitted nucleus associated with F . If A is tidy, then $v_F = u_d$, where $d = v_F(0) \in A$. Hence,

$$\alpha(d) = \nabla(u_d) = \nabla(v_F) = F.$$

and α is surjective.

- (2) Consider $a, b \in uA$ such that $\alpha(a) = \alpha(b)$, that is, $\nabla(u_a) = \nabla(u_b)$. By hypothesis, u_a, u_b are fitted nuclei, that is,

$$u_c \leq u_d \quad \text{y} \quad u_c \geq u_d.$$

Evaluating both nuclei at 0 we get $c = d$ and therefore α is injective.

- (3) Let A be tidy, $a \in uA$ and suppose that α is bijective. Consider $F = \nabla(u_a)$ and v_F the fitted nucleus associated with F . By efficiency, $v_F = u_d$, that is,

$$\alpha(a) = \nabla(u_a) = \nabla(v_F) = \nabla(u_d) = \alpha(d)$$

where $d = v_F(0)$.

By hypothesis, α is bijective, then, by injectivity, $a = d$. Therefore $v_F = u_a$ and u_a is fitted.

Conversely, since A is tidy, then by 1), α is surjective. By hypothesis, u_a is fitted and, by 2), α is injective. Therefore α is bijective. □

Using KC frames, we have the following result.

Corollary 9.5. *Let A be a KC frame and $F \in A^\wedge$. Then α is bijective if and only if for all $j \in [v_F, w_F]$, j is fitted.*

Proof. Suppose that A is a KC frame and let $F \in A^\wedge$. By definition, for all $j \in [v_F, w_F]$, we have $j = u_a$ for some $a \in A$. Since A is a KC frame, it is efficient and by the previous lemma, α is bijective if and only if for all $j \in [v_F, w_F]$, j is fitted. □

Proposition 9.6. *Let A be a frame. A is KC if and only if A is tidy and $\mathcal{U} \circ v_F = v_\nabla \circ \mathcal{U}$.*

Proof.

$\Rightarrow$): Suppose that A is a KC frame. Trivially, KC implies efficient and thus, it remains to verify that KC implies that $\mathcal{U} \circ v_F = v_\nabla \circ \mathcal{U}$ or equivalently, by properties of the right adjoint,

$$v_F = \mathcal{U}_* \circ v_\nabla \circ \mathcal{U}.$$

If A is a KC frame, then for any $F \in A^\wedge$, it holds that for all $j \in [v_F, w_F]$, $j = u_a$ for some $a \in A$. In particular,

$$v_F = u_d, \quad \text{y} \quad \mathcal{U}_* \circ v_\nabla \circ \mathcal{U} = u_a$$

where $d, a \in A$.

By construction, always holds that $u_d \leq u_a$. To see the other inequality, we verify that the function α is injective. Note that $d \leq a$

$\Leftarrow$): Suppose that A is efficient and $\mathcal{U} \circ v_F = v_\nabla \circ \mathcal{U}$ for all HM triples (F, Q, ∇) . First, if A is efficient, then A is T_1 . Moreover, if A is efficient and $\mathcal{U} \circ v_F = v_\nabla \circ \mathcal{U}$, then A is strongly stacked. Therefore, A is T_1

and strongly stacked. The above implies that $v_F = w_F$ and, by efficiency, $v_F = u_d$, where $d = v_F(0)$. Thus, for all $j \in [v_F, w_F]$, $j = u_d$, meaning that A is KC .

□

10. EXAMPLES

10.1. **The frame** $[0, 1]$. Following the idea presented in [Sim06b] (or, alternatively, in [Sim04]), let us consider the linearly ordered frame

$$A = [0, 1].$$

In this case the following observations hold.

- (1) For all $a, b \in A$ the implication $(a \succ b)$ is given by

$$(a \succ b) = \begin{cases} 1, & \text{si } a \leq b, \\ b, & \text{si } a > b. \end{cases}$$

- (2) For $a, x \in A$, the distinguished nuclei u_a, v_a, w_a are given by

$$u_a(x) = \begin{cases} x & \text{si } a \leq x, \\ a & \text{si } x < a, \end{cases}, \quad v_a(x) = \begin{cases} 1 & \text{si } x \geq a, \\ a & \text{si } x < a, \end{cases}, \quad w_a(x) = \begin{cases} 1 & \text{si } x > a, \\ a & \text{si } x \leq a. \end{cases}$$

- (3) If $F \subseteq A$ is a filter, then F is always admissible and has the form

$$F = \nabla(w_a) = (a, 1] \quad \text{o} \quad F = \nabla(v_a) = [a, 1]$$

for some $a \in A$. In particular, $F \in A^\wedge$ if and only if $F = (a, 1]$.

- (4) If $F \in A^\wedge$, the adjusted nucleus associated v_F is calculated as

$$(16) \quad v_F(x) = l_m(x) = \begin{cases} 1 & \text{if } x > m, \\ x & \text{if } x \leq m. \end{cases}$$

where $l_m = v_m \wedge w_m$ and $m = \bigwedge F$.

- (5) For $S = \text{pt } A$ it holds that $S = [0, 1)$ and the open sets in $\mathcal{O}S$ are of the form

$$\mathcal{U}_A(x) = \{p \in S \mid p < x\} = [0, x).$$

- (6) The frame A is not T_1 , since no $p \in S$ is maximal. Consequently, A does not satisfy any stronger separation property in Frm .

- (7) For $F \in A^\wedge$ the compact saturated set associated $Q \in \mathcal{Q}S$ is of the form $Q = [0, m]$.

- (8) It holds that $[Q'] = v_F$; therefore, S is strongly stacked.

- (9) Since $[0, x) \in \mathcal{O}S$, then $[x, 1] \in \mathcal{C}S$ and $Q = [0, m]$. Therefore, the compact subsets are not closed; that is, S is not packed.

- (10) The frame A is not efficient.

The above resume some characteristics about the frame $[0, 1]$. Now, we will perform the analysis of the patch constructions of A and its point space S .

Remember that for $A \in \text{Frm}$ and $S \in \text{Top}$ we have

$$\text{Pbase} = \{u_a \wedge v_F \mid a \in A, F \in A^\wedge\} \quad \text{y} \quad \text{pbase} = \{U \cap Q' \mid U \in \mathcal{O}S, Q \in \mathcal{Q}S\}$$

where Pbase generates the patch frame of A and pbase generates the topology of the patch space of S . By (16) we have that

$$\text{Pbase} = \{u_a \wedge l_m \mid a \in [0, 1], m \in [0, 1]\}.$$

For $x \in A$ we have

$$(u_a \wedge l_m)(x) = u_a(x) \wedge l_m(x).$$

Let us analyze the value of this map according to the relative position of x and m .

Case 1: If $x < m$, then

$$(u_a \wedge l_m)(x) = u_a(x) \wedge x = x.$$

Case 2: If $x = m$, then

$$(u_a \wedge l_m)(x) = u_a(m) \wedge m = x.$$

Case 3: If $x > m$, then

$$(u_a \wedge l_m)(x) = u_a(x) \wedge 1 = u_a(x).$$

In the first two cases, the value of a does not affect the evaluation. Thus, the only non-trivial case occurs when $m < x$, where the value of $(u_a \wedge l_m)(x)$ depends on the relative position of a and x . We therefore distinguish the following two subcases.

Case 3.1: If $m < x < a$, then

$$(u_a \wedge l_m)(x) = a.$$

Case 3.2: If $m < a \leq x$, then

$$(u_a \wedge l_m)(x) = x.$$

From the preceding analysis, we conclude that the non-trivial generators of Pbase correspond precisely to the pairs (m, a) with $m < a$. Consequently, the frame PA is determined by the open intervals of the form (m, a) , where $m \in \text{pt } A$ and $a \in A$.

We now turn to the question of whether PA satisfies any separation property. To this end, we recall the relationship between the point-free patch construction and its spatial counterpart.

Let A be an arbitrary frame and let $S = \text{pt } A$ be its space of points. The two patch constructions are related by the following commutative diagram:

$$\begin{array}{ccccc} A & \longrightarrow & PA & \hookrightarrow & NA \\ \downarrow u_A & & \downarrow P(u_A) & & \downarrow N(u_A) \\ \mathcal{O}S & \longrightarrow & P\mathcal{O}S & \hookrightarrow & N\mathcal{O}S \\ & \searrow & \downarrow \pi & & \downarrow \sigma \\ & & \mathcal{O}^p S & \hookrightarrow & \mathcal{O}^f S \end{array}$$

where $\mathcal{O}^f S$ is the Skula topology (or front topology) of S . In [SS06b] is named *the patch assembly diagram* and more details about it can also be found in [Sex03].

In our case, $\mathcal{U}_A$ is an isomorphism. Moreover, under $\mathcal{U}_A$, the patch construction turns out to be functorial and, therefore, $PA \cong POS$. Similarly, $NA \cong NOS$. To describe explicitly POS and NOS , we use the following result, which can be consulted in [ÁBMZ20].

Proposition 10.1. *Let S be a partially ordered set.*

- (1) *If S has no infinite antichains, then NOS is spatial.*
- (2) *If S is totally ordered, then NOS is spatial.*

Thus, by Proposition 10.1 we have that NOS is spatial and therefore $NOS \cong \mathcal{O}^f S$. With all this information, the patch assembly diagram, in our particular case is as follows.

$$\begin{array}{ccccc} A & \longrightarrow & PA & \longrightarrow & NA \\ \cong \downarrow & & \downarrow \cong & & \downarrow \cong \\ \mathcal{O}S & \longrightarrow & \mathcal{O}^p S & \longrightarrow & \mathcal{O}^f S \end{array}$$

Therefore, if we want to study the frame PA , we can study the patch topology $\mathcal{O}^p S$. Remember that

$$\text{pbase} = \{U \cap Q' \mid U \in \mathcal{O}S, Q \in \mathcal{Q}S\}.$$

In our case,

$$\text{pbase} = \{[0, a) \cap [m, 1) \mid a \in [0, 1], m \in [0, 1)\} = \{(m, a)\}.$$

Since the topology generated by the intervals (m, a) coincides with the usual topology on $[0, 1)$, the space ${}^p S$ is Hausdorff. Consequently, $\mathcal{O}^p S \cong PA$ satisfies **(H)**, even when A is not T_1 nor tidy.

Lemma 10.2. *Let A be a frame, if the morphism $pU: pA \rightarrow p\mathcal{O}S$ is an isomorphism, then A is strongly stacked.*

Proof. Consider any open filter F on A and the compact saturated determined by it, say Q , thus we have the nuclei v_F and $U_*[Q']U = Q[U]$, we need to observe the action of pU on these two nuclei, in order to do that remember that we have a commutative diagram:

$$\begin{array}{ccccc} A & \longrightarrow & pA & \xrightarrow{\iota} & NA \\ \mathcal{U}_A \downarrow & & \downarrow p(\mathcal{U}_A) & & \downarrow N(\mathcal{U}_A) \\ \mathcal{O}S & \longrightarrow & p\mathcal{O}S & \xrightarrow{\iota} & NOS \\ & \searrow & \downarrow \pi & & \downarrow \sigma \\ & & \mathcal{O}^p S & \longrightarrow & \mathcal{O}^f S \end{array}$$

since $N(\mathcal{U})\iota = v\mathcal{U}$, that is, $p\mathcal{U}$ is the restriction of $N(\mathcal{U})$ to pA , then

$$\text{p}\mathcal{U}(v_F) = N(\mathcal{U})(v_F) = \bigvee \{u_{\mathcal{U}(v_F)} \wedge v_{\mathcal{U}(x)} \mid x \in A\}$$

since the morphism $\mathcal{U}$ transforms open filters we have

$$\bigvee \{u_{\mathcal{U}(v_F)} \wedge v_{\mathcal{U}(x)} \mid x \in A\} = \bigvee \{u_{v_{\mathcal{U}F}} \wedge v_{\mathcal{U}(x)} \mid x \in A\}$$

But this representation is the corresponding to the nucleus $v_{\mathcal{U}F}$.

□

REFERENCES

- [ÁBMZ20] Francisco Ávila, Guram Bezhanishvili, PJ Morandi, and Angel Zaldívar, *The frame of nuclei on an alexandorff space*, Order (2020).
- [Hoc69] Melvin Hochster, *Prime ideal structure in commutative rings*, Transactions of the American Mathematical Society **142** (1969), 43–60.
- [Joh82] P. T. Johnstone, *Stone spaces*, Cambridge Studies in Advanced Mathematics, vol. 3, Cambridge University Press, Cambridge, 1982. MR 698074
- [PP12] J. Picado and A. Pultr, *Frames and locales: Topology without points*, Frontiers in Mathematics, Springer Basel, 2012.
- [PP21] Jorge Picado and Aleš Pultr, *Separation in point-free topology*, Springer, 2021.
- [Sex03] Rosemary A Sexton, *A point-free and point-sensitive analysis of the patch assembly*, The University of Manchester (United Kingdom), 2003.
- [Sim04] Harold Simmons, *The vietoris modifications of a frame*, Unpublished manuscript, 79pp., available online at <http://www.cs.man.ac.uk/hsimmons> (2004).
- [Sim06a] ———, *The assembly of a frame*, University of Manchester (2006).
- [Sim06b] ———, *Regularity, fitness, and the block structure of frames*, Applied Categorical Structures **14** (2006), 1–34.
- [SS06a] RA Sexton and H Simmons, *An ordinal indexed hierarchy of separation properties*, to appear (2006).
- [SS06b] ———, *Point-sensitive and point-free patch constructions*, Journal of Pure and Applied Algebra **207** (2006), no. 2, 433–468.