

Six concyclic points satisfy the Euclidean Ramsey tensor criterion

JD Jones

October 2026

Abstract

A finite set of points in Euclidean space is Ramsey if every finite colouring of a sufficiently high-dimensional Euclidean space contains a monochromatic congruent copy of it. The classification of finite Euclidean Ramsey configurations announced in the openai/math release of 2026 characterises the Ramsey sets that affinely span their space by a tensor criterion over $F \otimes_{\mathbb{Q}} F$, F the field generated by the coordinates, and proves that every set of at most five points on a circle is Ramsey; a seven-point circle set is known not to be. We show that every set of six distinct concyclic points satisfies the criterion, by an explicit certificate, and more generally that every spherical set of at most $m + 4$ points, m the dimension of its affine span, satisfies the criterion in its own space and field; for $m \geq 2$ the bound is sharp. We also show that every spherical set with algebraic coordinates satisfies the criterion, and that the criterion always implies sphericity. Given the classification, every spherical set of at most six points is Ramsey, seven is the least size of a non-Ramsey spherical set and, for $m \geq 2$, $m + 5$ the least size of one of affine dimension m , and the spherical conjecture of Erdős, Graham, Montgomery, Rothschild, Spencer and Straus holds for spanning configurations with algebraic coordinates (and, through an algebraic spanning congruent copy, for all of them). The unconditional statements are proved in Lean 4, and the Ramsey statements with the classification as an explicit hypothesis; the heptagon's failure of the criterion, the sharpness of the bound, the extension of the algebraic case to non-spanning configurations and the corollaries on least sizes are on paper, the first with an exact checker.

1 The criterion

Let $A = \{a_1, \dots, a_s\} \subset \mathbb{R}^d$ be a finite set of distinct points, $F = \mathbb{Q}(a_{ij}) \subset \mathbb{R}$ the field generated by their coordinates, $B = F \otimes_{\mathbb{Q}} F$, and $m: B \rightarrow F$ the multiplication map $x \otimes y \mapsto xy$. Write $p_i = (1, a_i) \in F^{d+1}$ and index $(d + 1) \times (d + 1)$ matrices by $\{0, 1, \dots, d\}$, the index 0 being the constant coordinate.

Definition 1 (the tensor criterion). A satisfies the criterion if there is $P \in \text{Mat}_{d+1}(B)$ with

$$\sum_{\alpha, \beta} (p_{i\alpha} \otimes 1) P_{\alpha\beta} (1 \otimes p_{i\beta}) = 0 \quad \text{for every } i, \quad m(P_{\alpha\beta}) = \delta_{\alpha\beta} \quad \text{for } \alpha, \beta \geq 1.$$

The constant row and column of $m(P)$ are unconstrained. Applying m to the first identity gives $p_i^{\top} m(P) p_i = 0$: the points lie on the F -rational quadric $|x|^2 - 2\langle c', x \rangle + C = 0$ read off from $m(P)$, which is a sphere. So the criterion asks that an F -rational sphere through the points lift to B with the evaluation identities holding before multiplication.

The classification paper [1] states (Theorem 1.1 there) that a set of at least two points that affinely spans $\mathbb{R}^d$ is Ramsey if and only if it satisfies the criterion, and proves (Corollary 7.4)

that every set of at most five points on a circle is Ramsey: for at most five concyclic points the quadratic evaluation rows $p_i p_i^\top$ are linearly independent over F (a product of two chords isolates each point), and independent rows lift by an adjugate (Proposition 7.3). Six concyclic points are the first case this does not reach: six points of a conic have rows spanning a space of dimension five. The heptagon paper [2] gives seven points on a circle that are not Ramsey. We use the classification only as a hypothesis in Section 6.

In Lean the definitions are those of the classification's challenge file, restated character for character; the repository is `six-concyclic-lean`, and the Lean name of each statement is given in brackets.

2 Six points on a circle

Theorem 2 (`six_concyclic_fieldCriterion`). *Every set of six distinct points on a circle in $\mathbb{R}^2$ satisfies the criterion.*

Proof. Let $a_0, \dots, a_5$ be the points. Since the real circle through three of them is unique and its centre is a rational function of their coordinates, there is an F -rational circle $|x|^2 - 2\langle c', x \rangle + C = 0$ through all six (in Lean this comes from an F -linear retraction $\mathbb{R} \rightarrow F$ fixing 1, applied to the real equation; `exists_rational_sphere`). Let H be its matrix: spatial block I_2 , constant row and column $-c'$, corner C .

Parametrisation. Take the base point $p = a_5$ and $e = p - c'$, which is nonzero because the points are distinct (`chord_dot_radius_ne_zero`). For a direction $x = (u, w) \in F^2 \setminus 0$ the second intersection of the line $p + tx$ with the circle is $q = p - 2 \frac{\langle e, x \rangle}{|x|^2} x$, and homogeneously

$$|x|^2 (1, q) = \Lambda (u^2, uw, w^2)^\top, \quad \Lambda = \begin{pmatrix} 1 & 0 & 1 \\ p_0 - 2e_0 & -2e_1 & p_0 \\ p_1 & -2e_0 & p_1 - 2e_1 \end{pmatrix},$$

with $\det \Lambda = 4|e|^2 \neq 0$ (`Lam_ver`, `Winv_mul_Lam`). The tangent direction $x = (-e_1, e_0)$ gives $|e|^2(1, p)$ (`Lam_tangent`). Put $x_j = a_j - p$ for $j < 5$ and $x_5 = (-e_1, e_0)$, so that $p_j = N_j^{-1} \Lambda v(x_j)$ with $N_j = |x_j|^2 \neq 0$ and $v(u, w) = (u^2, uw, w^2)$. Since $v(x)^\top \Lambda^\top H \Lambda v(x) = 0$ for every x and the symmetric form $\Lambda^\top H \Lambda$ vanishes on the Veronese curve, it is a multiple of the discriminant form $\delta(X, Y)^2$, $\delta = X_u Y_w - X_w Y_u$, whose matrix in the Veronese basis is $K = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -2 & 0 \\ 1 & 0 & 0 \end{pmatrix}$; explicitly $\Lambda^\top H \Lambda = -2|e|^2 K$ (`pullback`).

The bilinear forms. For three directions $x_j, j \in S$, let

$$\ell_S(X, Y) = \det \begin{pmatrix} X_u Y_u & X_u Y_w & X_w Y_u & X_w Y_w \\ x_{j,u} \otimes x_{j,u} & x_{j,u} \otimes x_{j,w} & x_{j,w} \otimes x_{j,u} & x_{j,w} \otimes x_{j,w} \end{pmatrix}_{j \in S}$$

a $(1, 1)$ -form in $X = x \otimes 1$ and $Y = 1 \otimes x$ with coefficients in B . It vanishes at $(x_j \otimes 1, 1 \otimes x_j)$ for $j \in S$ because the first row then repeats a row (`ell_vanishes`; this holds in any commutative ring). Put $q = \ell_{\{0,1,2\}} \ell_{\{3,4,5\}}$, a biquadratic form with a 3×3 Veronese matrix Q over B : $v(X)^\top Q v(Y) = q(X, Y)$ (`Qm_form`). Then q vanishes at all six points. Under m the two copies coincide; the corner coefficients of each ℓ_S vanish and the middle ones are $\pm V_S$, $V_S = \prod_{j < k \in S} \det(x_j, x_k)$ (`mult_coeffs`), so $m(Q) = V_1 V_2 K$. The brackets are nonzero: $\det(x_j, x_k)$ for two chords from p is twice the signed area of a triangle of three distinct points of a circle (`det_ne_zero`), and $\det(x_j, x_5) = \langle x_j, e \rangle = -|x_j|^2/2$ (`chord_dot_radius_ne_zero`).

The certificate. Let $W = \Lambda^{-1}$, $s = -2|e|^2/(V_1V_2) \in F$, and

$$P = (s \otimes 1)(W \otimes 1)^\top Q(1 \otimes W).$$

At the point i , $Wp_i = N_i^{-1}v(x_i)$, so the evaluation is $(sN_i^{-1} \otimes N_i^{-1})q(x_i \otimes 1, 1 \otimes x_i) = 0$ (`eval_P`). And $m(P) = sV_1V_2W^\top KW = W^\top(\Lambda^\top H\Lambda)W = H$ (`mult_P`), whose spatial block is I_2 . $\square$

No genericity is needed: the only inputs are that the points are distinct and concyclic. The certificate rests on universal polynomial identities in the left and right copies of the fourteen parameters (the base point, the centre and the five directions); certificate 0001 in `note/certificates` verifies these identities and twelve exact end-to-end evaluations independently of Lean.

3 Lifts through isolating quadrics

Lemma 3 (`fieldCriterion_of_isolators`). *Let H be an F -rational sphere matrix through A with spatial block I . If for every i there is an F -matrix q_i with $p_j^\top q_i p_j = \delta_{ij}$ for all j , then A satisfies the criterion.*

Proof. Let D be the $s \times (d+1)^2$ matrix over B with rows $(p_{i\alpha} \otimes 1)(1 \otimes p_{i\beta})$, S the $(d+1)^2 \times s$ matrix with columns $q_i \otimes 1$, $N = DS$, so that $m(N) = I$ by the isolating property, $b = H \otimes 1$, and $z = \det(N)b - S \operatorname{adj}(N)Db$. Then $Dz = \det(N)Db - N \operatorname{adj}(N)Db = 0$, and $m(z) = \det(mN)H - m(S) \operatorname{adj}(mN) m(Db) = H$ since $m(Db) = (p_i^\top H p_i)_i = 0$. Reshape z into P . $\square$

This is the argument of Proposition 7.3 of [1] without the choice of a minor. Isolating quadrics are products of affine functionals. On a sphere, three distinct points are never collinear (`gram_ne_zero`: if two chords from a common point were parallel, $w = \lambda u$, the sphere equation in the form $|x - p|^2 + 2\langle p - c', x - p \rangle = 0$ at both endpoints gives $\lambda(\lambda - 1)|u|^2 = 0$), so the line through two of the points has an F -rational affine equation that is nonzero at any third (`exists_line_functional`), and a point is separated from any other by $\langle a_i - a_j, x - a_j \rangle$.

Theorem 4 (`at_most_five_spherical_fieldCriterion`). *Every spherical set of at most five distinct points, in any dimension, satisfies the criterion.*

Proof. For each i split the other at most four points into two groups of at most two; the product of the two group functionals (a line, a point, or the constant 1) vanishes at the others and not at a_i . Apply Lemma 3 with the sphere of `exists_rational_sphere`. $\square$

Theorem 5 (`at_most_six_spherical_spanning_fieldCriterion`). *Every spherical set of at most six distinct points that affinely spans $\mathbb{R}^d$ satisfies the criterion.*

Proof. At most five points: Theorem 4. Six points: $d \leq 1$ is impossible, $d = 2$ is Theorem 2. For $d \geq 3$ and each i , some three of the other five points span a plane not containing a_i (`exists_isolating_triple`): fix two of them, a_j, a_k ; if the plane Π through a_j, a_k and a third point contains a_i , then any further point $a_m \notin \Pi$ gives a plane through a_j, a_k, a_m meeting Π in the line $a_j a_k$, which does not contain a_i ; and if every point lies in Π the set does not span. The affine function vanishing on that plane (the component of $a_i - a_j$ orthogonal to the plane, computed over F with the 2×2 Gram matrix of two chords; `exists_plane_functional`) times the line through the remaining two points isolates a_i ; apply Lemma 3. $\square$

Corollary 6 (`at_most_six_concyclic_fieldCriterion`). *Every set of at most six distinct points on a circle satisfies the criterion.*

4 The dimension theorem

Let $A \subset \mathbb{R}^D$ be a spherical set of s distinct points and m the dimension of its affine span.

Theorem 7 (`spherical_fieldCriterion_of_card_le_finrank_add_four`). *If $s \leq m + 4$, then A satisfies the criterion, in $\mathbb{R}^D$ and over its own coordinate field.*

For $m = 2$ this is Theorem 2 in any ambient space; it contains Theorems 4 and 5, and it gives `at_most_six_spherical_fieldCriterion`: every spherical set of at most six points, with any affine span, satisfies the criterion (six distinct points of a sphere are never collinear, so $m \geq 2$).

Proof. The real tensor ring. Write $R = \mathbb{R} \otimes_{\mathbb{Q}} \mathbb{R}$, with $\iota_l(x) = x \otimes 1$, $\iota_r(y) = 1 \otimes y$ and $m_R(x \otimes y) = xy$. Call $P \in \text{Mat}_{D+1}(R)$ a real certificate for A with multiplied matrix $H \in \text{Mat}_{D+1}(\mathbb{R})$ if $\sum_{\alpha, \beta} \iota_l(p_{i\alpha}) P_{\alpha\beta} \iota_r(p_{i\beta}) = 0$ for every i and $m_R(P) = H$ exactly (`RealCert`). We build a real certificate whose multiplied matrix is the real sphere matrix H_c of $|x - c|^2 = r^2$ (spatial block I) and descend it to F at the end.

Descent (`fieldCriterion_of_realCert`). Let (e_δ) be an F -basis of $\mathbb{R}$ and $\phi: \mathbb{R} \rightarrow F$ an F -linear map with $\phi(1) = 1$. The $\mathbb{Q}$ -bilinear map $(x, y) \mapsto \sum_\delta \phi(xe_\delta) \otimes y_\delta$, where $y = \sum_\delta y_\delta e_\delta$, induces $\Psi: R \rightarrow B$ with $\Psi((c \otimes 1)z) = (c \otimes 1)\Psi(z)$ and $\Psi((1 \otimes c)z) = (1 \otimes c)\Psi(z)$ for $c \in F$, and $m \circ \Psi = \phi \circ m_R$ (on $x \otimes y$: $\sum_\delta \phi(xe_\delta)y_\delta = \phi(xy)$). Applied entrywise to a real certificate, Ψ keeps the sandwiches zero, the $p_{i\alpha}$ being in F , and turns the spatial block I into $\phi(I) = I$. Nothing uses that B is a domain, which it need not be.

Four operations on real certificates. (i) A certificate over B pushes forward to R along $F \otimes F \rightarrow \mathbb{R} \otimes \mathbb{R}$ (`realCert_of_fieldCriterion`). (ii) The factor swap $\sigma(x \otimes y) = y \otimes x$ turns P into the certificate $(\sigma(P_{\beta\alpha}))_{\alpha\beta}$, whose multiplied matrix is $H^\top$, so the average has the symmetric multiplied matrix $(H + H^\top)/2$ (`realCert_symm`). (iii) If N is symmetric with zero spatial block and $p_i^\top N p_i = 0$ for every i , then with $l = (N_{00}/2, N_{01}, \dots, N_{0D})$ and $e_0 = (1, 0, \dots, 0)$ the matrix $(l \otimes 1)(1 \otimes e_0)^\top + (e_0 \otimes 1)(1 \otimes l)^\top$ has sandwiches $\langle l, p_i \rangle \otimes 1 + 1 \otimes \langle l, p_i \rangle = 0$, where $\langle l, p_i \rangle = p_i^\top N p_i / 2$ because the spatial block of N is zero, and multiplies exactly to N ; so H can be replaced by $H + N$ (`realCert_add_symm`). (iv) If $a_i = t + T b_i$ for a linear isometry $T: \mathbb{R}^d \rightarrow \mathbb{R}^D$ with matrix M ($M^\top M = I$) and the configuration (b_i) has a real certificate P whose multiplied matrix H has spatial block I_d , let L be the affine left inverse $(1, z) \mapsto (1, M^\top(z - t))$, so that $L p_i = (1, b_i)$, and G the matrix whose rows are the affine functionals $z \mapsto \langle (I - M M^\top) e_k, z - t \rangle$, which vanish at every a_i because $(I - M M^\top) M = 0$; then $(L \otimes 1)^\top P (1 \otimes L) + (G \otimes 1)^\top (1 \otimes G)$ is a real certificate for A with multiplied matrix $L^\top H L + G^\top G$, whose spatial block is $M M^\top + (I - M M^\top) = I$ (`realCert_of_isometric_image`).

Extension by a point (`realCert_insert`). Let $A' \subset A$ have a real certificate P with multiplied matrix H_c , and let $b \in A$ have p_b outside the linear span of the p_i , $i \in A'$. Choose a linear functional ℓ on $\mathbb{R}^{D+1}$ vanishing on that span with $\ell(p_b) = 1$, and let z be the sandwich of P at p_b . Then $P - z(\ell \otimes 1)(1 \otimes \ell)^\top$ has sandwich 0 at the points of A' and $z - z = 0$ at b , and it multiplies to $H_c - m_R(z) \ell \ell^\top = H_c$, because $m_R(z) = p_b^\top H_c p_b = 0$. Iterating (`realCert_of_finrank`): if the p_i , $i \notin A'$, add one dimension each to the span of the p_i , $i \in A'$, a real certificate of A' with multiplied matrix H_c extends to one of A .

The dichotomy and the rank argument. The vectors $p_i = (1, a_i)$ span $m + 1$ dimensions (`finrank_span_aug`). Either the evaluation map $Q \mapsto (p_j^\top Q p_j)_j$ on real matrices is surjective, so that every point has a real isolating quadric and H_c lifts by the adjugate of Lemma 3 over R (`realCert_of_isolators`); or a nonzero $\lambda \in \mathbb{R}^s$ annihilates its range, so that $\sum_i \lambda_i \langle p_i, w \rangle^2 = 0$ for every w (`isolators_or_dependence`). In the second case let $U = \{i : \lambda_i > 0\}$ and $V =$

$\{i : \lambda_i < 0\}$; both are nonempty, since $w = e_0$ gives $\sum_i \lambda_i = 0$. A linear functional vanishing on the p_i , $i \in U$, makes every term of the sum nonpositive, so it vanishes on the p_i , $i \in V$, as well; hence $\text{span}\{p_i : i \in U\} = \text{span}\{p_i : i \in V\} =: W$, of dimension $w \leq \min(|U|, |V|)$ (`rank_mem_span_of_dep`). Three distinct points of a sphere have linearly independent augmented vectors: a relation $x p_A + y p_B + z p_C = 0$ gives $x + y + z = 0$ and, paired with $A - c$, $B - c$, $C - c$, the system $y|AB|^2 + z|AC|^2 = x|AB|^2 + z|BC|^2 = x|AC|^2 + y|BC|^2 = 0$, whose determinant is $2|AB|^2|AC|^2|BC|^2 \neq 0$ (`rank_three_indep`); and $|U| = |V| = 1$ would force $a_u = a_v$, so W contains the augmented vectors of at least three distinct points, and $w \geq 3$. Every p_i lies in W or among the $s - |U| - |V|$ remaining vectors, so $m + 1 \leq w + s - |U| - |V| \leq w + m + 4 - |U| - |V|$, that is $|U| + |V| \leq w + 3$; and $|U| + |V| \geq 2w \geq w + 3$. Hence $|U| = |V| = w = 3$: the six points $S = U \cup V$ have augmented vectors spanning three dimensions, and $m + 1 = 3 + (s - 6)$ (`exists_six_of_dependence`).

The six points. The affine span of S has dimension two, so $S = t + T_2(S')$ for six distinct concyclic points $S' \subset \mathbb{R}^2$ and a linear isometry $T_2: \mathbb{R}^2 \rightarrow \mathbb{R}^D$ (the spanning representative of Section 6 with the isometry made explicit, `exists_spanning_representative`). Theorem 2 and (i), (iv), (ii) give a real certificate for S in $\mathbb{R}^D$ with a symmetric multiplied matrix H_0 of spatial block I ; the matrix $N = H_c - H_0$ is symmetric with zero spatial block and vanishes at the six points (both H_c and H_0 do), so (iii) gives a real certificate for S with multiplied matrix H_c (`realCert_six_of_planar`). Since $m + 1 = 3 + (s - 6)$, the remaining points add one dimension each, the extension lemma gives a real certificate for A with multiplied matrix H_c , and descent finishes. $\square$

The first branch is the argument of Proposition 7.3 of [1], run over $\mathbb{R} \otimes_{\mathbb{Q}} \mathbb{R}$; the congruence invariance of the criterion (a remark in Section 2 of [1]) and the descent of certificates between coefficient rings (in the proof of the equivalence proposition there) are stated for spanning sets in a fixed dimension. The extension lemma is the analogue, on the side of the criterion, of the one-point extension theorems of [3, 4] (a Ramsey set plus a point outside its affine hull is Ramsey), with the sphere carried along.

5 Algebraic coordinates

Theorem 8 (`algebraic_spherical_fieldCriterion`). *Every spherical set whose coordinates are algebraic numbers satisfies the criterion, whatever its size.*

Proof. F is a number field, so $B = F \otimes_{\mathbb{Q}} F$ is a finite étale F -algebra and the multiplication map splits: there is an idempotent $t \in B$ with $(1 \otimes x - x \otimes 1)t = 0$ for every $x \in F$ and $m(t) = 1$ (the separability idempotent; in Mathlib, `Algebra.FormallyUnramified.iff_exists_tensorProduct`). For an F -rational sphere matrix H put $P = t \cdot (H \otimes 1)$. In the sandwich, t moves every right factor to the left: $(x \otimes 1)t(H_{\alpha\beta} \otimes 1)(1 \otimes y) = t \cdot (xH_{\alpha\beta}y \otimes 1)$, so the evaluation at a_i is $t \cdot (p_i^T H p_i \otimes 1) = 0$; and $m(P) = H$. $\square$

Theorem 9 (`fieldCriterion_spherical`, `algebraic_fieldCriterion_iff_spherical`). *The criterion implies sphericity; for algebraic coordinates the criterion is exactly sphericity.*

Proof. The first is the remark after Definition 1, with the F -rational quadric read in $\mathbb{R}$: $|x - c'|^2 = |c'|^2 - C$, a sphere since a real point satisfies it. The second adds Theorem 8. $\square$

So the content of Theorems 2–7 lies in the transcendental case: for a coordinate field that is not algebraic over $\mathbb{Q}$ no separability idempotent exists (a field that is a separable $\mathbb{Q}$ -algebra is finite over $\mathbb{Q}$), though B may still have other idempotents, as for $F = \mathbb{Q}(t, \sqrt{2})$; when $\mathbb{Q}$ is algebraically closed in F , B is a domain. Every known non-Ramsey spherical set has a transcendental coordinate, as it must if (S) below holds.

6 Consequences given the classification

Write (S) for the sufficiency half of the classification: a configuration of at least two distinct points that affinely spans $\mathbb{R}^d$, $d \geq 1$, and satisfies the criterion is Ramsey. The Ramsey property passes to subsets and to congruent configurations, and any configuration of at least two distinct spherical points is congruent to one that spans a space of positive dimension and is still spherical (translate, take the span, map it isometrically; the centre projects orthogonally onto the span; `exists_spanning_representative`).

Theorem 10 (`at_most_six_spherical_ramsey_of_sufficiency`). *Given (S), every spherical set of at most six points is Ramsey, and every spherical set of at most $m+4$ points, m the dimension of its affine span, is Ramsey (`spherical_ramsey_of_sufficiency_of_card_le_finrank_add_four`).*

Proof. Sets of at most one point are Ramsey trivially. Otherwise Theorem 5, respectively Theorem 7, on the spanning representative (which has the same m), then (S) and congruence. $\square$

Theorem 11 (`algebraic_spherical_ramsey_of_sufficiency`). *Given (S), every spanning spherical set with algebraic coordinates is Ramsey; given both halves of the classification, a spanning set with algebraic coordinates is Ramsey if and only if it is spherical (the Lean name of the second statement is `algebraic_ramsey_iff_spherical_of_classification`).*

The first statement is the spherical conjecture (asked in [5], stated as a conjecture in [6]) for spanning configurations with algebraic coordinates, given (S); a non-spanning configuration with algebraic coordinates has a spanning congruent copy with algebraic coordinates (Gram–Schmidt over the real algebraic numbers), which is not formalised, so the conjecture follows for all algebraic configurations on paper. With the theorem of [5] that Ramsey sets are spherical, the second statement is the equivalence. Neither [1] nor its scope note states either.

Proposition 12. *The seven-point set $\{(0,0)\} \cup \{(j, \pm\sqrt{j(2r-j)}) : j \in \{1, 3, 4\}\}$ of [2], $r > 2$ transcendental, does not satisfy the criterion.*

Proof. Its coordinate field is $F = \mathbb{Q}(r, y_1, y_3, y_4)$ with $y_j^2 = j(2r-j)$; let D be the derivation of F with $D(r) = 1$, so $D(y_j) = j/y_j$, and $\Phi(x \otimes y) = D(x)D(y)$, which is well defined on B since it is $\mathbb{Q}$ -bilinear. Give the origin weight 2 and the pairs $j = 1, 3, 4$ weights $-2, 2, -1$. Then, exactly,

$$\sum \lambda_i \tilde{p}_i \tilde{p}_i^\top = 0, \quad \sum \lambda_i D(\tilde{p}_i) \tilde{p}_i^\top = 0, \quad \sum \lambda_i |D\tilde{p}_i|^2 = \frac{-24}{(2r-1)(2r-3)(2r-4)} \neq 0,$$

with $\tilde{p}_i = (1, a_i)$ (certificate 0002). Besides Φ and m , the $\mathbb{Q}$ -bilinear maps $\Psi_1(u \otimes v) = D(u)v$ and $\Psi_2(u \otimes v) = uD(v)$ are well defined on B , and for every $z \in B$ the Leibniz rule gives $\Phi((x \otimes 1)z(1 \otimes y)) = D(x)D(y)m(z) + D(x)y\Psi_2(z) + xD(y)\Psi_1(z) + xy\Phi(z)$ (check it on pure tensors and extend by linearity). For any P satisfying the evaluation identities, sum this against λ_i over the points and over α, β with $x = \tilde{p}_{i\alpha}$, $y = \tilde{p}_{i\beta}$, $z = P_{\alpha\beta}$: the first moment identity kills the Φ term, the second and its transpose kill the two mixed terms, and what remains is $\sum_{\alpha\beta} (\sum_i \lambda_i D(\tilde{p}_{i\alpha})D(\tilde{p}_{i\beta}))m(P_{\alpha\beta})$. With the spatial block of $m(P)$ equal to I (and $D(1) = 0$ on the constant row) this is $\sum_i \lambda_i |D\tilde{p}_i|^2 \neq 0$, while the left side is $\Phi(0) = 0$. $\square$

Proposition 13 (sharpness). *For every $d \geq 2$ there is a spherical set of $d + 5$ points whose affine span has dimension d , which fails the criterion and, by Theorem 1 of [2], is not Ramsey: the heptagon of Proposition 12 in the plane $x_3 = \dots = x_d = 0$ of $\mathbb{R}^d$, with centre $c = (r, 0, \dots, 0)$ and radius r , together with the points $c + re_j$, $3 \leq j \leq d$.*

Proof. The added points lie on the same sphere and contribute the $d - 2$ independent directions e_j , so the set has $d + 5$ points and affine dimension d ; its coordinate field is that of the heptagon, since $r = (y_1^2 + 1)/2 \in F$. If P were a certificate for it, the principal 3×3 block of P on the coordinates $0, 1, 2$ would have vanishing sandwiches at the heptagon's points, whose remaining coordinates are 0, and spatial block I_2 : a certificate for the planar heptagon over its own field, contradicting Proposition 12. The set contains the heptagon, so it is not Ramsey by Theorem 1 of [2], the Ramsey property passing to subsets. $\square$

Corollary 14. *Given (S) , and either the necessity half of the classification or Theorem 1 of [2], the least size of a non-Ramsey spherical set is seven, and for every $d \geq 2$ the least size of a non-Ramsey spherical set whose affine span has dimension d is $d + 5$.*

Proof. At most six points, or at most $d + 4$: Theorem 10. The heptagon is not Ramsey by Proposition 12 and the necessity half of the classification, or by Theorem 1 of [2] directly, and so is the set of Proposition 13. $\square$

Remark 15. The thresholds are about the existence of a failing set, not about every set of that size. Exact computations made during the development, not included here, decided the criterion over the field $\mathbb{Q}(s)$ (where $F \otimes_{\mathbb{Q}} F$ localised at the kernel of m is a discrete valuation ring) and found seven- and eight-point sets on the unit circle with stereographic parameters $c_i s + d_i$, c_i, d_i rational, that satisfy the criterion, while the nine-point and twelve-point sets of [1] fail it. The heptagon's failure rests on an algebraic relation among its coordinates; that almost every seven-point subset of a fixed circle is not Ramsey (by a derivation certificate, as for the heptagon) is Theorem A.3 of the appendix of [2], which its author marks as unverified output of a language model.

7 Prior art and provenance

As of 2026-10-07 (UTC) no statement that six concyclic points are Ramsey or satisfy the criterion, no statement of the criterion for algebraic configurations, and no positive bound depending on the dimension beyond simplices [9] was found in [1] (its text, scope note and Lean tree, whose positive results stop at five concyclic points), in [2] (whose appendix, marked by its author as unverified, treats derivation certificates: none refutes at most six points, and its non-Ramsey constructions in dimension d have $2d + 3$ points for $d \geq 2$ or $\binom{d+2}{2}$ points for $d \geq 3$), on arXiv, MathOverflow, erdosproblems.com (problem 174), the formal-conjectures repository, the Palomar registry, Hexagon, or GitHub. The one-point extension theorems [3, 4] are the nearest results. The classical positive results are the regular hexagon and all sets with a soluble transitive symmetry group [7], trapezoids [8], simplices [9] and regular polytopes [10]; the status of general cyclic quadrilaterals before 2026 and the subtransitivity conjecture are in [11].

Claude (Anthropic) was used throughout; the theorems, proofs, searches and checkers arose in that process, and the repository was written by Claude. A GPT model (OpenAI), consulted on the repository's first, eleven-theorem version, proposed Theorem 7 with an outline of its proof; the proof above and its Lean development were produced with Claude. The Lean development

was checked by the Lean 4 kernel [12] against Mathlib [13] at tag v4.35.0-rc2. The author directed the project and takes responsibility for the claims. This note is licensed CC BY-SA 4.0; the repository is MIT, with OpenAI’s Apache-2.0 notice on the restated definitions.

References

- [1] OpenAI, *A classification of finite Euclidean Ramsey configurations*, openai/math, family 172, preprint dated 23 September 2026, commit `adc7f124` (6 October 2026).
- [2] D. Pálvölgyi, *A cyclic non-Ramsey heptagon*, arXiv:2609.23327 (20 September 2026).
- [3] K. Moore, *A pyramid with a Ramsey base is Ramsey*, arXiv:2608.09649 (10 August 2026).
- [4] M. Mirabi, *One-point extensions of Euclidean Ramsey sets*, arXiv:2608.11736 (12 August 2026).
- [5] P. Erdős, R. L. Graham, P. Montgomery, B. L. Rothschild, J. Spencer, E. G. Straus, Euclidean Ramsey theorems. I, J. Combin. Theory Ser. A 14 (1973) 341–363.
- [6] R. L. Graham, Recent trends in Euclidean Ramsey theory, Discrete Math. 136 (1994) 119–127.
- [7] I. Kříž, Permutation groups in Euclidean Ramsey theory, Proc. Amer. Math. Soc. 112 (1991) 899–907.
- [8] I. Kříž, All trapezoids are Ramsey, Discrete Math. 108 (1992) 59–62.
- [9] P. Frankl, V. Rödl, A partition property of simplices in Euclidean space, J. Amer. Math. Soc. 3 (1990) 1–7.
- [10] K. Cantwell, All regular polytopes are Ramsey, J. Combin. Theory Ser. A 114 (2007) 555–562.
- [11] I. Leader, P. A. Russell, M. Walters, Transitive sets and cyclic quadrilaterals, J. Combin. 2 (2011) 457–462; Transitive sets in Euclidean Ramsey theory, J. Combin. Theory Ser. A 119 (2012) 382–396.
- [12] L. de Moura, S. Ullrich, The Lean 4 theorem prover and programming language, CADE 28, LNCS, Springer (2021) 625–635.
- [13] The mathlib Community, The Lean mathematical library, CPP 2020, 367–381.