

SLIM (Phentermine)

Brand: ProLab

Pharmaceutical Form: capsules

Presentation: Package with 50 capsules of 37.5 mg each

COMPOSITION:

Each SLIM tablet contains 37.5 mg of Phentermine, a powerful appetite suppressant that acts on the central nervous system. This compound is designed to help with weight loss, especially in patients who have a high body mass index (BMI) and who have not been successful in reducing weight through diet and physical exercise. SLIM's formulation includes excipients that optimize the stability of the drug, guaranteeing its efficacy and safety. These excipients allow for proper dissolution and absorption in the body, favoring a rapid onset of action and a prolonged duration of the effect.

THERAPEUTIC INDICATIONS:

SLIM is indicated as part of a comprehensive weight management program, which includes a low-calorie diet and a physical exercise regimen. This medication is primarily intended for people with obesity who have a BMI of 30 or more, or for those with a BMI of 27 or more and who have obesity-related medical conditions, such as hypertension, type 2 diabetes, or dyslipidemia.

Phentermine's action on the body results in a significant reduction in hunger and an improvement in the feeling of satiety, which makes it easier to adhere to a calorie-controlled eating plan. By using SLIM, patients can observe an improvement in their quality of life, as well as in their general well-being, thanks to effective and sustainable weight loss.

RECOMMENDED DOSAGE:

The usual recommended dosage of SLIM is one 37.5 mg tablet per day. It is suggested to take the tablet in the morning, approximately 30 minutes before breakfast. This morning administration allows to take full advantage of the appetite suppressant effect during the day, when patients tend to feel more hungry. It is essential to avoid taking SLIM in the afternoon or at night, as it can interfere with sleep due to its stimulant action.

In some cases, the doctor may adjust the dose according to the patient's individual response to treatment, but it is not recommended to exceed the daily dose of 37.5 mg. SLIM should be used short-term, generally for a period of 12 weeks, as prolonged use can lead to tolerance and decreased effectiveness of the drug. At the end of this cycle, it is essential to evaluate progress and determine the next step in the treatment of obesity.

MODE OF ADMINISTRATION:

SLIM is administered orally, and it is important to swallow the tablet whole with a glass of water. The tablet should not be crushed, chewed or dissolved, as this can alter its controlled release and reduce its effectiveness. To maximize the absorption and effects of the drug, it is recommended to take it on an empty stomach, 30 minutes before a meal.

It is crucial to follow the doctor or pharmacist's instructions regarding the duration of treatment and compliance with the diet and exercise plan. In addition, it is advised to maintain adequate hydration and eat balanced and nutritious meals to support the weight loss process.

CONTRAINDICATIONS:

SLIM is contraindicated in patients with known hypersensitivity to Phentermine or any of the product's excipients. It should also be avoided in people with a history of cardiovascular disease, uncontrolled hypertension, heart disease, as well as those with a history of substance abuse or significant psychiatric disorders.

The use of SLIM is also contraindicated during pregnancy and breastfeeding, as it may affect the fetus or newborn. It is also prohibited in patients with glaucoma, thyroid hyperactivity or in those who are using monoamine oxidase inhibitors (MAOIs) in the last two weeks. Women of childbearing age should use effective contraception during treatment.

Special Warnings and Precautions:

The use of SLIM requires caution in patients with pre-existing medical conditions such as diabetes, hypertension, heart disease, or psychiatric disorders. Regular monitoring of blood pressure and heart rate is essential during treatment, as Phentermine may cause increases in blood pressure and changes in heart rate.

When starting treatment with SLIM, patients should be informed about the possibility of experiencing side effects, including insomnia, nervousness, dry mouth, and mood changes. If significant or concerning adverse effects occur, it is crucial to consult with a health care professional immediately.

