

PROVI (Mesterolone)

Brand: ProLab

Pharmaceutical Form: Capsules

Presentation: Packaging with 50 capsules of 25 mg each

COMPOSITION:

Each PROVI tablet contains 25 mg of Mesterolone, a non-aromatizable androgen. This means that, unlike other anabolic steroids that can convert to estrogens and cause side effects such as gynecomastia (development of breast tissue in men), Mesterolone does not present a risk of aromatization. This characteristic makes PROVI especially useful for athletes looking to improve their muscle definition, as it allows for muscle hardening without the risk of fluid retention or increased body fat associated with estrogen. In addition, PROVI can improve the bioavailability of other hormones by optimizing the body's ability to use circulating testosterone. Its androgenic profile is moderate, making it a safe choice for those who need to increase androgen levels without exposing themselves to the more aggressive side effects of other steroids.

THERAPEUTIC INDICATIONS:

PROVI is indicated for the treatment of conditions resulting from an androgen deficiency, such as male hypogonadism, erectile dysfunction, and lack of libido. Due to its ability to enhance androgenic function without negatively affecting estrogen levels, it is also highly valued in the athletic arena. Athletes and bodybuilders use PROVI during cutting cycles, when the primary goal is to maximize muscle definition, hardness, and vascularity without the bloat or water retention commonly associated with other anabolic steroids. PROVI also acts as a natural anti-estrogenic agent, reducing side effects related to the use of aromatizing compounds, such as gynecomastia. Furthermore, due to its positive effects on libido and overall well-being, PROVI is frequently used as part of a Post-Cycle Therapy (PCT) to help restore natural testosterone levels and maintain muscle gains after a cycle.

RECOMMENDED DOSAGE:

The recommended dosage of PROVI varies considerably depending on the user's goals. In the therapeutic treatment of androgen deficiencies, the standard dosage usually ranges from 25 mg to 50 mg daily, spread over one or two daily doses. In this case, the goal is to restore adequate testosterone levels and improve symptoms related to androgen deficiency, such as fatigue, erectile dysfunction, or loss of muscle mass. For users looking to improve their athletic performance or physical appearance, the dosage can increase to a range of 50 mg to 100 mg daily, spread over two or three doses throughout the day. These higher doses help to optimize muscle hardness and improve the body's ability to metabolize other anabolic hormones that are being used simultaneously. For advanced users, who wish to maximize their results in terms of muscle definition and strength, doses can be raised up to 150 mg daily, although this should be done with caution and under medical supervision, as there may be a higher risk of negatively affecting the lipid profile or prostate function. It is not recommended to exceed this amount to avoid health risks. The cycle of use of PROVI is usually 6 to 8 weeks, and should be accompanied by an appropriate PCT to ensure full recovery of the hormonal axis.

ADMINISTRATION MODE:

PROVI should be administered orally, and it is recommended to take the tablets with an adequate amount of liquid. To optimize absorption and minimize the possibility of gastric irritation, it is preferable to ingest it with food. Healthy fats, such as those found in avocados, nuts, or natural oils, can improve the bioavailability of Mesterolone, promoting its transport and utilization by the body. Due to its relatively short half-life, it is advisable to divide the daily dose into several doses to maintain stable levels of the compound in the blood throughout the day. This not only optimizes its effectiveness in terms of anabolism, but also minimizes the risk of unwanted spikes in androgenic levels, which could trigger adverse effects such as aggression or mood swings. Users who train intensely should schedule their doses so that one of them is taken approximately 30 to 60 minutes before physical activity, as this could improve strength and endurance during exercise.

CONTRAINDICATIONS:

PROVI is contraindicated in patients with known hypersensitivity to Mesterolone or any of the components of its formulation. It should not be used in individuals with a history of severe liver disease, such as hepatitis, cirrhosis or any other active liver pathology, as it may affect hepatic metabolism. It is also contraindicated in individuals with prostate or breast cancer, especially in men, as it could exacerbate the progression of these conditions due to its action on androgen receptors. Its use should also be avoided in patients with severe hypertension or uncontrolled cardiovascular disease, as androgens may influence the lipid profile and increase the risk of complications such as atherosclerosis or heart failure. PROVI should not be used in women, especially during pregnancy or lactation, due to the risk of masculinization of the fetus or serious hormonal alterations in the infant. In addition, regular monitoring of liver and prostate status is recommended in users who use PROVI for prolonged periods or at high doses to avoid long-term complications.

