

PRIMAX^{PRO}

PRIMAX (Methenolone Enanthate)

Brand: ProLab

Pharmaceutical Form: Injectable Solution

Presentation: 100ml/mg concentrated vial of 10 ml

COMPOSITION:

Each ml of Primax contains 100 mg of methenolone enanthate, an anabolic steroid known to be mild but highly effective for preserving muscle mass and improving definition in cutting cycles. Methenolone enanthate has a long half-life, making it an excellent choice for multi-week cycles, where the main goal is to preserve muscle while reducing body fat percentage.

Primax is recognized for its low androgenic properties, making it suitable for both men and women looking to improve muscle definition without the risk of side effects associated with more potent steroids. This compound is highly valued in the bodybuilding community for its ability to provide quality gains without causing excessive water retention, making it an ideal choice for pre-contest periods or cutting phases.

THERAPEUTIC INDICATIONS:

Primax is indicated for those looking to preserve lean muscle mass during calorie-restrictive diets or cutting cycles. It is also used to improve strength and physical endurance in athletes, providing a more defined and toned appearance without the harsh androgenic side effects of other steroids. It is ideal for those looking to minimize water and fat retention while maximizing the development of quality muscle.

Because of its mild profile, Primax is frequently used by bodybuilders and athletes who wish to keep their testosterone levels in a healthy range without resorting to more aggressive steroids. Methenolone enanthate offers a gradual release, maintaining stable levels of the hormone in the bloodstream, resulting in a more controlled experience and lower risk of unwanted side effects such as gynecomastia or water retention.

RECOMMENDED DOSAGE:

The recommended dosage of Primax varies depending on the user's experience and cycle goals. For beginners, a dosage of 400 mg per week is suggested, spread over one or two applications to ensure a steady release of the compound. In more advanced users, dosages can be increased up to 600-800 mg per week, although it is important to remember that higher doses also increase the risk of side effects, such as suppression of endogenous testosterone production.

For women, PRIMAX is a safe choice due to its low androgenic activity. The recommended dosage for women ranges between 50-100 mg per week, allowing for improved muscle definition and tone without causing virilization or other side effects associated with the use of more potent steroids.

PRIMAX cycles typically last between 8-12 weeks. Since Methenolone Enanthate is a slow-acting compound, it is important to give it time to kick in and start showing visible results. At the end of the cycle, it is recommended to perform a proper Post Cycle Therapy (PCT) to help regain natural hormone production and minimize the loss of muscle gains.

MODE OF ADMINISTRATION:

PRIMAX is administered via deep intramuscular injection, preferably in areas such as the glutes or quads. Since its half-life is approximately 7-10 days, it is only necessary to apply the product once or twice a week to maintain stable blood levels. Alternating injection sites is key to avoid local irritation or infection.

This dosing pattern makes Primax a convenient choice for those who want a long cycle without the need for frequent injections. Additionally, due to its slow-release formula, it is ideal for cutting or maintenance cycles where muscle quality and definition are a priority.

CONTRAINDICATIONS:

Primax is contraindicated in individuals with hypersensitivity to methenolone or any of the excipients present in the formulation. It should not be used in people with breast or prostate cancer, or in those with severe liver disease. It is also contraindicated in patients with uncontrolled heart or kidney failure, as well as in people with a predisposition to cardiovascular problems or stroke.

The use of Primax in pregnant or lactating women is strictly prohibited. Although Primax has low androgenic activity, the risk of virilization remains, so women should exercise caution when using this compound and adjust doses as necessary to avoid adverse effects.

