



SLIM MAX
Brand: ProLab
Pharmaceutical Form: Capsules
Presentation: Package with 50 capsules of 850 mg each

COMPOSITION:

Each Slim Max tablet contains 850 mg of Metformin, a widely used drug for the treatment of type 2 diabetes due to its ability to improve insulin sensitivity and reduce blood glucose levels. Metformin works by decreasing the production of glucose in the liver, reducing its absorption in the intestine and improving the uptake and use of glucose by peripheral tissues.

In addition to its use in patients with diabetes, Slim Max has been used in weight loss treatments due to its ability to reduce insulin resistance, improve fat burning and reduce calorie absorption, making it an ideal option for people looking for support in managing their weight.

THERAPEUTIC INDICATIONS:

Slim Max is indicated for the treatment of type 2 diabetes, helping to control blood glucose levels when diet and exercise are not enough. In addition, it has been used in weight loss protocols for people with insulin resistance or in whom excess body fat interferes with proper blood sugar regulation.

Due to its action in improving insulin sensitivity, Slim Max has also been used in the treatment of conditions such as polycystic ovary syndrome (PCOS), where more effective hormonal regulation and an improvement in metabolic processes related to weight control and fertility are sought.

RECOMMENDED DOSAGE:

The standard dosage of Slim Max for the treatment of type 2 diabetes in adults is 850 mg once or twice a day, preferably administered with meals to minimize gastrointestinal side effects. For people using Slim Max as a support in weight loss or in the treatment of other metabolic conditions, doses may be adjusted under medical supervision.

It is recommended to start with a low dose of 850 mg per day to assess the patient's tolerance and then gradually increase the dose to a maximum of 2550 mg daily (divided into three doses) if necessary and well tolerated.

It is essential that the administration of Slim Max be accompanied by an adequate diet and a physical exercise regimen to maximize its effects. In people seeking weight loss, constant monitoring of glucose levels and periodic evaluation of progress are essential to adjust the dose or duration of treatment.

ADMINISTRATION MODE:

Slim Max should be administered orally with a glass of water and preferably with meals to reduce the possibility of gastrointestinal discomfort. The tablets should not be chewed or crushed, but swallowed whole to ensure a controlled release of the medication.

In case a dose is missed, it is advisable to take it as soon as remembered, but if it is close to the next dose, the missed dose should be skipped to avoid overdose.

CONTRAINDICATIONS:

The use of Slim Max is contraindicated in people with known hypersensitivity to metformin or any of the components of the formula. It should also not be used in patients with severe renal insufficiency (creatinine clearance less than 30 ml/min) due to the risk of accumulation of the drug and the possible appearance of lactic acidosis, a rare but serious complication.

Slim Max is contraindicated in people with uncontrolled heart failure, severe liver problems, or in those who suffer from diseases that can cause hypoxia, such as acute myocardial infarction or shock. Its use is not recommended in people who consume large amounts of alcohol, as this may increase the risk of lactic acidosis.

