

BOLDEX[®]PRO

BOLDEX (Boldenone Undecylenate)

Brand: ProLab

Pharmaceutical Form: Injectable Solution

Presentation: 200ml/mg concentrated vial of 10 ml

COMPOSITION:

Each ml of Boldex contains 200 mg of boldenone undecylenate, a very popular anabolic steroid among athletes and bodybuilders due to its ability to improve lean muscle growth and increase strength without excessive fluid retention. Its long half-life allows for longer intervals between injections, making it a convenient option for long cycles.

Boldenone is known for its balanced effect between anabolic and androgenic properties, making it useful for both bulking and cutting cycles. In addition, Boldex promotes a gradual increase in muscle mass, with less risk of water and fat retention compared to other steroids.

THERAPEUTIC INDICATIONS:

Boldex is indicated for those seeking a sustained increase in lean muscle mass with minimal water retention, ideal for both clean bulking and cutting phases. Due to its ability to stimulate protein synthesis and increase appetite, it is widely used in long-term cycles to build quality muscle mass.

Boldenone is also known for its ability to enhance red blood cell production, which increases oxygenation of the muscles and therefore improves endurance during intense workouts. This added benefit makes Boldex especially useful for athletes who practice endurance sports or bodybuilders who wish to maintain high levels of intensity in the gym.

RECOMMENDED DOSAGE:

The recommended dosage of Boldex for male users ranges from 300-600 mg per week. Due to the nature of its undecylenate ester, injections can be spaced out every 7-10 days, making it easier to use in longer cycles. Standard beginner dosages start at 300-400mg per week, while advanced users can increase the dosage to maximize muscle gains.

For women, the dosage should be much lower, typically between 50-100mg per week, due to the risk of androgenic side effects such as virilization. Women using Boldex should pay attention to any signs of virilization, such as body hair growth or voice changes, and adjust dosages accordingly.

Boldex cycles typically last between 10-12 weeks, due to its gradual and sustained action. It is commonly used in combination with other steroids, such as testosterone or winstrol, to maximize gains and reduce side effects. At the end of the cycle, it is advisable to perform a proper Post Cycle Therapy (PCT) to restore natural hormone levels.

MODE OF ADMINISTRATION:

Boldex is administered by deep intramuscular injection, usually in the glutes or quadriceps. Since Boldenone undecylenate has a long half-life, injections can be administered weekly or every 10 days. This feature makes it attractive to users who prefer a less frequent injection regimen, but still desire sustained and stable results.

The compound is released slowly into the body, ensuring consistent levels of Boldenone in the bloodstream over a long period of time. While this makes it easier to administer, users should keep in mind that Boldex requires longer cycles to achieve maximum benefits, as results are not as immediate as with other steroids.

CONTRAINDICATIONS:

Boldex is contraindicated in individuals with hypersensitivity to Boldenone or any component of the formula. It should not be used in individuals with a history of severe liver disease, prostate or breast cancer, as well as those with severe cardiovascular disorders. It is also not recommended for use in pregnant or breastfeeding women, due to the risk of androgenic effects on the fetus or infant.

Users with a genetic predisposition to baldness or prostate problems should take special caution when using Boldex, as although its androgenic profile is relatively low, it is still capable of causing these side effects in some susceptible individuals.

