



Oral toxicity prediction results for input compound




Predicted LD50: 2500mg/kg

Predicted Toxicity Class: 5



Average similarity: 42.04%

Prediction accuracy: 54.26%



Name	C[C@@]1(C(=O)N2CCc3c4cccc4)cc3C2)CCNC(=O)C1
Molweight	338.4
Number of hydrogen bond acceptors	4
Number of hydrogen bond donors	1
Number of atoms	25
Number of bonds	28
Number of rotatable bonds	3
Molecular refractivity	101.97
Topological Polar Surface Area	62.55
octanol/water partition coefficient(logP)	3.01

Toxicity Model Report

[Copy](#) [Excel](#) [CSV](#) [PDF](#)

Classification	Target	Shorthand	Prediction	Probability
Organ toxicity	Hepatotoxicity	dili	Inactive	0.85
Organ toxicity	Neurotoxicity	neuro	Active	0.78
Organ toxicity	Nephrotoxicity	nephro	Inactive	0.77
Organ toxicity	Respiratory toxicity	respi	Active	0.86
Organ toxicity	Cardiotoxicity	cardio	Inactive	0.91
Toxicity end points	Carcinogenicity	carcino	Inactive	0.54
Toxicity end points	Immunotoxicity	immuno	Inactive	0.97
Toxicity end points	Mutagenicity	mutagen	Inactive	0.55
Toxicity end points	Cytotoxicity	cyto	Inactive	0.64
Toxicity end points	BBB-barrier	bbb	Active	0.90
Toxicity end points	Ecotoxicity	eco	Inactive	0.51
Toxicity end points	Clinical toxicity	clinical	Active	0.69
Toxicity end points	Nutritional toxicity	nutri	Inactive	0.59
Tox21-Nuclear receptor signalling pathways	Aryl hydrocarbon Receptor (AhR)	nr_ahr	Inactive	0.87
Tox21-Nuclear receptor signalling pathways	Androgen Receptor (AR)	nr_ar	Inactive	0.95
Tox21-Nuclear receptor signalling pathways	Androgen Receptor Ligand Binding Domain (AR-LBD)	nr_ar_lbd	Inactive	0.98
Tox21-Nuclear receptor signalling pathways	Aromatase	nr_aromatase	Inactive	0.84
Tox21-Nuclear receptor signalling pathways	Estrogen Receptor Alpha (ER)	nr_er	Inactive	0.81
Tox21-Nuclear receptor signalling pathways	Estrogen Receptor Ligand Binding Domain (ER-LBD)	nr_er_lbd	Inactive	0.97
Tox21-Nuclear receptor signalling pathways	Peroxisome Proliferator Activated Receptor Gamma (PPAR-Gamma)	nr_ppar_gamma	Inactive	0.98
Tox21-Stress response pathways	Nuclear factor (erythroid-derived 2)-like 2/antioxidant responsive element (nrf2/ARE)	sr_are	Inactive	0.97
Tox21-Stress response pathways	Heat shock factor response element (HSE)	sr_hse	Inactive	0.97
Tox21-Stress response pathways	Mitochondrial Membrane Potential (MMP)	sr_mmp	Inactive	0.86
Tox21-Stress response pathways	Phosphoprotein (Tumor Suppressor) p53	sr_p53	Inactive	0.85
Tox21-Stress response pathways	ATPase family AAA domain-containing, protein 5 (ATAD5)	sr_atad5	Inactive	0.94
Molecular Initiating Events	Thyroid hormone receptor alpha (THRα)	mie_thr_alpha	Inactive	0.9
Molecular Initiating Events	Thyroid hormone receptor beta (THRβ)	mie_thr_beta	Inactive	0.87
Molecular Initiating Events	Transthyretin (TTR)	mie_ttr	Inactive	0.96
Molecular Initiating Events	Ryanodine receptor (RYR)	mie_ryr	Inactive	0.86

Classification	Target	Shorthand	Prediction	Probability
Molecular Initiating Events	<u>GABA receptor (GABAR)</u>	mie_gabar	Active	0.57
Molecular Initiating Events	<u>Glutamate N-methyl-D-aspartate receptor (NMDAR)</u>	mie_nmdar	Inactive	0.85
Molecular Initiating Events	<u>alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate receptor (AMPA)</u>	mie_ampar	Inactive	0.99
Molecular Initiating Events	<u>Kainate receptor (KAR)</u>	mie_kar	Inactive	1.0
Molecular Initiating Events	<u>Achetylcholinesterase (AChE)</u>	mie_ache	Active	0.58
Molecular Initiating Events	<u>Constitutive androstane receptor (CAR)</u>	mie_car	Inactive	0.99
Molecular Initiating Events	<u>Pregnane X receptor (PXR)</u>	mie_pxr	Inactive	0.54
Molecular Initiating Events	<u>NADH-quinone oxidoreductase (NADHox)</u>	mie_nadhox	Inactive	0.96
Molecular Initiating Events	<u>Voltage-gated sodium channel (VGSC)</u>	mie_vgsc	Inactive	0.76
Molecular Initiating Events	<u>Na⁺/I⁻ symporter (NIS)</u>	mie_nis	Inactive	0.86
Metabolism	<u>Cytochrome CYP1A2</u>	CYP1A2	Inactive	0.95
Metabolism	<u>Cytochrome CYP2C19</u>	CYP2C19	Inactive	0.79
Metabolism	<u>Cytochrome CYP2C9</u>	CYP2C9	Inactive	0.70
Metabolism	<u>Cytochrome CYP2D6</u>	CYP2D6	Active	0.52
Metabolism	<u>Cytochrome CYP3A4</u>	CYP3A4	Inactive	0.64
Metabolism	<u>Cytochrome CYP2E1</u>	CYP2E1	Inactive	0.99

Toxicity targets

Possible binding to toxicity targets is shown below. For more information on the targets, please click on the individual abbreviations.



<u>AA2AR</u>	<u>ADRB2</u>	<u>ANDR</u>	<u>AOFA</u>	<u>CRFR1</u>	<u>DRD3</u>	<u>ESR1</u>	<u>ESR2</u>	<u>GCR</u>	<u>HRH1</u>	<u>NR12</u>	<u>OPRK</u>	<u>OPRM</u>	<u>PDE4D</u>	<u>PGH1</u>	<u>PRGR</u>