

Axiom 1.1.1 : Properties of Equivalence

The following is assumed to be true for all α, β , and γ in the real numbers.

$$\forall \alpha, \beta, \gamma \in \mathbb{R}$$

An element α is equal to itself, this property is called reflexivity. If an element α is equal to an element β then β is also equal to α , this property is called symmetry. If an element α is equivalent to an element β , and β is also equivalent to γ then α is equivalent to γ , this property is called transitivity.

- (i) $\alpha = \alpha$, (reflexivity)
- (ii) $\alpha = \beta \Rightarrow \beta = \alpha$, (symmetry)
- (iii) $\alpha = \beta \wedge \beta = \gamma \Rightarrow \alpha = \gamma$, (transitivity)

Remark 1.1.2 : Well Defined Operation

An operation is said to be well defined if given the same inputs, the operation will always give the same outputs. Or symbolically,

$$\forall \alpha, \beta \in \mathbb{R}, f \text{ is an operation, } \alpha = \beta \Rightarrow f[\alpha] = f[\beta]$$

at the moment we will take for granted the fact that the normal operations of numbers is well defined, that is addition, subtraction, multiplication, and division of 2 numbers.

Example 1.1.3 : Isolate x

We can apply the remark about well defined operations to manipulate our equations in the following manner.

$$x + 2 = 3 \tag{1}$$

$$x + 2 - 2 = 3 - 2 \tag{2}$$

$$x = 1 \tag{3}$$

Remark 1.1.4 : Interpretation of Well Defined Operation in Terms of Equation Manipulation

For those that are not convinced by the manipulation above, or how the definition of a well defined operation is related to it. Another way to interpret a well defined operation is as an operation where if you apply the operation to a single number, you should always get the same result.

In step (1) we know that we have two equivalent objects $x + 2$ and 3, the well defined operation that we apply to the two equivalent objects is "minus 2". Since minus is well defined, we conclude that $x + 2 - 2$ is equivalent to $3 - 2$, we collect the terms and conclude that x is equivalent to 1.

Example 1.1.5 : Isolate x

$$\begin{aligned} 2x &= 6 \\ \frac{1}{2} \cdot 2x &= \frac{1}{2} \cdot 6 \\ x &= 3 \end{aligned}$$

Example 1.1.6 : Isolate x

$$\begin{aligned} \frac{x}{3} &= 4 \\ \frac{x}{3} \cdot 3 &= 4 \cdot 3 \\ x &= 12 \end{aligned}$$

Example 1.1.7 : Isolate y

$$\begin{aligned} \frac{y+2}{3} &= 4x \\ 3 \cdot \frac{y+2}{3} &= 3 \cdot 4x \\ y+2 &= 12x \\ y &= 12x - 2 \end{aligned}$$

Example 1.1.8 : Isolate y

$$\begin{aligned} \frac{3-2y}{5} - y &= x \\ \frac{3-2y}{5} &= x + y \\ 3-2y &= 5x + 5y \\ 3-5x &= 7y \\ y &= \frac{3-5x}{7} \end{aligned}$$

Remark 1.1.9 : Simplest Form

When we are dealing with equations with more than one variable, isolating one variable will result creating an equation in the form of a variable written in terms of another variable.

$$y = 2x + 3, x = 15$$

However that form is not simple like a single number, there is nothing wrong with that. It just happens that the simpler form cannot contain all of the information of the original equation.

Axiom 1.1.10 : Distribution

For α, β, γ the reals, left and right multiplication is distributed over addition.

$$\forall \alpha, \beta, \gamma \in \mathbb{R}$$

$$\alpha(\beta + \gamma) = \alpha\beta + \alpha\gamma$$

Remark 1.1.11 : Geometric Demonstration of Distribution * this is not proof
If we accept that the the area of a rectangle is its length multiplied by its width, and that its area is also the sum of the parts its made of, then it follows that multiplication is distributed over addition.

illustration here

Area = $\alpha(\beta + \gamma)$, length by width

Area = $\alpha\beta + \alpha\gamma$, sum of area of parts

Remark 1.1.12 : Number Line (optional)

A number line is a visual tool that is able to visually represent a variable that takes on some value. It consists of a single axis, with regularly spaced markings of numbers that increase towards the right with a point on the marking on which its the variable's value.

Example 1.1.13 : $x = 4$

insert illustration here

Remark 1.1.14 : Cartesian Plane (optional)

A Cartesian plane is an extended version of a number line that is able to visually represent two variables which take on some value. It consists of two perpendicular axes, both with markings like the number line. There is a point on the intersection of the two markings that take the values of the two variables intersect.

Example 1.1.15 : $x = 3 \wedge y = 4$

Often abbreviated as $(x, y) = (3, 4)$

insert illustration here

Remark 1.1.16 : Linear Equation

A linear equation is an equation of the form $\alpha x + \beta y + \gamma = 0$, where α, β, γ are

in the reals.

Remark 1.1.17 : Slope Intercept Form of a Linear Equation

From the original form of the linear equation, we can derive the following alternate form of a linear equation.

$$\begin{aligned}\alpha x + \beta y + \gamma &= 0 \\ \beta y &= -\alpha x - \gamma \\ y &= -\frac{\alpha}{\beta}x - \frac{\gamma}{\beta}\end{aligned}$$

Suppose we substitute $\sigma = -\frac{\alpha}{\beta}, \iota = -\frac{\gamma}{\beta}$, then we can rewrite our equation as

$$y = \sigma x + \iota$$

Remark 1.1.18 : Linear Equation Constructor (optional)

Consider a constructor for linear equations

$$\begin{aligned}\texttt{LinearEquation} : \mathbb{R}^3 &\mapsto \mathbf{Equation} \\ (\alpha, \beta, \gamma) &\rightarrow \alpha x + \beta y + \gamma = 0\end{aligned}$$

which is able to construct linear equations of standard form, the following alternate constructor is able to construct linear equation of slope intercept form

$$\begin{aligned}\mathbf{LinearEquation} : \mathbb{R}^2 &\mapsto \mathbf{Equation} \\ (\sigma, \iota) &\rightarrow y = \sigma x + \iota\end{aligned}$$

Remark 1.1.19 : Substitution

In an equation of more than one variables, we can suppose without proof that one or more variables takes on a concrete value in order to find the implied values of other variables. Consider the following equation

$$\alpha = \frac{\beta + \gamma + 5}{2}$$

from this equation, we can only find the value of variables in terms of other variables. Suppose that $\gamma = 2\beta$, then

$$\alpha = \frac{3\beta}{5}$$

Suppose that $\beta = 5$, then

$$\alpha = 3$$

So it turns out that in our original equation, if we suppose that $\gamma = 2\beta \wedge \beta = 5$, then $\alpha = 3$

This technique will become useful when we try to understand why equations of a certain form are called linear equations.

Remark 1.1.20 : Linear Equation Represented by Cartesian Plane

On a cartesian plane, a point requires the value of two variables. A linear equation is an equation of two variables, and thing to note about equations of two variables is that if you know the value of one variable, you can find out the value of the other.

Suppose: $\text{LinearEquation}(\alpha = -1, \beta = 1, \gamma = 0)$ which gives $-1 \cdot x + 1 \cdot y + 0 = 0$, is our equation

$$\text{Infer: } y = -\frac{-1}{1} \cdot x - \frac{0}{1}$$

Infer: $y = x$, is the slope intercept form of our equation

Suppose: $x = 0$

Infer: $y = 0$

Suppose: $x = 2$

Infer: $y = 2$

Suppose: $x = 2.5$

Infer: $y = 2.5$

using this method we can generate the pairs of numbers required to plot on the cartesian plane, see the pairs generated using this equation create pairs that fall on a line, hence the name linear equation

Remark 1.1.21 : Slope

Slope is a measure of how steep the a linear equation is, defined as increase in y for each increase in x

* increase y by 1 unit, see how much x increases * take 2 points (why does this work) * look at σ

Remark 1.1.22 : Intercept

where the graph intercepts the axis, y intecept is in the equation, how to find x intercept?

Remark 1.1.23 : Standard Form of Linear Equation