M328K Homework 8

Joshua Dong

March 21, 2014

0.1 4.4.6

Find all solutions of $x^8 - x^4 + 1001 \equiv 0 \pmod{539}$.

$$\begin{array}{l} x^8-x^4+462\equiv 0\pmod{7^211}.\\ \therefore x^8-x^4+462\equiv 0\pmod{7^2} \text{ and } x^8-x^4+462\equiv 0\pmod{11}\\ \text{We first find solutions to the first congruence.}\\ x^8-x^4+21\equiv 0\pmod{7^2} \end{array}$$

We now find solutions to the second congruence.

$$\begin{array}{l} x^8 - x^4 + 0 \equiv 0 \pmod{11} \\ x^4 (x^4 - 1) \equiv 0 \pmod{11} \end{array}$$

0.2 4.3.20.b

Find $x \in \mathbb{Z}$ such that

0.3 4.3.20.e

```
Find x \in \mathbb{Z} such that x \equiv 7 \pmod{9}, x \equiv 2 \pmod{10}, x \equiv 3 \pmod{12}, and x \equiv 6 \pmod{15}.
```

We derive an equivalent, minimal set of congruences using the Chinese remainder theorem.

```
\begin{array}{l} x \equiv 7 \pmod{9} \\ x \equiv 2 \pmod{2} \\ x \equiv 2 \pmod{5} \\ x \equiv 3 \pmod{5} \\ x \equiv 3 \pmod{4} \\ x \equiv 6 \pmod{5} \\ \text{These reduce to:} \\ x \equiv 0 \pmod{5} \\ \text{These reduce to:} \\ x \equiv 0 \pmod{2}, \\ x \equiv 0 \pmod{3}, \\ x \equiv 0 \pmod{4}, \\ x \equiv 1 \pmod{5}, \\ x \equiv 2 \pmod{5}, \\ x \equiv 7 \pmod{9}. \end{array}
```

There are no solutions to this contradictory set of congruences. (The intersection of the system is the null set).

$0.4 \quad 4.3.32$

Show that the system

```
x \equiv 1 \pmod{2},
x \equiv 0 \pmod{4},
x \equiv 0 \pmod{3},
x \equiv 2 \pmod{12},
x \equiv 2 \pmod{8},
x \equiv 22 \pmod{24}
is a covering set of congruences.
We derive an equivalent set of congruences using the Chinese remainder theo-
rem.
x \equiv 1 \pmod{2},
x \equiv 0 \pmod{4},
x \equiv 0 \pmod{3},
x \equiv 2 \pmod{3},
x \equiv 2 \pmod{4},
x \equiv 2 \pmod{8},
x \equiv 22 \pmod{3},
x \equiv 22 \pmod{8}.
These reduce to:
x \equiv 1 \pmod{2},
x \equiv 0 \pmod{3},
x \equiv 1 \pmod{3},
x \equiv 2 \pmod{3},
x \equiv 0 \pmod{4},
x \equiv 2 \pmod{4},
x \equiv 2 \pmod{8},
x \equiv 6 \pmod{8}.
The union of:
x \equiv 0 \pmod{3},
x \equiv 1 \pmod{3},
x \equiv 2 \pmod{3}
is U, the universal set of all integers.
Therefore the system is a covering set of congruences.
```

$0.5 \quad 4.3.36$

 $x^2 + 6x - 31 \equiv 0 \pmod{2^3}$.

```
Find all solutions of the congruence x^2 + 6x - 31 \equiv 0 \pmod{2^3 3^2}.
```

```
x^2 + 6x + 9 \equiv 0 \pmod{8}
(x+3)^2 \equiv 0 \pmod{8}.
\therefore x \equiv 1 \pmod{8} or
x \equiv 5 \pmod{8}.
These are equivalent to the congruence x \equiv 1 \pmod{4}.
x^2 + 6x - 31 \equiv 0 \pmod{3^2}.
x^2 - 3x - 4 \equiv 0 \pmod{9}
(x-4)(x+1) \equiv 0 \pmod{9}.
\therefore x \equiv 4 \pmod{9} or
x \equiv 8 \pmod{9}.
We want the intersection of the modulo 8 and modulo 9 solutions:
x \equiv 1 \pmod{4} and
x \equiv 4 \pmod{9} or x \equiv 8 \pmod{9}.
This is equivalent to:
x \equiv 1 \pmod{4} and x \equiv 4 \pmod{9} or
x \equiv 1 \pmod{4} and x \equiv 8 \pmod{9}.
Now we solve the systems of congruences.
x \equiv 1 \pmod{4},
x \equiv 4 \pmod{9}.
Using the first congruence,
Let q \in \mathbb{Z} where x = 4q + 1.
Substituting into the second congruence,
4q + 1 \equiv 4 \pmod{9}.
It follows that q \equiv (\bar{4})(3) \equiv 3 \pmod{9}.
Let r \in \mathbb{Z} where q = 9r + 3.
Now we have x = 4(9r + 3) + 1 = 36r + 13.
x \equiv 1 \pmod{4},
x \equiv 8 \pmod{9}.
Using the first congruence,
Let q \in \mathbb{Z} where x = 4q + 1.
Substituting into the second congruence,
4q + 1 \equiv 8 \pmod{9}.
It follows that q \equiv (\bar{4})(7) \equiv 4 \pmod{9}.
Let r \in \mathbb{Z} where q = 9r + 4.
```

Now we have x = 4(9r + 4) + 1 = 36r + 17.