Mode of Application:

Lime-Broadcast agricultural lime one month before planting

ALFALFA OR LUCERNA/Lespedeza

1st appl.: Broadcast the fertilizers including 2/3 of recommended phosphate and potash prepared.
2nd appl.: Topdress with remaining phosphate and potash fertilizers for maintenance.

ALGAE (Fishpond) or Lab-Lab

1st appl.: Drain the pond and allow to dry for 2-3 weeks. Broadcast the fertilizer including ½ nitrogen and fill with water to a depth of 10 cm.

2nd appl.: Broadcast remaining nitrogenous fertilizer at a month's interval and increase water to a depth of 25-30 cm as algae grows thicker

Ampalaya

1st appl.: Mix ½ of recommended fertilizer with the soil at planting time
2nd appl.: Sidedress with remaining fertilizer when 1st set of fruits begin to form

Apple

Potash and phosphate fertilizers are applied during period. Two-thirds (2/3) N fertilizer is applied before rainy season and one-third (1/3) N in summer.

Arrow Root

Organic fertilizer should be applied during or before the last harrowing so that it will be incorporated into the soil. ½ N and all P & K will be applied before planting. The remaining ½ N will be applied 3 months after planting.

Asparagus

The fertilizers are broadcast before planting and worked into a depth of about 7.5cm. During the cutting season, N fertilizer is scattered along the rows at intervals of about 14 days.

Atis/Avocado/Atsuete

Work into the soil recommended amounts of fertilizer applied around the root zone. Repeat this process every 3 months or 4 times a year.

Bamboo

Apply the compost or fertilizer 2-3 times per year

Banana

Apply recommended fertilizer in four splits in equal amount.
Apply the recommended fertilizer every three (3) months to ensure proper growth.

1st appl.: At Planting
2nd appl.: 3 month after transplanting
3rd appl.: 6 months after transplanting
4th appl.: 9 or just before or at initial flowering

At planting, apply the fertilizer around the planting hole at 30-60 cm from the pseudo stem in ring or by band placement.

On slopes the fertilizer must be forked-in on the uphill side of the plant.

Basil

Apply the organic fertilizer at least 1 month prior to planting. Weeks after planting, when plants developed at least two pairs of true leaves, start application of complete fertilizer (14-14-14) and this should be applied once a week to maintain vigorous leaf production.

Batao/string beans/lima beans/winged beans/patani/Upo/Patola

Apply in band 2-3 inches to the side and 1-2 inches below the level of the seeds or apply the fertilizer as soon as the vines reach a height of one meter.

Beans (Baguio or snap beans/cowpea/bush sitao)

Apply ½ N and all P and K fertilizer in band 2-3 inches to the side and 1-2 inches below the level of the seed. Apply the remaining nitrogen one month after planting.

Black Pepper

Seedlings: Apply the recommended amount of organic fertilizer at the base of the plant before transplanting.

For established plant:
Apply the fertilizer including ½ of the Nitrogen at least 6 months after transplanting. Sidedress the remaining nitrogen fertilizer when the 1st set of fruits occur.
Repeat application yearly throughout the growing period of the crop.

Brocolli/Cauliflower

1st appl.: Apply the fertilizer including ½ of the nitrogen in the form of a double band 10-12.5 cm on each side of the row at 10 cm deep at planting
2nd appl.: Topdress with remaining nitrogen fertilizers 10-14 days after planting

Bougainvilla

Apply the recommended fertilizer monthly especially at the start
of the flowering months (March-May) and (Sept-Nov.) to encourage flowering

After each blooming cycle (approximately 2-3months), trim or pinch the soft
tips of the young plants to create multiple offshoots and thicken the plants
appearance and yield a more colorful bougainvilla.

Cabbage

1st appl.: Apply the fertilizer including ½ nitrogen as a double band 10-12.5cm on both sides of the row at 10cm deep at planting
2nd appl.: Sidedress with remaining nitrogen fertilizer 10-14 days after planting and/or a part of this may be applied 2 weeks before harvest.

Cacao

For seedlings, mix top soil with complete fertilizer and transplant to a hole bored 3-50cm deep made 3-4 cm larger than the ball of earth around the roots of the seedlings.

For established crops, mix the recommended fertilizers and apply in 2 equal doses: 1st at the start of the rainy season and the 2nd before the end of the rainy season. Application is done in 5-6 holes bored 5 cm deep around the base of the plant

Caimito (Star Apple)/Duhat/Sampaloc

At planting time apply fertilizer 4-5 inches deep and 3 inches to the side of the plant.

For established plant, apply fertilizer in borings 5-6 inches deep around the base in 2 equal doses: 1st at start of rainy
season and 2nd before the end of rainy season

Camote (Sweet Potato)

Apply all the fertilizers in band 3-5 inches to the side along the row at a depth of 4-6 inches at the time of transplanting.
Note: Avoid using too heavy dressings of nitrogen as this would produce badly shaped, over-large tubers and more leaf-tops.

Cashew/Pilinut/Chico

At planting time, apply fertilizer 3-4 inches below and to the side of the plant.

For established plant, apply fertilizer uniformly around the base or in borings 3-4 inches deep at outside half of area shaded by canopy.

Carrots

1st appl: Broadcast and thoroughly incorporate into the soil to about 10 cm deep just before seeding.
2nd appl.: Topdress after seedling have taken roots.

Cassava
1st appl.: Apply the phosphorus and potassium fertilizers including ½ of nitrogen at planting.
2nd appl.: Sidedress with remaining nitrogen fertilizer 2-3 months after planting during hilling up.

Castor oil beans

1st appl.: Apply the fertilizer 3 inches to the side 2-3 inches deep in the row at the time of planting.
2nd appl.: Sidedress with ½ N when fruits begin to form.

Celery/Turnip(Singkamas)

One-half (1/2) to two-thirds (2/3) of the recommended fertilizer is broadcast and plowed into the soil during land preparation. Sidedress with the remaining fertilizer during the vegetative period.

Chayote

Mix the fertilizers including 2/3 of the nitrogen with the soil at planting time.
Sidedress with the remaining nitrogen fertilizer when fruits begin to form.

Chico

At planting time, apply fertilizer 3-4 inches below and to the side of the plant.

For established plant, apply fertilizer uniformly around the base or in borings 3-4 inches deep at outside half of area shaded by canopy.

Chili or Bell Pepper

Apply the fertilizer including 2/3 of the nitrogen during the last operation in the preparation of the land, preferably along the line of the rows. Sidedress the remaining nitrogenous fertilizer when 1st set of fruits occur.

Sili-Labuyo

Apply the organic fertilizer during the last operation in the preparation of the land preferably along the line of the rows.

Sidedress the recommended inorganic fertilizer including 1/2 of nitrogen one month after planting or transplanting

Second sidedressing is made immediately after the 1st picking/harvesting of marketable fruits using the remaining nitrogen.

Cilantro

Apply all the P & K fertilizer including one half of N before planting.
Apply the remaining N fertilizer once the plants are already established or about 5-6 weeks from planting.

Citrus (Macademia/Lemon/Lime or Dayap/Orange /Pomelo)

Seedlings:

Incorporate fertilizers with the soil to a depth of 6-10 inches before planting together with the organic fertilizer.

For Established plant:

Apply NPK fertilizers 2 times a year.
The fertilizers should be applied in 6-8 holes, 2-3 inches deep, equally spaced around the tree (outside half of the area shaded by the canopy).
1st appl.: At the start of the rainy season or when irrigation is possible.

2nd appl.: At the end of rainy season.

Calamansi

Apply the organic fertilizer at the base of the planting holes and place the plant before filling up the hole with soil preferably at the start of the rainy season. One month after transplanting, apply the recommended inorganic fertilizers about 5-10 cm deep around the base of the plant.

Second application of the remaining fertilizers will be applied before the end of the rainy season.

For bearing plants, apply the fertilizers every after harvesting of the fruits.

Coconut

Transplant

Apply the fertilizer in the planting hole including organic fertilizer and fill in the hole with soil preferably at the start of the rainy season.

1-5yrs

Start application of fertilizers 6 months after transplanting and one year thereafter for 5 years.

B**earing**

Application of recommended fertilizer are made annually. The fertilizer are applied in at least ten (10) holes bored 8-16 inches deep at a distance of 1 meter radius from the base of the tree.

Coffee

Seedlings: Incorporate fertilizers with the soil to a depth of 6-10 inches before planting

Matured trees: Mix recommended fertilizers and apply in 2 equal doses: 1st at the start of the rainy season and the 2nd before the end of the rainy season. Application is done in 5-6 holes bored 5 cm deep around the base of the plant

Corn (Wet season planting)

1st appl.: Apply ½ of the N and K recommended fertilizer and all P in band 5 cm to the side and 5 cm below the seed

2nd appl.: Sidedress the remaining fertilizer when plants are about 4 weeks old or at knee-high followed by hilling up and irrigation.

Corn (Dry season planting)

Apply all the fertilizers at planting or 3 weeks after planting.

Cotton

1st appl.: All potash and phosphate fertilizers and two thirds (2/3) of the nitrogenous fertilizer were applied 2 inches to the side of the seed at a depth of 2-4 inches at sowing time.
2nd appl.: Sidedress with the remaining nitrogenous fertilizer before flowering.
Note: With irrigated cotton, the potash dressing should also be divided.

Cucumber

Broadcast or apply the fertilizer including 2/3 on nitrogen in a band 3-4 inches to the side of the row 2 inches deeper than the seed before sowing. Topdress with remaining nitrogenous fertilizer when the 1st fruits are half formed.

Dragon Fruit

Apply fertilizer around the base of the plant 3 months after planting. Repeat application of fertilizer every 3 to 4 months during the 1st year of growth.
Organic fertilizer: Incorporate organic fertilizer 3-4 inches away from the base of the plant.

For established crop: Apply fertilizer uniformly around the base of the plant at the start of rainy season and second application at the end of the rainy season.

Eggplant

Apply all P and K fertilizers including ½ of nitrogen into the furrows at transplanting time. Sidedress with ¼ nitrogenous fertilizer about a month after transplanting. A second sidredressing is made immediately after the 2nd picking of marketable fruits using the remaining ¼ Nitrogen.

Eucalyptus

One time application of fertilizer per year
Apply the same amount of fertilizer per year for production maintenance

Forest trees/jathropa/Bamboo

Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below, at transplanting.

For established crop, apply fertilizer uniformly around the base in boring 3-4 inches deep at outside half of area shaded by canopy; phosphate fertilizer may be applied once every 3 years.

Gabi

1st appl.: Apply the recommended fertilizers including half nitrogen, 3-5 inches to the side at a depth of 3 inches at transplanting time
2nd appl.: Sidedress with the remaining nitrogen a month after transplanting

Garlic

Broadcast the fertilizer immediately prior to the leveling operation of the field to effect thorough incorporation into the soil.

Ginger
1st appl.: Apply the fertilizer including ½ of the nitrogen in the furrow 2-3 inches to the side of seed at planting time.
2nd appl.: Apply the remaining nitrogen in the furrow at hilling up.

G-Melina
Apply fertilizer 30 days after transplanting, 100 kg of complete (14-14-14) fertilizer per hectare will be applied for continued growth and higher percentage of survival. Cover the fertilizer with soil placed in a trench. The succeeding fertilization should be done 3 months after transplanting.

Grape vine (young plant)

Work into the soil or apply in boring to a depth of 8-10 inches the fertilizer.

Grape vine (bearing plant)

1st appl.: Apply as above the fertilizer including half of the N just after harvest
2nd appl.: Apply the remaining N in small doses after the fruits have begun to form.

Grape vine (for new plants)

1st appl.: Mix recommended fertilizers with soil compost or organic manure prior to filling in the hole before planting
2nd appl.: At planting, dissolve the recommended nitrogen fertilizer in one gallon of water and use to water the plant after planting

3rd appl.: Apply in holes bored 3-6 inches from the base 1-2 months after planting
4th appl.: Mix recommended fertilizers and apply in holes bored 5-6 inches from the base 3-4 months after planting
Application will be repeated every two months up to pruning age.

Grass (pasture)/Citronela/Napier/forage crop

1st appl.: Broadcast phosphate and potassium fertilizers together with the organic fertilizers before growth starts.
2nd appl.: Top-dress with half of the nitrogen fertilizer once growth starts
3rd appl.: Top-dress with remaining nitrogen fertilizer 5-8 weeks after the 2nd application.

Grass (Vertiver)
1st appl.: Broadcast half of the recommended fertilizer together with the organic fertilizer before growth.
2nd appl.: Top-dress the remaining fertilizer once growth starts.
3rd appl.: Application of additional N-containing fertilizer about 1 bag/ha maybe applied 5-8 weeks after 2nd application.

Green Manures, legumes/Palawan cherry

1st appl.: Broadcast the fertilizer and plow into the soil thoroughly during seedbed preparation
2nd appl.: Top-dress with more phosphate or potash fertilizer if necessary.

Guava (Guaple)

Newly planted:
Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below, at transplanting together with the organic fertilizer.

Young Trees:
1st appl.: Apply all phosphate and potash with light dressing nitrogen before the onset of the rainy season
2nd appl.: Apply the remaining light dressing of nitrogen when fruits have set.

Bearing:
1st appl.: Apply ½ of the recommended fertilizer at the start of rainy season in 6-8 holes around the tree
2nd appl.: Apply the remaining ½ of the recommended fertilizer before the end of rainy season in 6-8 holes around the tree

Guyabano

Work into the soil or apply the recommended amounts of fertilizers in small doses. Repeat this process during the first few years after planting.

Herbs (Basil/Sage/Rosemary)

Apply the organic fertilizer at least 1 month prior to planting.
Weeks after planting, when plants developed at least two pairs of true leaves, start application of complete fertilizer (14-14-14) and this should be applied once a week to maintain vigorous leaf production.

Ipil-Ipil-
Normally not fertilized.Can tolerate up to 7 months of dry season. However, it does not tolerate water logged soils or extended periods of flood (>3 weeks).
Prefers a soil pH 5.0-8.0

Jackfruit (Langka)

New/transplant
Incorporate fertilizer 3 inches below and 5 inches to the side of the plant at planting

Established (1-3 years)
Apply fertilizer in boring 5-6 inches deep around the base 2 times a year:
Start of rainy season and before the end of rainy season

Bearing trees (4 years and above)
Application of fertilizer will be done annually at the start of the rainy season

Jute

Apply the fertilizer including half the nitrogen in the furrows before sowing. Topdress with remaining nitrogen fertilizer 6-7 weeks after sowing.

Kangkong

1st appl.: Before sowing or transplanting, incorporate 2 kgs/m^2 animal manure (chicken manure)
2nd appl.: 15 days later from sowing, topdress or sidedress 1.0kg (46-0-0)/m^2)

Kudzu

Broadcast the fertilizer as the seedbed is being prepared for a thorough incorporation into the soil.

Kulitis (Spinach)

Apply the fertilizer including ¾ of the nitrogen at transplanting time.
Sidedress with remaining nitrogenous fertilizer two (2) weeks before harvest.

Lanzones/Durian/Longan

To be planted:
Incorporate fertilizer 3 inches below and 5 inches to the side of the plant at planting time.

For established plant: [1-5 years]
Apply fertilizer in boring 5-6 inches deep around the base 2 times a year start of rainy season and before the end of rainy season

Bearing : 6 years and above
Apply 3 times a year as follows:
1st appl.: ¼ of total fertilizer 1-2 months before flowering
2nd appl.: ¼ of total fertilizer at early fruit development (pea stage)
3rd appl.: ½ of total fertilizer after harvesting

Leafy vegetable (Lettuce/Mustard/Pechay)

Sensitive to heavy application of nitrogen. Apply 1/3 of the nitrogen fertilizer with the potash and phosphate at planting. Sidedress the remaining 2/3 of nitrogen into two equal doses at 14 days and at 30 days after planting.

Lemon Grass

1st appl.: Broadcast phosphate and potassium fertilizers before growth starts.
2nd appl.: Top-dress with held nitrogenous fertilizer once growth starts
3rd appl.: Top-dress with remaining nitrogenous fertilizer 5-8 weeks after the 2nd application.

Linseed

Apply all recommended phosphate and potash and 2/3 N fertilizer 2-3 inches to the side and 2-4 inches deep at planting time.
Sidedress with remaining N fertilizer before flowering.

Maguey

At planting, apply recommended fertilizer 3-4 inches deep to the side of the plant.

For established crop, apply fertilizer uniformly around base within radius of 2 feet.

Malunggay (Acid Ipil-5-22-2013)
1st appl. Apply all Phosphate and Potash and ½ Nitrogen 3-4 inches away from the base and 5-6 inches deep at 75 days after sowing.
2nd appl.: Apply the remaining Nitrogen at 150-160 days after sowing.

Note: Do not disturb or shake the plants because it may result to dying of plants

Mango/Mangosteen/Duhat (seedlings)(01-18-16 Revise)
Apply the fertilizer 3 inches below the roots and 5 inches to the side of the plants at planting time

Mango/Mangosteen/Duhat (Young trees)
Mix all organic and inorganic fertilizers and apply in two equal doses in 4-5 holes around the base of the tree 1.5meter radius.

1st appl.: At the start of the rainy season
2nd appl.: Before the end of rainy season.

Mango/Mangosteen/Duhat (bearing trees)
1st appl.: Apply ½ of the recommended fertilizer at the start of rainy season in 6-8 holes around the tree
2nd appl.: Apply the remaining ½ of the recommended fertilizer before the end of rainy season in 6-8 holes around the tree

Mango/Mangosteen/Duhat(Revised August 7, 2015)
Mango **(seedlings):** Apply the fertilizer 5 inches below the roots and 3 inches to side of the plants at planting time together with the Organic Fertilizer.

Mango **(Young trees):** Apply one half of the recommended Inorganic Fertilizer in holes (4-5 inches deep) about 1.5 meters away from the base and 4-5 inches below together with the Organic Fertilizer at the start of the rainy season.

Mango **(Bearing trees):** Apply the remaining fertilizer in holes (4-5 inches deep) about 1.5 meters away from the base and 4-5 inches below before the end of the rainy season.

Millet

Apply half of the recommended fertilizer in the furrow at planting time and the remaining half a month after planting.

Mongo/Mungbean

Broadcast uniformly the recommended fertilizer and plow thoroughly into the soil during seedbed preparation.

Muskmelon/Strawberry

Apply the fertilizer including half of nitrogen in the form of a broad band beneath the seed row at the preparation of the seedbed. Topdress with remaining N fertilizer when the first fruits are half grown. This may be repeated once more or two weeks before harvest.

Narra

General recommendation of (14-14-14) 71 grams/tree is to be applied during the grooming season.
Biofertilizer may also be applied. If liquid fertilizer is to be used, (1/2 tablespoon per gallon of water) is to be sprayed every two to three weeks at seedling stage before planting.

Okra

1st appl.: apply the fertilizer including half of the nitrogen 2-3 inches to the side and 2 inches below the seed at sowing
2nd appl.: Sidedress with remaining nitrogen late in the season at the time "forms" or "blooms" are concentrated in top of the plant.

Okra(Revised August 7, 2015)

1st application: Apply the fertilizer including half of the nitrogen 2-3 inches to the side and 2 inches below the seed at sowing
2nd application: Sidedress with the remaining fertilizer before hilling –up operation.

Onion

Broadcast the fertilizers including half the nitrogen during the last operation in the preparation of the land.
Sidedress with remaining nitrogen at banking time.

Ornamental, flowering/herbs

Apply the recommended fertilizer bi-weekly; water the plants daily.

Ornamental plants (non-flowering)

Apply the recommended fertilizer monthly or as the plants leaves begin to discolor; water the plants daily.

Palm, Date

Mix the fertilizer thoroughly with the soil, or the recommended phosphate is mixed with compost and worked into the soil. No fertilizer containing chloride should be used.

Papaya (Updated)

Field planting; apply around the root zone 30-40cm from the trunk.
8 months from field planting: apply within the root feeding zone.
14 months from field planting: apply within the root feeding zone.

2-4 months
Field planting; apply around the root zone 30-40cm from the trunk.

6-12 months
8 months from field planting: apply within the root feeding zone.

1-2 years
14 months from field planting: apply within the root feeding zone.

Balitbitan

Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below together with the organic fertilizer

Alibangbang

Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below together with the organic fertilizer

Kamuning

Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below together with the organic fertilizer

Papaya (Revised August 7, 2015)

Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below together with the organic fertilizer.

Parnship

Potash and phosphate fertilizer and one-third (1/3) nitrogenous fertilizer are applied before sowing. The remaining N fertilizer is applied into two (2) dressings at one (1) month interval after planting.

Parsley

Apply the recommended amount of fertilizer 2 weeks after planting and should be done monthly thereafter.

Passion Fruit

1st appl.: Apply at flowering
2nd appl.; Apply when fruits are half grown

Patola

Mix the fertilizer including 2/3 of the nitrogen with the soil at planting time
Sidedress with remaining N fertilizer when fruits begin to be formed.

Paper tree (seedlings)
Incorporate fertilizer with the soil to a depth of 6-10 inches
before planting together with the organic fertilizer

Paper tree (established plant)
The fertilizer should be applied in 6-8 holes, 2-3 inches deep, equally
spaced around the tree (outside half of the area shaded by the canopy)
Peanut

Apply the fertilizer in the furrow and cover with 2-3 inches of soil before sowing the seeds. Inoculation of the seeds with nitrogen-fixing bacteria increases the yield and quality of the crops.

Peas

Broadcast or incorporate by tillage the recommended fertilizer including half the nitrogen either at the time of sowing or soon after they come up. Topdress with remaining N fertilizer after flowering but not during the 1st blooming.

Pepper (similar to chili)

Pineapple

Apply the fertilizer including 1/3 the nitrogen 1-2 inches below the soil surface and about 1-2 inches to the side of the plant at planting time.
Topdress with 1/3 n fertilizer three (3) months after planting.
Repeat topdressing with remaining N in the 6th-9th month, but at least before the appearance of the flower buds.

Potatoes

1st appl.; Apply fertilizer including half of the nitrogen in the furrow 3 inches on the side and 3 inches below the seedlings at planting time
2nd appl.: Sidedress with remaining N fertilizer before hilling up 2 months after planting.

Radish

Broadcast the fertilizer including half the N and work into the soil 2-3 inches deep thoroughly at seedling time. Topdress with remaining N fertilizer after seedlings have taken root.

Rambutan/Lychees

Newly planted:
Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below, at transplanting together with the organic fertilizer.

1-5 years
Apply in two equal doses: 1st at start of rainy season and 2nd before the rainy season ends.

6 years and above

Apply 3 times a year as follows:
1st appl.: ¼ of total fertilizer 1-2 months before flowering
2nd appl.: ¼ of total fertilizer at early fruit development (pea stage)
3rd appl.: ½ of total fertilizer after harvesting

Ramie

Potash and phosphate dressings are best given before sowing or disturbed annually at the commencement of the growing period. Nitrogen dressing should be given 2-3 weeks after each cut.

Rice (Updated-7-22-16)
1st appl.: Basal (10-14 Days after Transplanting), apply ½ of the recommended Nitrogen(N) fertilizer with phosphorus and potash 10-14 days after transplanting.
2nd appl.: Active Tillering (Mid tillering stage). Topdress ¼ of N fertilizer within 23-27 days after transplanting.
3rd appl.: Panicle Initiation (33-37 days aftertransplanting). Topdress the remaining ¼ of N and Potash if any to satisfy the required fertilizer requirement at panicle initiation stage.

Rice-HYV-hybrid

1st appl.: Broadcast ½ of the recommended Nitrogen(N) fertilizer with phosphorus and potash during the last harrowing before transplanting.
2nd appl.: Topdress ¼ of N fertilizer within 15 to 25 days after transplanting.
3rd appl.: Topdress the remaining ¼ of N and Potash if any to satisfy the required fertilizer requirement at panicle initiation stage.

Rice (Upland)

1st appl.: Broadcast ½ of nitrogenous fertilizer with phosphorus and potash before or 10 days after sowing
2nd appl.: Topdress with remaining half of nitrogen 50-60 days before harvest.

Roses (similar to ornamentals (flowering)

Rubber

For nursery beds: Mix the fertilizer with compost or sieved soil and apply when the 1st tier of the leaves are formed. Another similar dressing is given one (1) month before budding or grafting.

To be transplanted: [1-2 years]

Apply fertilizer 3-4 inches away from the base of the plant and 4-5 inches below, at transplanting together with the organic fertilizer.

For established crop: apply fertilizer uniformly around the base in 3-4 inches deep at outside half of area shaded by canopy. Application is done at the start of rainy season and onset of dry season at two equal doses.

Santol

Seedlings: Apply the fertilizer 3 inches below the roots and 3-5 inches to the side of the plants at planting.

Young trees: Apply in 2 equal doses 1st at start of rainy season an 2nd before the end of rainy season

Bearing trees: Apply in 2 equal doses in holes around the tree; 1st at start of rainy season and 2nd before the end of rainy season

Seguidillas (see beans, winged)

Sisal

Fertilizer application is made at the commencement of rainy season either broadcast, distributed in band in the neighborhood of the plant or distributed uniformly over the area shaded by the plant leaves.

Sitao (see beans, string)

Squash

Apply the fertilizer including half of the N at planting time, mixing thoroughly with the soil. Sidedress with remaining N when fruits begin to be formed.

Sorghum

1st appl.: Apply ½ of recommended fertilizers at planting
2nd appl.: Apply remaining ½ of the recommended fertilizer 1 month after planting

Soybean

A light dressing of nitrogen is worked into the soil with potash and phosphate fertilizers at sowing such that the fertilizer is 2 inches below and slightly to the side of the seed.

Spinach (see Kulitis)

Star Apple (See Caimito)

Strawberry (similar to coffee)

Sugarcane

Apply the full dose of phosphorus with half portions of the nitrogen and potassium fertilizers either one side or on both sides of the row at planting. Sidedress the balance of the nitrogen and potassium fertilizers about 2-3 months later.

Tamarind

Tamarind Trees bear fruit well even without fertilizer application. However, fertilizer application is recommended to keep the trees in healthy condition. Apply 50 g of 16-20-0/ tree and 100 g of 14-14-14/tree one month after planting. The amount of fertilizer is gradually increased as the trees grow. For early bearing fruits, 5 kg of 14-14-14/tree twice a year. A full bearing tree may need at least 3 kg of 14-14-14/year.

For ratoon cropping - follow the same as that of plant cane crop.
Tapilan

Work into the soil the recommended fertilizer such that it is placed 2 inches below and 3 inches to the side of the seed.

Tobacco

1st appl.: Apply the fertilizer 2 inches below the plant and 3 inches to the side at transplanting
2nd appl. Dissolve Nitrogen fertilizer and apply 7-10 days after transplanting

Tropical bloom (Hibiscus Plant)
Apply the recommended fertilizer monthly or as the plant leaves begin to discolor; water the plants daily.

Tugi/Ubi

Apply the recommended fertilizers including half the nitrogen in band, 3-5 inches to the side at a depth of 3 inches at transplanting time. Sidedress with the remaining nitrogen during hilling up or a month after transplanting

Tomato

1st appl.: Apply the fertilizer including half of the nitrogen 3 inches away from the base of the plant and 3 inches deeper than the root system immediately before or at planting.
2nd appl.: Sidedress with remaining nitrogen fertilizer when the plants start to bloom or when there are some small fruits developing.
Note: Tomatoes are to too heavy application of nitrogen and may result in delayed ripening and fruits that are soft and split easily.

Mint Herb

1st appl.: Apply the fertilizer including ½ of the nitrogen 3 inches away from the base of the plant and 3 inches deeper than the root system immediately before or at planting.
2nd appl.: Sidedress with remaining nitrogen fertilizer one month after planting.

Ubi (similar to Tugi)

Upo

Apply half of the recommended fertilizer uniformly mixed with soil at planting time.
Sidedress with remaining fertilizer when the fruits begin to form.

Watermelon

Apply the fertilizer including half of the recommended nitrogen in trenches near the row or in a strip where the row is to be located during planting time.
Sidedress with ¼ of N when vine begins to run, and again with the rest of the ¼ N when the fruits are half-grown.
Note: Watermelons are sensitive to chlorides

Yam (similar to tugi)

By Ms. Betty Magno-2-15-2014

Mahogany/Fire Tree

Yearly applications of complete fertilizer for the first few years will speed up growth in mahogany trees

African Tulip

Fertilize the seedlings after they have emerged and exhibit two true leaves. Topdress the soil with organic fertilizers once or twice per year.Mulch around the base of the tulip tree with organic materials that will break down and release macro and micronutrients into the soil year-round. Water your African tulip tree when you press your finger into the soil and it is almost dry. Don't let the plant dry out entirely-the tulip tree favors moist growing conditions.

Ba Grass/Bermuda Grass/Carabao (Notification)

Application of organic fertilizer is best recommended in growing Ba Grass. This will also help in conditioning the soil to increase its pH level. Ba Grass thrives well in soil pH 6-7. In using nitrogen fertilizers, avoid the use of ammonium sulfate or other sulfate containing fertilizers to avoid making the soil in acidic condition.

Sage/Rosemary/Basil

As with other herbs, sage should be grown in your soil as this will aid in the production of essential oils,and give the herbs a stronger flavor. If fertilized, the plants will grow prolifically, but will have little taste

Mungbean/Monggo (Revise 04/24/19)

1st application: Apply fertilizer including1/2 of the nitrogen in the furrow 3 inches on the side and 3 inches below the seedlings at planting time.
2nd application: Side dress with the remaining nitrogen fertilizer before hilling up.

Ficus Tree/Maki Tree

1st application: Apply the fertilizer including ½ nitrogen with the soil at planting time.
2nd application: Side dress with remaining nitrogen fertilizer 2-3 months after planting.