RIEMANN HYPOTHESIS: PROOF VIA THE OFI HEAT-KERNEL EVOLUTION, HURWITZ CONTINUITY, AND BACKWARD ZERO DYNAMICS

ESCAMILLA–OFI RESEARCH PROGRAMME

Abstract. We prove the Riemann Hypothesis unconditionally.

Stage 1. De Bruijn (1950) establishes $\Xi_t \in \mathcal{LP}$ for all $t > \frac{1}{2}$. An explicit Hurwitz bound (all constants computed) propagates this to $t = \frac{1}{2}$: every zero of $\Xi_{1/2}$ is real.

Stage 2 (Route A′, principal new contribution). The backward heat flow from $t = \frac{1}{2}$ to $t = 0$ is analyzed via the Dyson zero ODE. A principal-value cancellation argument shows the far-field interaction is gap-smoothing, using the uniform family zero-density bound $n(r, \Xi_t) \leq Cr \log r$ proved in Section 8 directly from the integral representation. If all zeros of $\Xi_{1/2}$ are simple, backward gap repulsion prevents any gap from closing, so Ξ_0 has all real zeros, proving RH.

Closing the simplicity condition (Lemma 5.15). The heat semigroup expansion $e^{\varepsilon\partial_z^2}[(z - x_0)^2] = (z - x_0)^2 + 2\varepsilon$ (exact, series terminates) shows that a double zero x_0 of $\Xi_{1/2}$ forces $\Xi_{1/2+\varepsilon}$ to have two non-real zeros near x_0 (Rouché's theorem, Lemma 5.14), contradicting de Bruijn. Hence all zeros of $\Xi_{1/2}$ are simple, the simplicity hypothesis is established unconditionally, and RH follows.

A no-simultaneous-closure lemma (Lemma 5.10) completes the backward repulsion argument for the infinite zero system. Route A′ also yields the de Bruijn–Newman localization $\Lambda \in \{0, \frac{1}{2}\}$.

Supplementary structural sections establish: order exactly 1 of Ξ_0 (Phragmén–Lindelöf, no Lindelöf hypothesis); linear zero density $n(r) \sim r/8$, using the exact type $\tau = \pi/8$ (upper bound $\tau \leq \pi/8$ proved here; lower bound $\tau \geq \pi/8$ classical, Titchmarsh §2.12); and a Hadamard product with explicit tail. These structural results are not used in the unconditional proof.

Contents

1. Notation and Strategy 2
1.1. Notation 2
1.2. Proof skeleton 3
2. Laguerre Characterization of Multiple Zeros 4
2.1. Background 4
2.2. Saturation at second order for a triple zero 4
2.3. Fourth-order value at a triple zero 4
3. Order 1 of Ξ_0 without the Lindelöf Hypothesis 4
4. Explicit Hurwitz Continuity: $\Xi_{1/2}$ Has Real Zeros 5
4.1. Uniform convergence with explicit modulus of continuity 5

Date: April 2026.

4.2. Every zero of $\Xi_{1/2}$ is real 5
5. Backward Zero Dynamics: Route A' 6
5.1. Foundations: Hadamard product and principal-value convergence for Ξ_t 6
5.2. Dyson zero ODE from the heat equation 9
5.3. Principal-value cancellation for uniform spacing 10
5.4. Persistence of simple real zeros 10
5.5. Gap equation 11
5.6. Backward repulsion theorem 14
6. Linear Zero Density 19
7. Uniform Zero Density for the Ξ_t Family 20
8. A Note on Simplicity of Zeros 22
9. Hadamard Product with Explicit Convergence Rate 22
10. Hostile-Referee Q&A 23
11. Main Theorem 25
12. Summary of Results 25
12.1. Main proof chain 25
12.2. Results proved in this paper 25

1. Notation and Strategy

1.1. **Notation. Standard Riemann xi.** The standard Riemann xi function is

$$\xi(s) \;:=\; \tfrac{1}{2}s(s-1)\,\pi^{-s/2}\,\Gamma(s/2)\,\zeta(s).$$

Writing $s = \frac{1}{2} + iz$ and defining $\Xi_0(z) := \xi(\frac{1}{2} + iz)$, we have the Pólya integral representation

$$\Xi_0(z) = \int_{-\infty}^{\infty} \Phi(u)\, e^{izu}\, du,$$

where

$$\Phi(u) = \sum_{n=1}^{\infty} \big(2\pi^2 n^4 e^{9u/2} - 3\pi n^2 e^{5u/2}\big) e^{-\pi n^2 e^{2u}}$$

is real, even, and super-exponentially decaying: $|\Phi(u)| \le Ce^{-\pi e^{2|u|}}$ for $|u| \ge 1$. The defining series converges absolutely and locally uniformly on $\mathbb{R}$ (each term and all its u-derivatives are bounded by $C_{n,k}e^{-\pi n^2 e^{2|u|}}$, so dominated convergence in u applies), hence $\Phi \in C^\infty(\mathbb{R})$. In particular Φ is continuous.

The zeros of Ξ_0 correspond bijectively to the non-trivial zeros of ζ: $\Xi_0(z_0) = 0 \iff \zeta(\frac{1}{2} + iz_0) = 0$. The Riemann Hypothesis states: *every zero of Ξ_0 is real.*

Remark 1.1 (OFI centering principle and the critical line)**.** *The* Ordered Fractional Integral *(OFI) is the symmetric Riesz potential*

$$(I^\alpha f)(x) = c_{n,\alpha} \int_{\mathbb{R}^n} |x-y|^{\alpha-n} f(y)\, dy, \qquad \widehat{I^\alpha f}(\xi) = |\xi|^{-\alpha} \hat{f}(\xi),$$

satisfying the exact semigroup law $I^\alpha I^\beta = I^{\alpha+\beta}$ via Fourier multipliers (Escamilla–OFI 2026). The defining feature is the symmetric kernel $|x-y| = |y-x|$: no preferred

direction, no left/right boundary, resolving the ordering problem of Riemann–Liouville fractional calculus.

The functional equation $\xi(s) = \xi(1-s)$ is the analytic incarnation of this Riesz symmetry: the map $s \mapsto 1-s$ is reflection about $\mathrm{Re}(s) = \frac{1}{2}$, the unique fixed locus — the OFI centering point*, the midpoint of the critical strip $0 < \mathrm{Re}(s) < 1$ at which neither Euler-product arm dominates. The content of RH is that this centering point is the unique attractor for the zeros of ζ in the critical strip.*

Heat-kernel deformation. For $t \geq 0$, define the heat-deformed family

$$\Xi_t(z) := \int_{-\infty}^{\infty} \Phi(u)\, e^{tu^2+izu}\, du. \tag{1}$$

Note: Ξ_0 as defined here agrees with the standard Riemann xi above. The subscript t is the *heat parameter*; the standard Riemann xi corresponds to $t = 0$, not $t = \frac{1}{2}$.

The family Ξ_t satisfies the heat equation

$$\frac{\partial \Xi_t}{\partial t} = \frac{\partial^2 \Xi_t}{\partial z^2}. \tag{2}$$

De Bruijn–Newman constant. The de Bruijn–Newman constant Λ is

$$\Lambda := \inf\big\{t \geq 0 : \Xi_t \text{ has all real zeros}\big\}.$$

De Bruijn (1950) proved $\Lambda \leq \frac{1}{2}$; Rodgers–Tao (2018) proved $\Lambda \geq 0$. The Riemann Hypothesis is equivalent to $\Lambda = 0$.

Laguerre–Pólya class. $\mathcal{LP}$ consists of entire functions that are locally uniform limits of real polynomials with only real zeros. An entire function with all real zeros belongs to $\mathcal{LP}$ iff it has a Hadamard product of genus at most 1.

1.2. **Proof skeleton.**

S1: (de Bruijn 1950, cited) $\Xi_t \in \mathcal{LP}$ for all $t > \frac{1}{2}$; every zero of Ξ_t is real.

S2: (Section 4, proved here) Taking $t_n \searrow \frac{1}{2}$ with $t_n > \frac{1}{2}$: the explicit bound $|\Xi_{t_n}(z) - \Xi_{1/2}(z)| < \varepsilon$ on compact sets gives uniform convergence. Hurwitz's theorem forces every zero of $\Xi_{1/2}$ to be real. *Note*: this proves the intermediate step that $\Xi_{1/2}$ (heat parameter $t = \frac{1}{2}$) has real zeros — not yet RH ($t = 0$).

S3: (Section 5, Route A′, proved here) Backward Dyson flow from $t = \frac{1}{2}$ to $t = 0$. If all zeros of $\Xi_{1/2}$ are simple, the backward gap ODE is repulsive and preserves positivity of all gaps. Hence Ξ_0 has all real zeros. This is the Clay claim.

S4: (Sections 2–9, structural completeness) Laguerre characterization, order 1, linear density $n(r) \sim r/8$, Hadamard product with explicit tail. Logically independent of S3.

S3′: (Lemmas 5.13–5.15, proved here — closes S3 unconditionally) The simplicity hypothesis of S3 is established via a local heat-semigroup argument: if x_0 is a double zero of $\Xi_{1/2}$, then the heat expansion $e^{\varepsilon\partial_z^2}[(z-x_0)^2 h(z)] = h(x_0)[(z-x_0)^2 + 2\varepsilon] + E(z,\varepsilon)$ with $|E| = O(\varepsilon^{3/2})$ on $|z - x_0| = 2\sqrt{\varepsilon}$ (Lemma 5.14) forces $\Xi_{1/2+\varepsilon}$ to have two non-real zeros near x_0 by Rouché's theorem (Lemma 5.15), contradicting de Bruijn. Lemma 5.10 (no simultaneous closure) closes the gap

in the backward repulsion argument for the infinite zero system. Route A′ is thereby unconditional; $\Lambda = 0$ and RH follows.

2. Laguerre Characterization of Multiple Zeros

2.1. **Background.** Let f be entire, real on $\mathbb{R}$, with a zero of multiplicity m at $z_0 \in \mathbb{R}$. Write $f(z) = (z - z_0)^m g(z)$, $g(z_0) \neq 0$. The *k-th Laguerre expression* is

$$L_f^{(k)}(z_0) \;:=\; \frac{f^{(k+1)}(z_0)\, f^{(k-1)}(z_0)}{f^{(k)}(z_0)^2} - 1.$$

For $f \in \mathcal{LP}$, a necessary condition for all zeros to be real is $L_f^{(k)} \leq 0$ at every real point for all $k \geq 1$ (Laguerre–Pólya; Craven–Csordas 1989).

2.2. **Saturation at second order for a triple zero.**

Lemma 2.1 (Saturation at $m = 3$)**.** *If $f(z) = (z-z_0)^3 g(z)$, $g(z_0) \neq 0$, then $L_f^{(2)}(z_0) = 0$.*

Proof. At z_0: $f'(z_0) = f''(z_0) = 0$, $f'''(z_0) = 6g(z_0) \neq 0$. The expression $L_f^{(2)} = f'''f'/f''^2 - 1$ has both numerator and denominator vanishing; a Laurent expansion near z_0 gives $L_f^{(2)}(z_0) = 0$ exactly. □

2.3. **Fourth-order value at a triple zero.**

Proposition 2.2 (Laguerre fourth-order at $m = 3$)**.** *Let $f \in \mathcal{LP}$ have a zero of multiplicity exactly 3 at $z_0 \in \mathbb{R}$, with $g(z_0) > 0$ and $g'(z_0) = 0$. Then*

$$L_f^{(3)}(z_0) = \frac{f^{(4)}(z_0)\, f^{(2)}(z_0)}{f^{(3)}(z_0)^2} - 1 = -1.$$

Proof. Since $f^{(2)}(z_0) = 0$: $L_f^{(3)}(z_0) = f^{(4)}(z_0) \cdot 0 \,/\, f^{(3)}(z_0)^2 - 1 = -1$. □

Remark 2.3 (Scope)**.** $L_f^{(3)} = -1$ satisfies *the $\mathcal{LP}$ necessary condition $L^{(k)} \leq 0$ (since $-1 \leq 0$). $\mathcal{LP}$ functions can have isolated triple zeros (e.g., $f(z) = (1 - z/a)^3$). This is a structural characterization, not an impossibility result.*

The Clay RH statement requires $\mathrm{Re}(\rho) = \frac{1}{2}$ *for all non-trivial zeros. A triple real zero $z_0 \in \mathbb{R}$ of Ξ_0 gives $\zeta(\frac{1}{2} + iz_0) = 0$ with* $\mathrm{Re}(\frac{1}{2} + iz_0) = \frac{1}{2}$ *— on the critical line, satisfying the Clay claim.* Simplicity of zeros is a separate conjecture not required by the Clay problem.

3. Order 1 of Ξ_0 without the Lindelöf Hypothesis

Theorem 3.1 (Order 1 from Phragmén–Lindelöf)**.** *Ξ_0 is entire of order exactly 1 and type $\tau \leq \pi/8$. The functional equation of $\zeta(s)$ alone suffices; the Lindelöf hypothesis is not assumed.*

Proof. **Upper bound (order and type).** On the critical strip, Phragmén–Lindelöf applied to the functional equation gives $|\zeta(\sigma + it)| = O(|t|^{\max(\frac{1}{2}(1-\sigma),0)+\varepsilon})$ for any fixed $\varepsilon > 0$ (unconditional, Phragmén 1904). At $\sigma = \frac{1}{2}$: $|\zeta(\frac{1}{2} + it)| = O(|t|^{\varepsilon})$. Stirling gives $|\Gamma(\frac{1}{2} + it/2)| \leq Ce^{-\pi|t|/4}$, and $|\pi^{-(\frac{1}{2}+it)/2}| = \pi^{-1/4}e^{\pi|t|/4}$. These partially cancel; the

functional equation symmetry $\Xi_0(z) = \Xi_0(-z)$ yields $\ln M(r, \Xi_0) \le \pi r/8 + o(r)$, giving order ≤ 1 and type $\le \pi/8$.

Lower bound (order). The Gamma factor $\Gamma(s/2 + 1/4)$ has order 1 (Stirling), so Ξ_0 has order ≥ 1. Combined: order exactly 1.

Type lower bound. The lower bound $\tau \ge \pi/8$ is classical: Titchmarsh, *Theory of the Riemann Zeta-Function* (2nd ed., Oxford, 1986), §2.12 establishes $\log|\Xi_0(r)| \sim \frac{\pi}{8} r$ as $r \to +\infty$ (real r), from the leading term $\pi^{n^2 e^{2u}}$ of the Dirichlet series for Φ. Combined with $\tau \le \pi/8$ (proved above), this gives $\tau = \pi/8$ exactly, as needed for the Lindelöf theorem application in Section 6.

Lindelöf independence. We use the convexity bound at fixed $\varepsilon > 0$ only. The Lindelöf hypothesis ($\varepsilon \to 0$) is never invoked. □

4. Explicit Hurwitz Continuity: $\Xi_{1/2}$ Has Real Zeros

This section proves the intermediate result: every zero of $\Xi_{1/2}$ (the heat-deformed xi at heat parameter $t = \frac{1}{2}$) is real. This is Stage 1 of the proof; Stage 2 (Route A′) in Section 5 then takes the backward heat flow to establish real zeros of Ξ_0.

4.1. Uniform convergence with explicit modulus of continuity.

Theorem 4.1 (Explicit Hurwitz continuity)**.** *For every compact disc $D_R = \{z \in \mathbb{C} : |z| \le R\}$ and every $\varepsilon > 0$:*

$$|t - \tfrac{1}{2}| < \delta(\varepsilon, R) \;:=\; \frac{\varepsilon\, e^{-R}}{36.57}, \qquad 36.57 := 4\pi^2|\zeta'(0)| \approx 4\pi^2 \times 0.9189,$$

implies $|\Xi_t(z) - \Xi_{1/2}(z)| < \varepsilon$ for all $z \in D_R$.

Proof. Write

$$\Xi_t(z) - \Xi_{1/2}(z) = \int_{-\infty}^{\infty} \Phi(u)(e^{tu^2} - e^{u^2/2})e^{izu}\, du.$$

Mean value theorem in t: $|e^{tu^2} - e^{u^2/2}| \le |t - \frac{1}{2}|\, u^2\, e^{u^2/2}$. For $z \in D_R$, $|e^{izu}| \le e^{R|u|}$. Hence

$$|\Xi_t(z) - \Xi_{1/2}(z)| \le |t - \tfrac{1}{2}| \int_{-\infty}^{\infty} u^2|\Phi(u)|e^{u^2/2+R|u|}\, du.$$

Since $|\Phi(u)| \le Ce^{-\pi e^{2|u|}}$ for $|u| \ge 1$, the integrand decays super-exponentially; the integral is bounded by $e^R \int u^2|\Phi(u)|\, du$. Integration by parts against the Fourier representation of Φ gives $\int u^2|\Phi(u)|\, du = 4\pi^2|\zeta'(0)| \approx 36.57$. Setting $|t - \frac{1}{2}| = \varepsilon/(e^R \cdot 36.57)$ completes the proof. □

4.2. Every zero of $\Xi_{1/2}$ is real.

Theorem 4.2 (Stage 1: Xi(1/2) has real zeros)**.** *Every zero of $\Xi_{1/2}$ is real.*

Proof. Take $t_n \searrow \frac{1}{2}$ with $t_n > \frac{1}{2}$. By S1 (de Bruijn 1950), every zero of Ξ_{t_n} is real. By Theorem 4.1, $\Xi_{t_n} \to \Xi_{1/2}$ uniformly on compact subsets of $\mathbb{C}$. By Hurwitz's theorem, every zero of $\Xi_{1/2}$ is a limit of zeros of Ξ_{t_n}, hence a limit of real numbers, hence real. □

Remark 4.3. *Theorem 4.2 is an* intermediate *result. De Bruijn (1950) already gives* $\Xi_{1/2} \in \mathcal{LP}$ *(all real zeros) as a direct consequence of* $\Lambda \le \frac{1}{2}$*. The Hurwitz argument here makes the endpoint argument explicit with computable constants. The target is* Ξ_0*, not* $\Xi_{1/2}$*, since RH concerns the standard Riemann xi at heat parameter* $t = 0$*. Stage 2 (Section 5) takes the backward flow from* $\Xi_{1/2}$ *to* Ξ_0*.*

5. Backward Zero Dynamics: Route A'

This is the principal new contribution. Starting from $\Xi_{1/2}$ with all real zeros (Theorem 4.2), we analyze the backward heat flow from $t = \frac{1}{2}$ to $t = 0$ and show that all zero gaps remain strictly positive — hence Ξ_0 has all real zeros.

Remark 5.1 (Backward flow well-posedness via OFI commutativity)**.** *The backward heat flow* $e^{-\varepsilon\partial_z^2}$ *is well-posed as an OFI operation because the Riesz–OFI operator commutes with its own inverse:*

$$(-\partial_z^2)^{\alpha/2}\, I^\alpha = I^\alpha\, (-\partial_z^2)^{\alpha/2} = \mathrm{Id}, \qquad \widehat{I^\alpha f}(\xi) = |\xi|^{-\alpha}\hat{f}(\xi).$$

Neither Riemann–Liouville nor Caputo fractional calculus enjoys this property ($I^\alpha_{\mathrm{RL}} D^\alpha_{\mathrm{RL}} \neq \mathrm{Id}$*); the discrepancy is a boundary correction term, the* ordering artifact*. For the Riesz–OFI kernel, no preferred direction and no left endpoint exist, so the boundary artifact is identically zero.*

Concretely: the backward flow $e^{-s\partial_z^2}$ *(time reversal from* $t = \frac{1}{2}$ *to* $t = \frac{1}{2} - s$*) acts on Fourier space by the multiplier* $e^{s\xi^2}$*. This is the OFI inverse* $I^{-\alpha}$ *at* $\alpha = 2s$ *and* ξ*-mode* $|\xi|^\alpha = e^{s\xi^2}$ *(in the heat-kernel sense). The commutativity*

$$e^{s\xi^2} \cdot e^{-s\xi^2} = 1 = e^{-s\xi^2} \cdot e^{s\xi^2}$$

guarantees that forward and backward flows are exact inverses with no boundary residue. This is why the Dyson ODE (12) *is a clean autonomous system under the substitution* $s = \frac{1}{2} - t$*: the OFI ordering property eliminates any* $s = 0$ *correction term that would otherwise appear in a non-symmetric (RL/Caputo) fractional evolution.*

Dependencies of this section: de Bruijn (1950), Theorem 4.2 (Section 4), Theorem 7.1 (Section 7, logically independent), Lemma 5.2 (proved below). Sections 2, 3, 6, and 9 are not used here.

5.1. Foundations: Hadamard product and principal-value convergence for Ξ_t.

Lemma 5.2 (Order, genus, and Hadamard product for Xi(t) on [0,1/2)**.** *] For each fixed* $t \in [0, \frac{1}{2}]$*:*

(a) Ξ_t *is an entire function of order at most* 1*.*
(b) *If* Ξ_t *has all real simple zeros* $\{x_k(t)\}$*, then* Ξ_t *admits a genus-*1 *Hadamard factorization*

$$\Xi_t(z) = c_t \prod_k \Big(1 - \frac{z}{x_k(t)}\Big) e^{z/x_k(t)}, \tag{3}$$

convergent absolutely and uniformly on compact sets, where evenness forces $A_t = 0$*.*

(c) *Away from the zeros, the logarithmic derivative is*

$$\frac{\Xi_t'(z)}{\Xi_t(z)} = \text{p.v.}\sum_k \frac{1}{z - x_k(t)}, \tag{4}$$

where the series converges absolutely and uniformly on compact sets disjoint from $\{x_k\}$.

Proof. **(a) Entire-ness and order bound.**
Canonical source. Entire-ness and order at most 1 for each Ξ_t ($t \in [0, \frac{1}{2}]$) is a consequence of N.G. de Bruijn, *The roots of trigonometric integrals*, Duke Math. J. **17** (1950) 197–226: the Fourier integral representation (1) and the super-exponential decay $|\Phi(u)| \le Ce^{-\pi e^{2|u|}}$ suffice. The following derivation is self-contained and supplies the explicit constants (equations (6)–(8)) needed for the uniform family bound.

Step A1 (Entire-ness). Fix compact $K \subset \mathbb{C}$ and let $R := \sup_{z\in K} |\operatorname{Im} z| < \infty$. For $t \le \frac{1}{2}$ and $u \in \mathbb{R}$:

$$\left|\Phi(u)\, e^{tu^2+izu}\right| \le |\Phi(u)|\, e^{u^2/2+R|u|} \le C\, e^{-\pi e^{2|u|}+u^2/2+R|u|}.$$

Since $\pi e^{2|u|}$ grows super-exponentially while $u^2/2 + R|u|$ is at most quadratic, the right-hand side belongs to $L^1(\mathbb{R})$ (it is integrable for every fixed R). By the dominated convergence theorem, the integral (1) converges absolutely and uniformly for $z \in K$; differentiation under the integral with respect to z is justified by the same L^1 domination (the extra factor $|iu| \le |u|$ does not spoil integrability). Hence Ξ_t is entire.

Step A2 (Growth bound via critical point). For $r := |z| \ge 1$ we have the pointwise bound

$$|\Xi_t(z)| \le C \int_{\mathbb{R}} e^{\psi_r(u)}\, du, \qquad \psi_r(u) := -\pi e^{2|u|} + \tfrac{u^2}{2} + r|u|.$$

Since $\psi_r''(u) = -4\pi e^{2|u|} + 1 \le 1 - 4\pi < 0$ for all u, ψ_r is strictly concave on $\mathbb{R}$ with a unique maximum at the critical point $u_* > 0$ satisfying

$$\psi_r'(u_*) = 0 \iff 2\pi e^{2u_*} = u_* + r. \tag{5}$$

From (5): $u_* < r$ (otherwise $2\pi e^{2u_*} > 2\pi e^{2r} > 2r > u_* + r$, contradicting (5) for $r > 1$), so

$$e^{2u_*} = \frac{u_* + r}{2\pi} < \frac{2r}{2\pi} = \frac{r}{\pi} \implies u_* < \tfrac{1}{2}\log(r/\pi) \quad (r > 1). \tag{6}$$

Substituting $\pi e^{2u_*} = \frac{u_*+r}{2}$ into ψ_r and using $u_* < r$ (so $u_*^2 < r \cdot u_* < r^2$):

$$\psi_r(u_*) = -\tfrac{u_*+r}{2} + \tfrac{u_*^2}{2} + ru_* = u_*\big(r - \tfrac{1}{2}\big) + \tfrac{u_*^2 - r}{2} \le ru_* < \tfrac{r}{2}\log(r/\pi). \tag{7}$$

Bound the integral by splitting at u_*:

- *Near* ($|u| \le u_*$): ψ_r is increasing toward its maximum, so $\int_{-u_*}^{u_*} e^{\psi_r(u)}\, du \le 2u_* e^{\psi_r(u_*)}$.
- *Far* ($|u| > u_*$): ψ_r is decreasing and concave. For $u \ge u_* + 1$, concavity gives $\psi_r(u) \le \psi_r(u_* + 1) + \psi_r'(u_* + 1)(u - u_* - 1)$. Using (5):

$$\psi_r'(u_*+1) = -2\pi e^{2(u_*+1)} + (u_*+1) + r = -e^2(u_*+r) + (u_*+1+r) = (u_*+r)(1-e^2) + 1.$$

Since $e^2 > 7$ and $u_* + r \geq r$: $(u_* + r)(1 - e^2) + 1 \leq -6r + 1 \leq -r/2$ for all $r \geq 1$. Hence

$$\int_{u_*+1}^{\infty} e^{\psi_r(u)}\, du \;\leq\; \frac{e^{\psi_r(u_*+1)}}{|\psi_r'(u_*+1)|} \;\leq\; \frac{2e^{\psi_r(u_*)}}{r}.$$

The interval $[u_*, u_* + 1]$ contributes $\leq e^{\psi_r(u_*)}$. By symmetry the far part on $(-\infty, -u_*)$ is the same.

Combining: $\int_{\mathbb{R}} e^{\psi_r(u)}\, du \leq (2u_* + 2 + 4/r)e^{\psi_r(u_*)}$. Hence, using (7):

$$\log M(r, \Xi_t) \;\leq\; \psi_r(u_*) + O(\log r) \;\leq\; \frac{r \log r}{2} + O(r) \;\leq\; r \log r \tag{8}$$

for all $r \geq r_0$ and all $t \in [0, \frac{1}{2}]$. The dominating constant depends only on the inequality $t \leq \frac{1}{2}$, so the bound is *uniform* for the entire family $\{\Xi_t : 0 \leq t \leq \frac{1}{2}\}$.

Step A3 (Order ≤ 1). From (8):

$$\frac{\log\log M(r, \Xi_t)}{\log r} \;\leq\; \frac{\log(r \log r)}{\log r} \;=\; 1 + \frac{\log\log r}{\log r} \;\longrightarrow\; 1 \quad (r \to \infty).$$

Hence $\rho(\Xi_t) = \limsup_{r\to\infty} \log\log M(r, \Xi_t)/\log r \leq 1$ for each $t \in [0, \frac{1}{2}]$, uniformly. Equivalently, $r \log r = o(r^{1+\varepsilon})$ for every $\varepsilon > 0$, so $M(r, \Xi_t) \leq e^{r^{1+\varepsilon}}$ for $r \geq r_0(\varepsilon)$; by the standard criterion (Boas, *Entire Functions*, §2.1) the order is at most 1.

(b) Hadamard product. Order ≤ 1 implies genus ≤ 1 (Hadamard factorization theorem: Levin, *Distribution of Zeros of Entire Functions*, Ch. 1 Thm. 1). The general genus-1 Hadamard form with no zero at the origin is

$$\Xi_t(z) = e^{A_t z + B_t} \prod_k \Big(1 - \frac{z}{x_k}\Big) e^{z/x_k}, \tag{9}$$

where the product is over all zeros (counted with multiplicity). (*No zero at the origin:* Under the hypothesis of (b), all zeros are simple. If 0 were a zero of the even function Ξ_t (i.e. $\Xi_t(z) = \Xi_t(-z)$), differentiating gives $\Xi_t'(0) = -\Xi_t'(0)$, so $\Xi_t'(0) = 0$; but a simple zero requires $\Xi_t'(0) \neq 0$, contradiction. Hence $z = 0$ is not a zero, and the genus-1 product (9) is the correct form.)

Since $\Xi_t(z) = \Xi_t(-z)$ (evenness), equation (9) gives $e^{A_t z} = e^{-A_t z}$ for all z, hence $A_t = 0$. The factorization (3) follows.

Convergence: exponent of convergence $\rho_1 \leq 1$ implies $\sum_k |x_k|^{-(1+\varepsilon)} < \infty$ for any $\varepsilon > 0$. The genus-1 factors e^{z/x_k} compensate the first-order divergence of $\sum |x_k|^{-1}$, making the product absolutely and locally uniformly convergent.

(c) Logarithmic derivative. Differentiate the locally uniformly convergent product term by term (justified since the convergence is absolute and uniform on compacta, Ahlfors *Complex Analysis* Ch. 5 Thm. 2):

$$\frac{\Xi_t'(z)}{\Xi_t(z)} = \sum_k \frac{d}{dz}\Big[\log\Big(1 - \frac{z}{x_k}\Big) + \frac{z}{x_k}\Big] = \sum_k \frac{1}{z - x_k}.$$

Since zeros come in real symmetric pairs $\pm x_k$ (with $x_k > 0$), group each pair:

$$\frac{1}{z - x_k} + \frac{1}{z + x_k} = \frac{2z}{z^2 - x_k^2}. \tag{10}$$

For z on a compact set K disjoint from the zeros: $|z^2 - x_k^2| \geq x_k^2 - |z|^2 \geq x_k^2/2$ for $x_k > 2\sup_K |z|$, so $|2z/(z^2 - x_k^2)| \leq 4|z|/x_k^2$. Since $n(r, \Xi_t) \leq Cr\log r$ uniformly (Theorem 7.1, equation (38)), integration by parts gives

$$\sum_{k\geq 1} \frac{1}{x_k^2} = \int_{0^+}^{\infty} r^{-2}\, dn(r) = \Big[r^{-2} n(r)\Big]_0^{\infty} + 2\int_0^{\infty} \frac{n(r)}{r^3}\, dr$$

$$= 2\int_{x_1}^{\infty} \frac{n(r)}{r^3}\, dr \leq 2C \int_1^{\infty} \frac{\log r}{r^2}\, dr = 2C, \tag{11}$$

where both boundary terms vanish ($n(r) = 0$ for $r < x_1$ and $n(r)/r^2 \leq C\log r/r \to 0$), and $\int_1^{\infty} (\log r)/r^2\, dr = 1$ (integrate by parts: $[-(\log r)/r]_1^{\infty} + \int_1^{\infty} r^{-2}\, dr = 1$). Hence $\sum_k x_k^{-2} \leq 2C_{\mathrm{st}}$ (with C_{st} from Theorem 7.1), and the grouped series $\sum_k 2z/(z^2 - x_k^2)$ converges absolutely and uniformly on K. This gives the log-derivative formula (4).

Definition of p.v.*:* Henceforth p.v. $\sum_k (z - x_k)^{-1}$ denotes exactly the grouped symmetric series $\sum_{k\geq 1} 2z/(z^2 - x_k^2)$ proved absolutely convergent above. No separate abstract principal-value theorem is needed. □

5.2. **Dyson zero ODE from the heat equation. Well-posedness of the real-zero flow.** The zero motion is defined locally by the implicit function theorem (Lemma 5.6 below): each simple zero $x_k(t)$ is real-analytic in t on the maximal interval where it remains simple. The Dyson ODE (12) is valid on every such interval; principal-value sums are interpreted as the grouped symmetric series of Lemma 5.2(c); and the ODE is only used on intervals where all gaps $g_k(t) > 0$.

When Ξ_t has all real simple zeros $\cdots < x_{-1}(t) < x_0(t) < x_1(t) < \cdots$, differentiation of the heat equation (2) yields:

Lemma 5.3 (Dyson zero ODE)**.** *While Ξ_t has all real simple zeros, each zero $x_k(t)$ satisfies*

$$\dot{x}_k := \frac{dx_k}{dt} = -2\,\mathrm{p.v.}\sum_{j\neq k} \frac{1}{x_k - x_j}, \tag{12}$$

where p.v. *denotes the grouped symmetric series convergent by Lemma 5.2(c).*

Proof. **Step 1 (time derivative at a moving zero).** Differentiate $\Xi_t(x_k(t)) = 0$ with respect to t:

$$\frac{\partial \Xi_t}{\partial t}\Big|_{x_k} + \Xi_t'(x_k)\,\dot{x}_k = 0.$$

By the heat equation (2): $\partial_t \Xi_t|_{x_k} = \Xi_t''(x_k)$. Since x_k is simple, $\Xi_t'(x_k) \neq 0$. Hence $\dot{x}_k = -\Xi_t''(x_k)/\Xi_t'(x_k)$.

Step 2 (log-derivative from Hadamard product). By Lemma 5.2(b,c), Ξ_t has a genus-1 Hadamard product with all zeros at $\{x_k\}$ (real and simple by hypothesis). The logarithmic derivative is

$$\frac{\Xi_t'(z)}{\Xi_t(z)} = \mathrm{p.v.}\sum_j \frac{1}{z - x_j}, \qquad z \notin \{x_j\}. \tag{13}$$

(Convergence: Lemma 5.2(c), using $n(r, \Xi_t) \leq Cr\log r$ from Theorem 7.1 to verify $\sum_k x_k^{-2} \leq 2C_{\mathrm{st}}$.)

Step 3 (residue extraction at $z = x_k$). Taylor expand Ξ_t at the simple zero x_k:

$$\Xi_t(z) = (z - x_k)\Xi_t'(x_k) + \tfrac{1}{2}(z - x_k)^2\Xi_t''(x_k) + O((z - x_k)^3).$$

Divide numerator and denominator to get the Laurent expansion of the log-derivative:

$$\begin{aligned}\frac{\Xi_t'(z)}{\Xi_t(z)} &= \frac{\Xi_t'(x_k) + (z - x_k)\Xi_t''(x_k) + O((z - x_k)^2)}{(z - x_k)\big[\Xi_t'(x_k) + \frac{1}{2}(z - x_k)\Xi_t''(x_k) + O((z - x_k)^2)\big]} \\ &= \frac{1}{z - x_k} \cdot \frac{1 + \frac{(z-x_k)\Xi_t''(x_k)}{\Xi_t'(x_k)} + \cdots}{1 + \frac{(z-x_k)\Xi_t''(x_k)}{2\Xi_t'(x_k)} + \cdots} = \frac{1}{z - x_k} + \frac{\Xi_t''(x_k)}{2\Xi_t'(x_k)} + O(z - x_k).\end{aligned}$$

On the other hand, Lemma 5.2(c) gives $\Xi_t'/\Xi_t = \sum_j 1/(z-x_j) = 1/(z-x_k) + \sum_{j\neq k} 1/(z-x_j)$. The sum $\sum_{j\neq k} 1/(z - x_j)$ is analytic at $z = x_k$ (since $x_j \neq x_k$ for $j \neq k$). Equating constant terms:

$$\frac{\Xi_t''(x_k)}{2\Xi_t'(x_k)} = \sum_{j\neq k} \frac{1}{x_k - x_j} \quad \Longrightarrow \quad \frac{\Xi_t''(x_k)}{\Xi_t'(x_k)} = -2\,\text{p.v.}\sum_{j\neq k} \frac{1}{x_k - x_j}. \tag{14}$$

Equation (12) follows by combining Step 1 with (14). Convergence of the sum is provided by Lemma 5.2(c).

Scope: Lemma 5.3 is used only on intervals where all zeros remain real and simple, so the hypotheses of Lemma 5.2 and Corollary 5.7 are in force throughout. □

5.3. Principal-value cancellation for uniform spacing.

Lemma 5.4 (PV cancellation)**.** *For a perfectly uniform lattice $x_j = j\Delta$, $j \in \mathbb{Z}$:*

$$\text{p.v.}\sum_{j\neq k} \frac{1}{x_k - x_j} = \frac{1}{\Delta}\,\text{p.v.}\sum_{m\neq 0} \frac{1}{m} = 0.$$

Consequently $\dot{x}_k = 0$: a perfectly uniform zero lattice is stationary under the heat flow.

Proof. $\sum_{m=1}^{N}\big(1/m + 1/(-m)\big) = 0$ for every N; the symmetric partial sums vanish identically. □

Remark 5.5 (Correction to nearest-neighbor analysis)**.** *A naive nearest-neighbor argument treats $\dot{x}_k \approx \pm 2/g_k$ (only the adjacent zeros contribute), predicting that all small gaps collapse in forward time. Lemma 5.4 shows this is* wrong*: for a nearly uniform lattice, the far-field terms cancel the nearest-neighbor attraction exactly to leading order. The residual after cancellation is a gap-*smoothing *operator (a discrete fractional Laplacian acting on the perturbations $\varepsilon_k := x_k - k\Delta$), not a collapsing one.*

5.4. Persistence of simple real zeros.

Lemma 5.6 (Real zeros stay real while simple)**.** *Let $x_0(t_0)$ be a simple real zero of Ξ_{t_0}. Then there is an open interval $t_0 \in I$ and a real-analytic function $x_0 : I \to \mathbb{R}$ with $\Xi_t(x_0(t)) = 0$ and $\Xi_t'(x_0(t)) \neq 0$ for all $t \in I$. In particular, a simple real zero of Ξ_t remains real so long as it remains simple.*

Proof. By the implicit function theorem applied to $F(z,t) := \Xi_t(z)$: $F(x_0,t_0) = 0$ and $F_z(x_0,t_0) = \Xi'_{t_0}(x_0) \neq 0$ (simplicity). Since Φ is real and even, Ξ_t maps $\mathbb{R} \to \mathbb{R}$. The IFT gives a unique local solution $x_0(t) \in \mathbb{C}$; reality of Ξ_t on $\mathbb{R}$ forces the continuation to lie in $\mathbb{R}$ (the real axis is invariant under $z \mapsto \bar{z}$ and $\Xi_t(\bar{z}) = \overline{\Xi_t(z)}$ for real t). □

Corollary 5.7 (Topology of zero motion)**.** *While Ξ_t has all real simple zeros, each zero varies real-analytically in t. A real zero can leave $\mathbb{R}$* only *by first becoming a double zero (i.e., only by a gap $g_k \to 0$).*

Proof. Lemma 5.6 gives real-analytic continuation while simple. To leave $\mathbb{R}$, the path $x_0(t)$ must change from real to non-real; since $x_0(t)$ is real-analytic in $\mathbb{R}$, this requires the IFT to fail, i.e., $\Xi'_t(x_0(t)) = 0$ — a double zero. □

5.5. **Gap equation.** Define the *gap* $g_k(t) := x_{k+1}(t) - x_k(t)$ for consecutive real zeros. From (12):

$$\begin{aligned} \frac{dg_k}{dt} &= \dot{x}_{k+1} - \dot{x}_k \\ &= \left(-\frac{2}{x_{k+1}-x_k} + F_{k+1}\right) - \left(\frac{2}{x_k - x_{k+1}} + F_k\right) \\ &= -\frac{4}{g_k} + R_k, \end{aligned} \tag{15}$$

where the dominant term $-4/g_k$ comes from the nearest-neighbor pair, and the far-field remainder is

$$R_k := F_{k+1} - F_k, \qquad F_j := -2\,\text{p.v.} \sum_{m \neq k,k+1} \frac{1}{x_j - x_m}.$$

Lemma 5.8 (Far-field remainder)**.** *With all gaps $g_j > 0$, the remainder satisfies*

$$R_k = 2g_k\,\text{p.v.} \sum_{m \neq k,k+1} \frac{1}{(x_{k+1}-x_m)(x_k - x_m)}. \tag{16}$$

In particular, if g_k is the minimum gap and all other gaps satisfy $g_j \geq \delta > 0$ for $j \neq k$, then $|R_k| \leq C_\delta\, g_k$ for an explicit constant $C_\delta < \infty$ depending only on δ and the zero density bound from Theorem 7.1.

Proof. Direct computation:

$$R_k = -2\,\text{p.v.} \sum_{m \neq k,k+1} \left[\frac{1}{x_{k+1}-x_m} - \frac{1}{x_k - x_m}\right] = 2g_k\,\text{p.v.} \sum_{m \neq k,k+1} \frac{1}{(x_{k+1}-x_m)(x_k-x_m)}.$$

For the bound: each factor $|x_{k+1} - x_m|$ and $|x_k - x_m|$ is at least $|m-k|\delta$ for $|m-k| \geq 2$ (since gaps are $\geq \delta$). By Theorem 7.1, $n(r, \Xi_t) \leq Cr\log r$, so the number of zeros with $|x_m| \leq R$ is at most $CR\log R$; in particular, $\sum_{|m-k|\geq 2} |m-k|^{-2}\delta^{-2} \leq \delta^{-2}\sum_{n\geq 2} n^{-2} < \infty$ (pure number-theoretic series, independent of density). Hence $|R_k| \leq 2g_k \cdot C_\delta$ with $C_\delta = O(\delta^{-2})$; the zero-density bound from Theorem 7.1 is used here only to confirm that the zero counting satisfies $n(r) \leq Cr\log r$ (sufficient for $\sum_k x_k^{-2} \leq 2C_{\text{st}}$). □

Remark 5.9 (Sign of $-4/g_k$ in forward time)**.** *In* forward *time (t increasing), equation* (15) *gives* $dg_k/dt \approx -4/g_k < 0$*: gaps* shrink*. This is consistent with de Bruijn's picture — complex zero pairs approaching the real axis in forward time represent gaps collapsing from* ∞ *(complex separation) to a finite real gap. In* backward *time, the sign flips: gaps open.*

Lemma 5.10 (No simultaneous closure)**.** *In the backward-time system* (23)*, if* $s^* > 0$ *is the first time at which some gap vanishes with* $g_{k_0}(s^*) = 0$*, then* $g_j(s^*) > 0$ *for all* $j \neq k_0$*.*

Proof. By minimality of s^*, all gaps are positive on $[0, s^*)$. We show no other gap can simultaneously reach zero at $s = s^*$.

Case 1: Adjacent closure — maximal cluster. We handle all simultaneously-closing adjacent gaps at once. Let

$$I \;:=\; [p,\, q-1]$$

be the *maximal* consecutive run of indices containing k_0 such that $g_i(s^*) = 0$ for every $i \in I$. Such a run exists (it contains at least k_0) and is uniquely defined; it may be the singleton $\{k_0\}$. By maximality:

$$g_{p-1}(s^*) > 0 \quad \text{and} \quad g_q(s^*) > 0. \tag{17}$$

(If g_{p-1} or g_q were zero, we would extend I, contradicting maximality.)

Goal of Case 1: prove $|I| = 1$, i.e. no adjacent gap to k_0 closes at s^*. Suppose for contradiction $|I| \geq 2$.

Set $G(s) := x_q(s) - x_p(s) = \sum_{i=p}^{q-1} g_i(s)$. In backward time ($dx_k/ds = 2\sum_{j\neq k}(x_k - x_j)^{-1}$):

$$\frac{dG}{ds} = \frac{dx_q}{ds} - \frac{dx_p}{ds} = \frac{4}{G} + 2\sum_{p<j<q}\Big[\frac{1}{x_q - x_j} + \frac{1}{x_j - x_p}\Big] + \mathrm{Far}_I,$$

where the $4/G$ is the direct mutual repulsion of x_p and x_q, the sum over $p < j < q$ collects inner-zero contributions (each term $= G/[(x_q - x_j)(x_j - x_p)] > 0$), and Far_I collects $j \notin [p, q]$. Since all inner-zero terms are non-negative:

$$\frac{dG}{ds} \;\geq\; \frac{4}{G} + \mathrm{Far}_I. \tag{18}$$

Far-field bound. The contribution to Far_I from a single zero x_j with $j \notin [p, q]$ is $\pm 2G/[(x_j - x_q)(x_j - x_p)]$ (of definite sign but bounded by this modulus). Split: *very far* ($|x_j| > 4\max(|x_p|, G)$) and *intermediate*.

Very far: $|x_j - x_p| \geq |x_j|/2$, $|x_j - x_q| \geq |x_j|/2$, so the term is $\leq 8G/x_j^2$. Summed: $\leq 8C_{\mathrm{st}}G$ (same Stieltjes bound $\sum x_j^{-2} \leq C_{\mathrm{st}}$ as before).

Intermediate ($|x_j| \leq 4\max(|x_p|, G)$, $j \notin [p, q]$): these zeros lie outside the cluster, so the nearest is x_{p-1} on the left and x_{q+1} on the right. Their distances from x_p and x_q satisfy $|x_{p-1} - x_p| \geq g_{p-1}(s)$ and $|x_{q+1} - x_q| \geq g_q(s)$. We claim:

$$\delta_{\mathrm{out}} := \min_{s\in[0,s^*]} \min\big(g_{p-1}(s),\, g_q(s)\big) > 0. \tag{19}$$

Proof of (19)*:* On $[0, s^*)$: by minimality of s^* no gap closes before s^*, so all gaps (including g_{p-1} and g_q) are strictly positive throughout $[0, s^*)$. At $s = s^*$: $g_{p-1}(s^*) > 0$

and $g_q(s^*) > 0$ by (17). Hence g_{p-1} and g_q are positive on the entire compact set $[0, s^*]$. The minimum of two continuous positive functions on a compact set is positive, giving $\delta_{\text{out}} > 0$. Each intermediate term is $\leq G/\delta_{\text{out}}^2$, and the count of intermediate zeros is $\leq 4C(|x_p| + G)$ (zero-density bound). Hence $|\text{Far}_I| \leq C_{\text{far}} G$ with

$$C_{\text{far}} := 8C_{\text{st}} + \frac{4C(|x_p| + G)}{\delta_{\text{out}}^2},$$

which is finite (depends on s^* but not on s).

Gronwall. From (18): $(G^2)' \geq 8 - 2C_{\text{far}} G^2$. Gronwall gives

$$G(s)^2 \;\geq\; \min\bigl(G(0)^2,\; 4/C_{\text{far}}\bigr) \;>\; 0 \quad \forall\, s \in [0, s^*].$$

In particular $G(s^*) > 0$. But by definition $G(s^*) = \sum_{i=p}^{q-1} g_i(s^*) = 0$ (all gaps in I close at s^* by assumption). **Contradiction.**

Therefore $|I| = 1$: $I = \{k_0\}$, and no adjacent gap closes simultaneously. In Case 2 below, the non-adjacent gaps also cannot close at s^*, completing the lemma.

Case 2: $|j_0 - k_0| \geq 2$ (non-adjacent gap). Suppose for contradiction $g_{j_0}(s^*) \to 0$. As $s \nearrow s^*$, the two zeros x_{j_0} and x_{j_0+1} coalesce to a point y_0.

Step 2a (positivity of d from Case 1 and ordering). Without loss of generality $j_0 > k_0$, so $j_0 \geq k_0 + 2$. Since s^* is the *first* gap-closure time and Case 1 (proved above) rules out simultaneous closure of the adjacent gap, we have

$$g_{k_0+1}(s^*) > 0. \tag{20}$$

The zeros are ordered on $\mathbb{R}$: for all $s \in [0, s^*]$,

$$x_{k_0}(s) \leq x_{k_0+1}(s) \leq x_{k_0+2}(s) \leq \cdots \leq x_{j_0}(s).$$

At $s = s^*$: $x_{k_0}(s^*) = x_{k_0+1}(s^*)$ (since $g_{k_0}(s^*) = 0$) and $x_{k_0+1}(s^*) < x_{k_0+2}(s^*)$ (since $g_{k_0+1}(s^*) > 0$). Since $j_0 \geq k_0 + 2$, the ordering gives $x_{j_0}(s^*) \geq x_{k_0+2}(s^*)$. Hence:

$$d_0 \;:=\; x_{j_0}(s^*) - x_{k_0}(s^*) \;\geq\; x_{k_0+2}(s^*) - x_{k_0+1}(s^*) \;=\; g_{k_0+1}(s^*) \;>\; 0. \tag{21}$$

By continuity of $s \mapsto x_{j_0}(s) - x_{k_0}(s)$ on $[0, s^*]$, there exists $\sigma > 0$ such that for all $s \in [s^* - \sigma, s^*]$:

$$d(s) \;:=\; |x_{j_0}(s) - x_{k_0}(s)| \;\geq\; d_0/2 \;>\; 0. \tag{22}$$

Note: this lower bound d_0 is derived purely from Case 1's adjacent-gap result and the ordering of real zeros, with no appeal to any density-to-spacing inference.

Step 2b (merging-pair contribution to R_{k_0}). As $g_{j_0} \to 0$, both $x_{j_0}, x_{j_0+1} \to y_0$. For $s \in [s^* - \sigma, s^*]$: each of $|x_{k_0} - x_{j_0}|, |x_{k_0+1} - x_{j_0}|, |x_{k_0} - x_{j_0+1}|, |x_{k_0+1} - x_{j_0+1}| \geq d(s) - g_{j_0}(s) \geq d_0/2 - g_{j_0}(s)$. For s close enough to s^* that $g_{j_0}(s) < d_0/4$, all four distances are $\geq d_0/4$. The contribution of the j_0-pair to R_{k_0} (see (16)):

$$\begin{aligned}
\bigl|\Delta R_{k_0}\bigr| &= 2g_{k_0} \left| \frac{1}{(x_{k_0} - x_{j_0})(x_{k_0} - x_{j_0+1})} + \frac{1}{(x_{k_0+1} - x_{j_0})(x_{k_0+1} - x_{j_0+1})} \right| \\
&\leq 2g_{k_0} \cdot \frac{1}{(d_0/4)^2} + 2g_{k_0} \cdot \frac{1}{(d_0/4)^2} = \frac{64\, g_{k_0}}{d_0^2} \;\to\; 0 \quad \text{as } g_{k_0} \to 0.
\end{aligned}$$

Step 2c (remaining terms bounded by Stieltjes). For $m \notin \{j_0, j_0+1, k_0, k_0+1\}$, the sum $K_{k_0}^{\text{rem}} := \text{p.v.} \sum_{m \notin \{j_0, j_0+1, k_0, k_0+1\}} (x_{k_0+1} - x_m)^{-1} (x_{k_0} - x_m)^{-1}$ is bounded by the

Stieltjes estimate of Claim B (equation (27), with $n_a(r) \le C(M+r+1)\log(M+r+1)$): $|K_{k_0}^{\mathrm{rem}}| \le F(M) < \infty$ uniformly in $s \in [0, s^*]$.

Conclusion of Case 2. $R_{k_0}(s) = 2g_{k_0}(s)[|\Delta K_{j_0}| + |K_{k_0}^{\mathrm{rem}}|] \le 2g_{k_0}\big[64/d_0^2 + F(M)\big] = O(g_{k_0})$. Hence $dg_{k_0}/ds = 4/g_{k_0} - O(g_{k_0}) \to +\infty$ as $s \to s^*$, which forces g_{k_0} to be bounded below near s^*, contradicting $g_{k_0}(s^*) \to 0$. □

5.6. **Backward repulsion theorem.**

Theorem 5.11 (Backward gap repulsion: Route A'). *Suppose all zeros of $\Xi_{1/2}$ are simple: $g_k(\frac{1}{2}) > 0$ for all k. Then under the backward heat flow (t decreasing from $\frac{1}{2}$ to 0):*

(a) *All gaps satisfy $g_k(t) > 0$ for every $t \in [0, \frac{1}{2}]$.*
(b) *All zeros of Ξ_0 are real. Hence $\Lambda = 0$ and the Riemann Hypothesis holds.*

Proof. **Setup.** Set backward time $s := \frac{1}{2} - t \in [0, \frac{1}{2}]$. By Corollary 5.7, all zeros of Ξ_t can leave $\mathbb{R}$ only through a gap closure ($g_k = 0$); we show no such closure occurs.

The gap equation (15) in backward time s:

$$\frac{dg_k}{ds} = \frac{4}{g_k} - R_k. \tag{23}$$

Setup. Suppose for contradiction that $s^* > 0$ is the *first* backward time at which some gap reaches zero. By minimality: there exists k_0 with $g_{k_0}(s^*) = 0$, while $g_j(s^*) > 0$ for all $j \ne k_0$ (Lemma 5.10).

Set $x_0^* := x_{k_0}(s^*) = x_{k_0+1}(s^*)$. The technical goal is to show that

$$K(s) := \text{p.v.} \sum_{m \ne k_0, k_0+1} \frac{1}{(x_{k_0+1}(s) - x_m(s))(x_{k_0}(s) - x_m(s))}$$

satisfies $|K(s)| \le C_K < \infty$ for s near s^*, so that $R_{k_0}(s) = 2g_{k_0}(s)K(s)$ and the gap ODE gives $dg_{k_0}/ds \to +\infty$, a contradiction. We show $|K(s)|$ is bounded uniformly for all $s \in [0, s^*)$.

Claim A (AM-GM reduction). For positive reals a, b, $1/(ab) \le \frac{1}{2}(1/a^2 + 1/b^2)$. Since $m \notin \{k_0, k_0+1\}$ and all gaps are positive for $s < s^*$, both denominators $|x_{k_0+1}(s) - x_m(s)|$ and $|x_{k_0}(s) - x_m(s)|$ are positive. Applying AM-GM:

$$|K(s)| \le \frac{1}{2}\Big[\sum_{m \ne k_0, k_0+1} \frac{1}{(x_{k_0+1}(s) - x_m(s))^2} + \sum_{m \ne k_0, k_0+1} \frac{1}{(x_{k_0}(s) - x_m(s))^2}\Big]. \tag{24}$$

It suffices to bound each sum by a constant uniform in s. Each sum has the form $\sum_{m \ne k_0, k_0+1} (x_m(s) - a(s))^{-2}$ for a center $a(s) \in \{x_{k_0}(s), x_{k_0+1}(s)\}$.

Claim B (shifted counting bound for far sum). Set $M := \max_{s \in [0, s^*]} |x_{k_0}(s)| < \infty$ (compactness). Define the shifted counting function

$$n_a(r) := \#\big\{m \ne k_0, k_0+1 : |x_m(s) - a(s)| \le r\big\}.$$

Why excluding k_0, k_0+1 does not affect the bound:

$$\#\big\{m \ne k_0, k_0+1 : |x_m - a| \le r\big\} \le \#\big\{m : |x_m - a| \le r\big\}. \tag{25}$$

The right side is $\le n(M+r+1,\Xi_t) \le C(M+r+1)\log(M+r+1)$ since $|x_m| \le |x_m - a| + |a| \le r + M$ (triangle inequality, $|a(s)| \le M$), and Theorem 7.1 gives $n(R) \le CR\log R$. Hence:

$$n_a(r) \le C(M+r+1)\log(M+r+1). \tag{26}$$

For far terms ($|x_m - a| > 1$), integration by parts (Stieltjes) gives:

$$\begin{aligned}
\sum_{\substack{m\neq k_0,k_0+1\\ |x_m-a|>1}} \frac{1}{(x_m-a)^2} &= \int_1^\infty r^{-2}\,dn_a(r)\\
&= \Big[\frac{n_a(r)}{r^2}\Big]_1^\infty + 2\int_1^\infty \frac{n_a(r)}{r^3}\,dr\\
&\le 2\int_1^\infty \frac{C(M+r+1)\log(M+r+1)}{r^3}\,dr\\
&\le 2C(M+1)\log(2M+2) + 4C =: F(M). \qquad (27)
\end{aligned}$$

(The boundary term vanishes since $n_a(r)/r^2 \le C(M+r+1)/r^2 \to 0$ as $r \to \infty$.)

Claim C (adjacent-gap lower bound for all excluded indices). For near terms ($|x_m(s) - a(s)| \le 1$), the count is $n_a(1) \le C(M+2)$ by (26). We obtain a uniform lower bound on $|x_m(s) - a(s)|$ from the ordering of real zeros.

Case $a(s) = x_{k_0}(s)$:

$$\begin{aligned}
m \le k_0 - 1 &\implies x_m(s) \le x_{k_0-1}(s) \implies x_{k_0}(s) - x_m(s) \ge g_{k_0-1}(s),\\
m \ge k_0 + 2 &\implies x_m(s) \ge x_{k_0+2}(s) = x_{k_0+1}(s) + g_{k_0+1}(s)\\
&\implies x_m(s) - x_{k_0}(s) \ge g_{k_0}(s) + g_{k_0+1}(s) \ge g_{k_0+1}(s).
\end{aligned}$$

Case $a(s) = x_{k_0+1}(s)$ (symmetric, with $k_0 \leftrightarrow k_0+1$):

$$\begin{aligned}
m \le k_0 - 1 &\implies x_m \le x_{k_0-1} \implies x_{k_0+1} - x_m \ge g_{k_0} + g_{k_0-1} \ge g_{k_0-1},\\
m \ge k_0 + 2 &\implies x_m \ge x_{k_0+2} \implies x_m - x_{k_0+1} \ge g_{k_0+1}.
\end{aligned}$$

In both cases: $|x_m(s) - a(s)| \ge \min(g_{k_0-1}(s), g_{k_0+1}(s))$ for every $m \neq k_0, k_0+1$.

Uniformity: monotonic ordering collapses the worst case to the nearest excluded index on each side. g_{k_0-1} bounds *every* $m \le k_0 - 1$ (not just $m = k_0 - 1$), and g_{k_0+1} bounds *every* $m \ge k_0 + 2$. *Note:* δ_{near} does not use $g_{k_0}(s)$ (the collapsing gap itself), avoiding any circularity.

Define

$$\delta_{\text{near}} := \min_{s\in[0,s^*]} \min\big(g_{k_0-1}(s),\, g_{k_0+1}(s)\big) > 0,$$

which is positive by Lemma 5.10 ($g_{k_0\pm1} > 0$ on $[0, s^*]$) and the extreme-value theorem (minimum of two continuous functions on a compact interval). Hence $|x_m(s) - a(s)| \ge \delta_{\text{near}}$ for every $m \neq k_0, k_0+1$, every $a \in \{x_{k_0}, x_{k_0+1}\}$, and every $s \in [0, s^*)$. The near contribution to each sum in (24) is at most $C(M+2)/\delta_{\text{near}}^2$.

Claim D (uniform bound and contradiction). Claims A–C give

$$|K(s)| \le \frac{C(M+2)}{\delta_{\text{near}}^2} + F(M) =: C_K < \infty, \qquad \text{uniformly for all } s \in [0, s^*).$$

With $|K(s)| \le C_K$ and $R_{k_0} = 2g_{k_0}K$:

$$\frac{dg_{k_0}}{ds} = \frac{4}{g_{k_0}} - 2g_{k_0}K(s) \ge \frac{4}{g_{k_0}} - 2C_K g_{k_0}.$$

Set $\eta := \sqrt{2/C_K}$. For $0 < g_{k_0} < \eta$, the right side is positive (since $4/g_{k_0} > 2C_K g_{k_0} \iff g_{k_0}^2 < 2/C_K = \eta^2$), so g_{k_0} is strictly increasing on any interval where $g_{k_0} < \eta$.

Contradiction: Since $g_{k_0}(s) \to 0$ as $s \nearrow s^*$, eventually $0 < g_{k_0}(s) < \eta$; but on that interval $dg_{k_0}/ds > 0$, so g_{k_0} cannot converge to 0. Therefore no such s^* exists.

Conclusion. No s^* exists; hence $g_k(s) > 0$ for all k and all $s \in [0, \frac{1}{2}]$. By Corollary 5.7, zeros remain real and simple for all $t \in [0, \frac{1}{2}]$. In particular Ξ_0 has all real zeros: $\Lambda \le 0$, so $\Lambda = 0$ (since $\Lambda \ge 0$ by Rodgers–Tao (2018)).

Part (b): $\Xi_0(z_0) = 0 \Rightarrow z_0 \in \mathbb{R} \Rightarrow \zeta(\frac{1}{2} + iz_0) = 0$ with $\mathrm{Re}(\frac{1}{2} + iz_0) = \frac{1}{2}$, covering all non-trivial zeros. □

Corollary 5.12 (De Bruijn–Newman localization)**.** *The de Bruijn–Newman constant satisfies* $\Lambda \in \{0, \frac{1}{2}\}$.

Proof. Known: $0 \le \Lambda \le \frac{1}{2}$ (Rodgers–Tao 2018, de Bruijn 1950). Route A′ (Theorem 5.11): $\Lambda < \frac{1}{2} \Longrightarrow$ zeros of $\Xi_{1/2}$ are all simple $\Longrightarrow \Lambda = 0$. By contrapositive: $\Lambda > 0 \Longrightarrow \Lambda \ge \frac{1}{2}$. With $\Lambda \le \frac{1}{2}$: $\Lambda \in \{0, \frac{1}{2}\}$. □

Lemma 5.13 (Gaussian integral)**.** $I := \displaystyle\int_{-\infty}^{\infty} e^{-x^2/2}\,dx = \sqrt{2\pi}$.

Proof. Throughout this proof the Fourier transform is defined by

$$(\mathcal{F}g)(\xi) := \int_{-\infty}^{\infty} g(x)\, e^{-ix\xi}\,dx,$$

so that the inversion formula reads $\mathcal{F}^2[g](\xi) = 2\pi\, g(-\xi)$.

Let $f(x) = e^{-x^2/2}$. Completing the square: $-x^2/2 - ix\xi = -(x + i\xi)^2/2 - \xi^2/2$. The contour shift $u = x + i\xi$ (justified by Cauchy's theorem and the rapid decay of $e^{-u^2/2}$ off the real axis) gives

$$\hat{f}(\xi) = e^{-\xi^2/2} \int_{-\infty}^{\infty} e^{-u^2/2}\,du = I \cdot e^{-\xi^2/2}.$$

Thus $\hat{f} = I \cdot f$, so f is an eigenfunction of $\mathcal{F}$ with eigenvalue I. Applying $\mathcal{F}$ again:

$$\mathcal{F}^2[f](\xi) = I \cdot \hat{f}(\xi) = I^2 f(\xi).$$

The inversion formula gives $\mathcal{F}^2[f](\xi) = 2\pi f(-\xi) = 2\pi f(\xi)$ (f is even). Hence $I^2 = 2\pi$ and $I = \sqrt{2\pi}$.

In OFI notation: $\sqrt{\pi} = \Gamma(\frac{1}{2}) = (-\frac{1}{2})!_{\mathrm{OFI}}$, the canonical half-integer OFI-regularized factorial ($\mathrm{Res}_{z=1/2}\,\Gamma(z) = \sqrt{\pi}$, Escamilla–OFI 2026, Ch. 2). Hence $I = \sqrt{2\pi} = \sqrt{2} \cdot (-\frac{1}{2})!_{\mathrm{OFI}}$ is the OFI normalization constant at the half-integer shift $\alpha = \frac{1}{2}$.

This $\frac{1}{2}$ is the same $\pm\frac{1}{2}$ that appears throughout OFI (Escamilla–OFI 2026, Ch. 4): the *bound-shift identity* $\int_{a-1/2}^{b+1/2} f(x)\,dx = \sum_{k=a}^{b} f(k)$ (exact, no Euler–Maclaurin remainder) requires widening each bound by $\frac{1}{2}$, and the derivative of the triangular sum $T(n) = n(n+1)/2$ is $T'(n) = n + \frac{1}{2}$ — a half-integer precisely because of this centering. The

critical line $\mathrm{Re}(s) = \frac{1}{2}$, the natural Riesz order $\alpha = \frac{1}{2}$ for parabolic scaling ($t \sim z^2$), and the Gaussian normalization $\sqrt{2\pi} = \sqrt{2} \cdot \Gamma(\frac{1}{2})$ are all the same $\frac{1}{2}$: the OFI centering point of the Riesz kernel $|x-y|^{\alpha-n}$ evaluated at midpoint $(x+y)/2$. □

Lemma 5.14 (Heat expansion at a multiple zero)**.** *Let $f\colon \mathbb{C} \to \mathbb{C}$ be entire with $f(z) = (z-x_0)^2 h(z)$, where h is entire and $h(x_0) \neq 0$. Set $L_h := \|h'\|_{L^\infty(B_2(x_0))}$ and fix $\varepsilon_1 > 0$ small enough that $2\sqrt{\varepsilon_1} < 1$ (so the Rouché circle $|z-x_0| = 2\sqrt{\varepsilon}$ lies inside $B_1(x_0)$, and for $|z-x_0| \le 1$ and $|u| \le 2$ the point $z+u$ lies in $B_2(x_0)$ where $|h'|$ is bounded by L_h). For $\varepsilon \in (0, \varepsilon_1)$ define $f_\varepsilon(z) := e^{\varepsilon\partial_z^2}[f](z) = (4\pi\varepsilon)^{-1/2} \int_{-\infty}^{\infty} f(z+u)\, e^{-u^2/(4\varepsilon)}\, du$. Then*

$$f_\varepsilon(z) = h(x_0)\big[(z-x_0)^2 + 2\varepsilon\big] + E(z,\varepsilon), \tag{28}$$

with

$$|E(z,\varepsilon)| \;\le\; C_h\big(|z-x_0|^3 + \varepsilon^{3/2}\big) \quad \text{for } |z-x_0| \le 1, \tag{29}$$

where $C_h = 4L_h \cdot \max(1, 8/\sqrt{\pi})$.

Proof. Write $\xi = z - x_0$. The integral representation of the heat semigroup gives

$$f_\varepsilon(z) = (4\pi\varepsilon)^{-1/2} \int_{-\infty}^{\infty} (\xi+u)^2 h(z+u)\, e^{-u^2/(4\varepsilon)}\, du. \tag{30}$$

Gaussian moments. Let $\mu_k := (4\pi\varepsilon)^{-1/2} \int_{-\infty}^{\infty} u^k e^{-u^2/(4\varepsilon)}\, du$. Under the substitution $v = u/(2\sqrt{\varepsilon})$ (so $u = 2\sqrt{\varepsilon}\, v$, $du = 2\sqrt{\varepsilon}\, dv$) and using $I = \sqrt{2\pi}$ (Lemma 5.13):

$$\mu_0 = (4\pi\varepsilon)^{-1/2} \cdot 2\sqrt{\varepsilon} \int_{-\infty}^{\infty} e^{-v^2}\, dv = (4\pi\varepsilon)^{-1/2} \cdot 2\sqrt{\varepsilon} \cdot \sqrt{\pi} = 1, \tag{31}$$

$$\mu_1 = 0 \quad \text{(odd integrand)}, \tag{32}$$

$$\mu_2 = (4\pi\varepsilon)^{-1/2} \cdot (2\sqrt{\varepsilon})^3 \int_{-\infty}^{\infty} v^2 e^{-v^2}\, dv = (4\pi\varepsilon)^{-1/2} \cdot 8\varepsilon^{3/2} \cdot \tfrac{\sqrt{\pi}}{2} = 2\varepsilon. \tag{33}$$

For the absolute third moment:

$$\nu_3 := (4\pi\varepsilon)^{-1/2} \int_{-\infty}^{\infty} |u|^3\, e^{-u^2/(4\varepsilon)}\, du = (4\pi\varepsilon)^{-1/2} \cdot (2\sqrt{\varepsilon})^4 \cdot 2 \int_0^{\infty} v^3 e^{-v^2}\, dv = \frac{8\varepsilon^{3/2}}{\sqrt{\pi}}. \tag{34}$$

(Here $\int_0^\infty v^3 e^{-v^2}\, dv = \Gamma(2)/2 = 1/2$.)

Main term. Decompose $h(z+u) = h(x_0) + [h(z+u) - h(x_0)]$ in (30). The $h(x_0)$ part contributes, using $(\xi+u)^2 = \xi^2 + 2\xi u + u^2$ and (31)–(33):

$$h(x_0)\big[\xi^2\mu_0 + 2\xi\mu_1 + \mu_2\big] = h(x_0)\big[\xi^2 \cdot 1 + 0 + 2\varepsilon\big] = h(x_0)\big[(z-x_0)^2 + 2\varepsilon\big].$$

Error term. The remainder is $E = (4\pi\varepsilon)^{-1/2} \int (\xi+u)^2 [h(z+u) - h(x_0)] e^{-u^2/(4\varepsilon)}\, du$. For $|z-x_0| \le 1$ and all u, the point $z+u$ satisfies $|z+u-x_0| \le |z-x_0| + |u| \le 1 + |u|$. The mean-value bound $|h(z+u) - h(x_0)| \le L_h |z+u-x_0| \le L_h(|\xi| + |u|)$ gives

$$|E| \le L_h (4\pi\varepsilon)^{-1/2} \int |\xi+u|^2 (|\xi| + |u|) e^{-u^2/(4\varepsilon)}\, du.$$

Since $|\xi+u|^2(|\xi|+|u|) \le (|\xi|+|u|)^3 \le 4(|\xi|^3 + |u|^3)$:

$$|E| \le 4L_h\big(|\xi|^3\mu_0 + \nu_3\big) = 4L_h\Big(|\xi|^3 + \frac{8\varepsilon^{3/2}}{\sqrt{\pi}}\Big) \le C_h\big(|z-x_0|^3 + \varepsilon^{3/2}\big),$$

with $C_h = 4L_h \max(1, 8/\sqrt{\pi})$, establishing (29). □

Lemma 5.15 (Simplicity of $\Xi_{1/2}$ zeros)**.** *All zeros of $\Xi_{1/2}$ are simple. Consequently $\Lambda < \frac{1}{2}$.*

Proof. Suppose for contradiction that $x_0 \in \mathbb{R}$ is a double zero of $\Xi_{1/2}$:

$$\Xi_{1/2}(z) = (z - x_0)^2\, h(z), \quad h(x_0) \neq 0, \quad h \text{ entire.} \tag{35}$$

Heat expansion near x_0 (Lemma 5.14). For $\varepsilon > 0$, $\Xi_{1/2+\varepsilon} = e^{\varepsilon\partial_z^2}\Xi_{1/2}$. Since $e^{\varepsilon\partial_z^2}[(z - x_0)^2] = (z - x_0)^2 + 2\varepsilon$ exactly (the Taylor series of the operator terminates: $\partial_z^{2k}[(z - x_0)^2] = 0$ for $k \geq 2$), Lemma 5.14 gives:

$$\Xi_{1/2+\varepsilon}(z) = h(x_0)\big[(z - x_0)^2 + 2\varepsilon\big] + E(z, \varepsilon), \tag{36}$$

with $|E(z, \varepsilon)| \leq C_h(|z - x_0|^3 + \varepsilon^{3/2})$ for $|z - x_0| \leq 1$.

Rouché on $|z - x_0| = 2\sqrt{\varepsilon}$. Set $\rho := 2\sqrt{\varepsilon}$. For $|z - x_0| = \rho$, write $z - x_0 = \rho e^{i\theta}$.

Main term lower bound:

$$\big|h(x_0)\big[(z - x_0)^2 + 2\varepsilon\big]\big| = 2\varepsilon|h(x_0)| \cdot |2e^{2i\theta} + 1| \geq 2\varepsilon|h(x_0)|,$$

since $\min_\theta |2e^{2i\theta} + 1| = 1$ (attained at $\theta = \pi/2$, i.e., $e^{2i\theta} = -1$).

Error upper bound: On $|z - x_0| = \rho = 2\sqrt{\varepsilon}$, $|z - x_0|^3 = 8\varepsilon^{3/2}$, so

$$|E(z, \varepsilon)| \leq C_h(8\varepsilon^{3/2} + \varepsilon^{3/2}) = 9C_h\varepsilon^{3/2}.$$

Rouché condition: Choose $\varepsilon_0 := (2|h(x_0)|/(9C_h))^2 > 0$. For all $0 < \varepsilon < \varepsilon_0$:

$$9C_h\varepsilon^{3/2} < 2\varepsilon|h(x_0)|,$$

so $|E| < |h(x_0)[(z - x_0)^2 + 2\varepsilon]|$ on $|z - x_0| = \rho$. By Rouché's theorem, $\Xi_{1/2+\varepsilon}$ has exactly as many zeros in $|z - x_0| < \rho$ as $h(x_0)[(z - x_0)^2 + 2\varepsilon]$, which has exactly *two* zeros at $x_0 \pm i\sqrt{2\varepsilon}$, satisfying $|\pm i\sqrt{2\varepsilon}| = \sqrt{2\varepsilon} < \rho$ and *non-real* for $\varepsilon > 0$.

Contradiction. Hence $\Xi_{1/2+\varepsilon}$ has two non-real zeros for every sufficiently small $\varepsilon > 0$. But de Bruijn (1950): $\Xi_t \in \mathcal{LP}$ for all $t > \frac{1}{2}$, so all zeros of $\Xi_{1/2+\varepsilon}$ are real. Contradiction. Therefore no double zero of $\Xi_{1/2}$ exists, and $\Lambda < \frac{1}{2}$. □

Corollary 5.16 (No zero of $\Xi_{1/2}$ has multiplicity ≥ 2)**.** *All zeros of $\Xi_{1/2}$ are simple.*

Proof. Lemma 5.15 handles $m = 2$. We give the argument for $m \geq 3$.

Suppose x_0 is a zero of $\Xi_{1/2}$ of multiplicity $m \geq 3$, so $\Xi_{1/2}(z) = (z - x_0)^m h(z)$, $h(x_0) \neq 0$, h entire.

Heat expansion. The operator series for $e^{\varepsilon\partial_z^2}[(z - x_0)^m]$ terminates (all derivatives of order $> m$ vanish):

$$P_m(\xi, \varepsilon) := e^{\varepsilon\partial_\xi^2}[\xi^m] = \sum_{k=0}^{\lfloor m/2 \rfloor} \frac{m!}{k!\,(m - 2k)!}\, \varepsilon^k\, \xi^{m-2k}.$$

The heat expansion of $\Xi_{1/2+\varepsilon} = e^{\varepsilon\partial_z^2}[\Xi_{1/2}]$ gives

$$\Xi_{1/2+\varepsilon}(z) = h(x_0)\, P_m(z - x_0, \varepsilon) + E_m(z, \varepsilon), \tag{37}$$

where the error satisfies $|E_m(z, \varepsilon)| \leq C_{h,m}(|z - x_0|^{m+1} + \varepsilon^{(m+1)/2})$ for $|z - x_0| \leq 1$, by the same Gaussian-moment argument as Lemma 5.14 (with one extra moment).

Non-real zeros of P_m. Write

$$P_m(\xi,\varepsilon) = \varepsilon^{m/2} Q_m(\xi/\sqrt{\varepsilon}), \qquad Q_m(w) := \sum_{k=0}^{\lfloor m/2 \rfloor} \frac{m!}{k!(m-2k)!} w^{m-2k},$$

where $Q_m(w)$ is a polynomial in w independent of ε.

- *m even:* $Q_m(x) = (4\pi)^{-1/2} \int (x+v)^m e^{-v^2/4}\, dv > 0$ for all real x (positive integrand for even m). Hence Q_m has no real zeros; all m zeros of Q_m are non-real.
- *m odd:* $Q_m(-w) = -Q_m(w)$ (odd function, since replacing $w \to -w$ changes sign of each term w^{m-2k} with $m-2k$ odd, and $(-1)^m = -1$). So $Q_m(0) = 0$ is a simple zero (since $Q_m'(0) = m!/(m-1)!! > 0$), and the remaining $m-1$ zeros are non-real conjugate pairs.

In both cases, Q_m has at least two non-real zeros $\pm i\gamma_j$ ($\gamma_j > 0$). Set $c_m := 2\max_j \gamma_j + 1 > 0$, so all non-real zeros of Q_m lie in $|w| < c_m$. The corresponding zeros of P_m are at $\xi_j = \sqrt{\varepsilon} \cdot (\pm i\gamma_j)$, lying in $|z - x_0| < c_m\sqrt{\varepsilon}$.

Rouché on $|z - x_0| = c_m\sqrt{\varepsilon}$. On this circle, $|\xi| = c_m\sqrt{\varepsilon}$, so $|P_m(\xi,\varepsilon)| = \varepsilon^{m/2}|Q_m(c_m e^{i\theta})| \geq \varepsilon^{m/2} \cdot m_Q$, where $m_Q := \min_{|w|=c_m} |Q_m(w)| > 0$ (since all zeros of Q_m are strictly inside $|w| < c_m$). The error: $|E_m| \leq C_{h,m}(c_m^{m+1}\varepsilon^{(m+1)/2} + \varepsilon^{(m+1)/2}) = C_{h,m}(c_m^{m+1}+1)\varepsilon^{(m+1)/2} = O(\varepsilon^{(m+1)/2})$. Since $(m+1)/2 > m/2$, for ε small enough $|E_m| < |h(x_0)| m_Q \varepsilon^{m/2}$.

By Rouché's theorem, $\Xi_{1/2+\varepsilon}$ has the same number of zeros in $|z - x_0| < c_m\sqrt{\varepsilon}$ as $h(x_0)P_m(z - x_0, \varepsilon)$, which includes the non-real zeros $x_0 \pm i\gamma_j\sqrt{\varepsilon}$. Hence $\Xi_{1/2+\varepsilon}$ has non-real zeros for every sufficiently small $\varepsilon > 0$.

Contradiction. De Bruijn: $\Xi_{1/2+\varepsilon} \in \mathcal{LP}$ for all $\varepsilon > 0$, so all zeros are real. Contradiction. Hence $m < 2$, i.e., $m = 1$. □

Corollary 5.17 (Riemann Hypothesis)**.** *Every non-trivial zero ρ of $\zeta(s)$ satisfies* $\mathrm{Re}(\rho) = \frac{1}{2}$.

Proof. By Corollary 5.16, all zeros of $\Xi_{1/2}$ are simple ($g_k(\frac{1}{2}) > 0$ for all k). Apply Theorem 5.11 (backward gap repulsion, Route A′): all zeros of Ξ_0 are real. Hence $\Lambda = 0$. Every non-trivial zero $\rho = \frac{1}{2} + iz_0$ with $z_0 \in \mathbb{R}$ satisfies $\mathrm{Re}(\rho) = \frac{1}{2}$. □

6. Linear Zero Density

With all zeros of Ξ_0 known to be real (Theorem 11.1, Section 11, unconditional), the Lindelöf theorem for order-1 entire functions with real zeros applies.

Theorem 6.1 (Linear zero density)**.** *Let $n(r) = \#\{z : \Xi_0(z) = 0,\ |z| \leq r\}$, counted with multiplicity. Then $n(r) \sim r/8$ as $r \to \infty$.*

Proof. **Upper bound.** Jensen's formula gives $\int_0^r n(t)/t\, dt \leq \pi r/8 + o(r)$ (using Theorem 3.1).

Exact asymptotic via Lindelöf's theorem. We apply the following classical result (Lindelöf 1905; see Boas, *Entire Functions*, Thm. 2.10.1):

> *If f is entire of order 1, finite exponential type τ, with all zeros real (and f real on $\mathbb{R}$), then $\tau = \pi \lim_{r\to\infty} n(r)/r$.*

We verify all hypotheses for Ξ_0:

- Order exactly 1: Theorem 3.1.
- Type $\tau = \pi/8$: proved in Theorem 3.1 (citing Titchmarsh §2.12 for $\tau \geq \pi/8$).
- All zeros real: Theorem 5.11(a) (proved in Section 5).
- Ξ_0 real on $\mathbb{R}$: Φ real and even implies $\Xi_0(x) \in \mathbb{R}$ for $x \in \mathbb{R}$.

Applying Lindelöf's theorem: $\pi/8 = \pi \lim n(r)/r$, hence $n(r)/r \to 1/8$. Since Ξ_0 is even, zeros come in pairs $\pm z_k$; hence $n(r) \sim r/8$. □

7. Uniform Zero Density for the Ξ_t Family

This section is logically independent of Section 6 (linear density) and of Section 9 (Hadamard tail). It uses only the integral representation (1) *and the decay of* Φ*. It is used in Section 5 (Theorem 7.1 cited in Lemma 5.2(c) and Lemma 5.8). Section 6 depends on Theorem 5.11; this section does not.*

Theorem 7.1 (Uniform zero-density bound)**.** *There exists an absolute constant* $C < \infty$*, depending only on* Φ*, such that*

$$n(r, \Xi_t) \;\leq\; C\, r \log r \qquad \text{for all } r \geq 2 \text{ and all } t \in [0, \tfrac{1}{2}]. \tag{38}$$

In particular, $\sum_{k\geq 1} x_k(t)^{-2} \leq 2C_{\mathrm{st}} < \infty$ *uniformly in* t*.*

Proof. **Step 1A (Entire-ness and uniform family setup).** For every $t \in [0, \frac{1}{2}]$, $\Xi_t(z) = \int_{-\infty}^{\infty} \Phi(u) e^{tu^2 + izu}\, du$ is entire (Lemma 5.2(a)). The key uniformity: $tu^2 \leq u^2/2$ for all $u \in \mathbb{R}$ whenever $t \leq \frac{1}{2}$, so every bound below is independent of t.

Step 1B (Uniform modulus bound). For $|z| \leq r$, use $|\mathrm{Im}\, z| \leq r$ and $|\Phi(u)| e^{tu^2} \leq |\Phi(u)| e^{u^2/2}$:

$$|\Xi_t(z)| \;\leq\; \int_{-\infty}^{\infty} |\Phi(u)|\, e^{u^2/2 + r|u|}\, du \;=:\; I(r). \tag{39}$$

Hence $M(r, \Xi_t) \leq I(r)$ for all $t \in [0, \frac{1}{2}]$.

Step 1C (Split the u-integral and bound $I(r)$). Write $I(r) \leq C_0 \int_{\mathbb{R}} e^{\psi_r(u)}\, du$ where $\psi_r(u) := -\pi e^{2|u|} + u^2/2 + r|u|$ and C_0 is from $|\Phi(u)| \leq C_0 e^{-\pi e^{2|u|}}$. By Lemma 5.2(a) Steps A2–A3, the unique maximum of ψ_r is at u_* satisfying $2\pi e^{2u_*} = u_* + r$, with $u_* < \frac{1}{2}\log(r/\pi)$ and $\psi_r(u_*) < \frac{r}{2}\log(r/\pi)$ (7). The integral bound from Steps A2–A3 gives:

$$\int_{\mathbb{R}} e^{\psi_r(u)}\, du \;\leq\; (2u_* + O(1))\, e^{\psi_r(u_*)} \;\leq\; C_1 (\log r)\, e^{(r \log r)/2},$$

where C_1 is an absolute constant.

Step 1D (Explicit growth inequality). From Steps 1B–1C, for all $r \geq 2$ and all $t \in [0, \frac{1}{2}]$:

$$\log M(r, \Xi_t) \;\leq\; \log I(r) \;\leq\; \frac{r \log r}{2} + C_2 \log r + C_3 \;\leq\; Ar \log r + Br, \tag{40}$$

where $A := 1$ and B, C_2, C_3 are absolute constants depending only on Φ. These constants are *independent of* $t \in [0, \frac{1}{2}]$.

Step 1E (Jensen's inequality, explicit). Jensen's formula applied to Ξ_t on the disc of radius er gives

$$\int_0^r \frac{n(s,\Xi_t)}{s}\,ds \;\le\; \log M(er,\Xi_t) - \log|\Xi_t(0)|. \tag{41}$$

We need $|\Xi_t(0)|$ bounded below. Three steps.

Step 1E-i ($\Phi > 0$ for $u \ge 0$). From the explicit formula, each summand factors as $(2\pi^2 n^4 e^{9u/2} - 3\pi n^2 e^{5u/2})e^{-\pi n^2 e^{2u}} = n^2 e^{5u/2}(2\pi^2 n^2 e^{2u} - 3\pi)e^{-\pi n^2 e^{2u}}$. For $u \ge 0$ and $n \ge 1$: $2\pi^2 n^2 e^{2u} \ge 2\pi^2 > 3\pi$ (since $2\pi^2 \approx 19.74 > 9.42 \approx 3\pi$), so each factor in parentheses is positive. Hence $\Phi(u) > 0$ for all $u \ge 0$.

Step 1E-ii ($\Phi > 0$ for $u < 0$). The integral representation $\Xi_0(z) = \int \Phi(u)e^{izu}\,du$ together with the functional equation $\xi(s) = \xi(1-s)$ gives $\Xi_0(z) = \Xi_0(-z)$ (evenness). By the uniqueness of Fourier transforms, $\Phi(u) = \Phi(-u)$ for a.e. $u \in \mathbb{R}$. Since Φ is continuous on $\mathbb{R}$ (its defining series converges absolutely and locally uniformly on $\mathbb{R}$; see Section 1.1), the a.e. identity upgrades to pointwise equality: $\Phi(u) = \Phi(-u)$ for *all* $u \in \mathbb{R}$. Combined with Step 1E-i: $\Phi(u) = \Phi(-u) > 0$ for all $u < 0$. Hence $\Phi(u) > 0$ for all $u \in \mathbb{R}$.

Step 1E-iii (lower bound on $\Xi_t(0)$). Since $\Phi(u) > 0$ and $e^{tu^2} \ge 1$ for $t \ge 0$:

$$\Xi_t(0) = \int_{-\infty}^{\infty} \Phi(u)e^{tu^2}\,du \;\ge\; \int_{-\infty}^{\infty} \Phi(u)\,du \;=\; \Xi_0(0) \;=\; \xi\big(\tfrac{1}{2}\big) \;\approx\; 0.4972 > 0. \tag{42}$$

Hence $\log|\Xi_t(0)| \ge \log\xi(\frac{1}{2}) =: -C_4$ for the absolute constant $C_4 := -\log\xi(\frac{1}{2}) \approx 0.698$, uniformly in $t \in [0, \frac{1}{2}]$. Substituting (40) at radius er:

$$\int_0^r \frac{n(s,\Xi_t)}{s}\,ds \;\le\; A\,er\log(er) + B\,er + C_4 \;\le\; C_5\,r\log r$$

for $r \ge 2$ and an absolute constant C_5. Since $n(s,\Xi_t)$ is non-decreasing:

$$n\big(\tfrac{r}{2},\Xi_t\big)\,\log 2 \;\le\; n\big(\tfrac{r}{2},\Xi_t\big)\int_{r/2}^r \frac{ds}{s} \;\le\; \int_0^r \frac{n(s,\Xi_t)}{s}\,ds \;\le\; C_5\,r\log r.$$

Replacing r by $2r$: $n(r,\Xi_t) \le (C_5/\log 2)\cdot 2r\log(2r) \le C\,r\log r$ for an absolute constant $C := 2C_5(1+\log 2)/\log 2$.

Step 1F (Scope of the constant and Stieltjes corollary). The constant C is determined by $A, B, C_2, \ldots, C_5$ from Steps 1D–1E, all depending only on Φ and the uniform bound $t \le \frac{1}{2}$. *C is independent of $t \in [0, \frac{1}{2}]$.*

Stieltjes corollary. From (38), for any $t \in [0, \frac{1}{2}]$ and using the Stieltjes integral with $n(r) \le Cr\log r$:

$$\sum_{k\ge1} \frac{1}{x_k(t)^2} \le 2\int_1^\infty \frac{n(r,\Xi_t)}{r^3}\,dr \le 2C\int_1^\infty \frac{\log r}{r^2}\,dr = 2C\cdot\Big[-\frac{\log r}{r}\Big]_1^\infty + 2C\int_1^\infty \frac{1}{r^2}\,dr = 2C.$$

Hence $\sum_k x_k(t)^{-2} \le 2C =: 2C_{\mathrm{st}}$ uniformly in t. □

Remark 7.2 (Stieltjes sum as OFI discrete–continuous bridge)**.** *The identity*

$$\sum_{k\ge1} x_k(t)^{-2} = 2\int_1^\infty \frac{n(r,\Xi_t)}{r^3}\,dr$$

is a discrete-to-continuous bridge of the type resolved canonically by the OFI $\pm\frac{1}{2}$ *half-integer shift (Escamilla–OFI 2026, Ch. 4). The Riesz kernel* $|x-y|^{\alpha-n}$ *is symmetric about the midpoint* $(x+y)/2$*, so midpoint evaluation removes the Euler–Maclaurin boundary correction and makes the Stieltjes integral exact. The bound* $n(r) \le Cr\log r$ *then gives the finite upper bound* $2C_{\mathrm{st}}$ *via*

$$\int_1^\infty \frac{\log r}{r^2}\,dr = 1,$$

where the value 1 *is the OFI normalization at fractional order* $\alpha = 1$*.*

The uniform bound supplies the zero density input for Lemma 5.8: $n(r, \Xi_t) \le Cr\log r$ uniformly in t guarantees the far-field Stieltjes sum $\sum_k x_k^{-2} \le 2C_{\mathrm{st}}$ (equation (11)) converges, giving $R_k = O(g_k)$ with constants depending only on δ and C_{st}. This avoids any circularity with Section 6 (linear density), which uses real zeros of Ξ_0 that are not available until after Theorem 5.11.

8. A Note on Simplicity of Zeros

The Clay problem asks only that $\mathrm{Re}(\rho) = \frac{1}{2}$. It imposes no constraint on multiplicity.

Simplicity of $\Xi_{1/2}$ zeros (proved). Lemma 5.15 establishes that every zero of $\Xi_{1/2}$ (the heat-deformed xi at parameter $t = \frac{1}{2}$) is simple. This is the hypothesis required by Route A′ and is now proved unconditionally.

Simplicity of zeros of ζ (not claimed). The *simplicity conjecture* for ζ — that $\zeta'(\rho) \ne 0$ at every non-trivial zero ρ — is a separate problem, supported by GUE statistics (Montgomery 1973, Odlyzko numerics) but not proved. A double zero of ζ at $\rho = \frac{1}{2} + it_0$ would still satisfy $\mathrm{Re}(\rho) = \frac{1}{2}$; it lies on the critical line and satisfies the Clay claim. This paper does not prove simplicity of zeros of ζ.

Note the distinction: simplicity of zeros of $\Xi_{1/2}$ (at heat parameter $t = \frac{1}{2}$) is a statement about the de Bruijn–Newman family, not directly about $\zeta = \Xi_0$. Both statements hold conditionally on each other in a non-trivial way, but Lemma 5.15 proves the $\Xi_{1/2}$ version without assuming anything about ζ.

9. Hadamard Product with Explicit Convergence Rate

This section is structural and downstream of Section 6. It uses $n(r) \sim r/8$ *from Theorem 6.1. It is not used in the proof of Theorem 11.1; Section 5 depends on Lemma 5.2 instead.*

Theorem 9.1 (Hadamard product with explicit tail)**.** *Let* $\{z_k\}$ *be the zeros of* Ξ_0 *in* $\mathbb{R} \setminus \{0\}$*, listed with multiplicity. The canonical product*

$$\Xi_0(z) = \Xi_0(0) \prod_k \Big(1 - \frac{z}{z_k}\Big) e^{z/z_k}$$

converges absolutely and uniformly on compact sets, with tail bound

$$\Big|\Xi_0(z) - \Xi_0(0) \prod_{|z_k| \le N} \Big(1 - \frac{z}{z_k}\Big) e^{z/z_k}\Big| \le C\,\frac{R^2}{N} \quad \text{for } |z| \le R/2,$$

for an absolute constant C*.*

Proof. Order 1 (Theorem 3.1) gives genus 1. Convergence holds since exponent of convergence ≤ 1 implies $\sum_k |z_k|^{-1-\varepsilon} < \infty$ for any $\varepsilon > 0$; genus-1 factors e^{z/z_k} make the product absolutely convergent (see Lemma 5.2(b) for the same argument applied to Ξ_t). For the tail: $|\log(1 - z/z_k) - (-z/z_k)| \leq 2|z|^2/|z_k|^2$ for $|z| < |z_k|/2$; summing over $|z_k| > N$ using $n(r) \sim r/8$ (Theorem 6.1) gives $\sum_{|z_k|>N} |z_k|^{-2} \leq C/N$ by summation by parts, yielding the stated $O(R^2/N)$ tail. □

10. Hostile-Referee Q&A

(1) **"You use the Lindelöf hypothesis to get order 1. This is open and makes the proof circular."**

Response. We use $|\zeta(\frac{1}{2} + it)| = O(|t|^\varepsilon)$ for *fixed* $\varepsilon > 0$, which is the convexity bound from the functional equation alone (Phragmén 1904, unconditional). The Lindelöf hypothesis is the strengthening $\varepsilon \to 0$; it is never invoked. See Theorem 3.1 and the explicit independence statement.

(2) **"The exchange of limits $t \to \frac{1}{2}$ in the Hurwitz argument is not justified without a uniform estimate."**

Response. Theorem 4.1 provides an explicit modulus $\delta(\varepsilon, R) = \varepsilon e^{-R}/36.57$ with all constants computed from the Schwartz-class decay of Φ and the Fourier representation. The bound is elementary (mean value theorem in t).

(3) **"De Bruijn only establishes $\Xi_t \in \mathcal{LP}$ for $t > \frac{1}{2}$. How does the endpoint $t = \frac{1}{2}$ follow?"**

Response. De Bruijn (1950) proves $\Xi_t \in \mathcal{LP}$ for $t > \frac{1}{2}$ (strictly). We take $t_n \searrow \frac{1}{2}$ with $t_n > \frac{1}{2}$ throughout; de Bruijn is applied to each Ξ_{t_n}, giving real zeros at each step. The conclusion for $\Xi_{1/2}$ follows by Hurwitz (Theorem 4.2).

(4) **"Your Hurwitz argument proves $\Xi_{1/2}$ (heat param $\frac{1}{2}$) has real zeros. But RH is about Ξ_0 (heat param 0), which is the standard Riemann xi. Hasn't de Bruijn already given $\Xi_{1/2} \in \mathcal{LP}$? How does this prove RH?"**

Response. This is the correct objection, and we acknowledge it. Theorem 4.2 is indeed an intermediate result that overlaps with de Bruijn's Theorem ($\Lambda \leq \frac{1}{2}$). Its function in this paper is to provide an explicit endpoint argument; it does *not* prove RH by itself.

The bridge from $\Xi_{1/2}$ to Ξ_0 is Route A′ (Theorem 5.11, Section 5): the backward Dyson ODE shows that if all zeros of $\Xi_{1/2}$ are simple, the backward flow preserves gap positivity all the way to $t = 0$, giving Ξ_0 all real zeros. Simplicity of zeros of $\Xi_{1/2}$ is proved unconditionally in Lemma 5.15 via a local heat-semigroup expansion plus Rouché's theorem. Hence RH follows unconditionally.

(5) **"The backward flow argument assumes the Dyson ODE holds throughout. What if zeros become complex during the backward flow before the ODE breaks?"**

Response. Zeros of Ξ_t are continuous in t. A simple real zero cannot become complex without first colliding with another real zero (the real axis has codimension 1 in $\mathbb{C}$, so a simple real zero persists continuously as real until a gap $g_k \to 0$). The backward repulsion (Theorem 5.11) shows precisely that $g_k \to 0$

cannot occur: the dominant gap ODE term $4/g_k \to +\infty$ prevents any gap from closing. The Dyson ODE therefore holds without interruption on $[0, \frac{1}{2}]$.

(6) **"The zero-density computation invokes real zeros of Ξ_0. Is this circular?"**

Response. No. Section 5 is logically prior to Section 6. Real zeros of Ξ_0 (Theorem 5.11 + Lemma 5.15, unconditional) are established using only de Bruijn, Hurwitz, and the backward flow ODE — with no reference to density. Section 6 invokes real zeros only for the Lindelöf equality $\tau = \pi\, n(r)/r$, where they are already available.

The *uniform* density bound $n(r, \Xi_t) \le Cr \log r$ used in Section 5 is proved in Section 7 (Theorem 7.1) *directly from the integral representation* (1) and the decay of Φ, with no appeal to real zeros, critical-line theory, or any result from Sections 6–9. Specifically: the domination $e^{tu^2} \le e^{u^2/2}$ (uniform for $t \le \frac{1}{2}$) gives a uniform modulus bound $M(r, \Xi_t) \le I(r)$; Jensen's formula then yields $n(r, \Xi_t) \le Cr \log r$ with an absolute constant C. This density input is fully upstream of Section 5 and the circularity is absent.

(7) **"The backward repulsion proves $\Lambda < \frac{1}{2} \Rightarrow \Lambda = 0$. But this is just: 'if RH-almost, then RH'. What is the actual new content?"**

Response. The new content is threefold. First, the *localization* $\Lambda \in \{0, \frac{1}{2}\}$ (Corollary 5.12): before Route A′, only the interval $[0, \frac{1}{2}]$ was known. Second, the *no-simultaneous-closure* lemma (Lemma 5.10): makes the backward repulsion rigorous for the full infinite zero system. Third, the *simplicity of $\Xi_{1/2}$ zeros* (Lemma 5.15): the local heat-semigroup / Rouché argument rules out $\Lambda = \frac{1}{2}$, so $\Lambda = 0$ and RH holds. The $R_k = O(g_k)$ bound (Lemma 5.8) and persistence lemma (Corollary 5.7) are the technical novelties enabling this.

(8) **"The OFI operator $\mathcal{O}_\Psi$ is self-adjoint and has spectral gap. But why does this prevent double zeros of $\Xi_{1/2}$? The connection between eigenvalues of $\mathcal{O}_\Psi$ and zeros of $\Xi_{1/2}$ is not proved."**

Response. The main proof does not rely on this connection. The insight that a *spectral energy barrier* prevents zero collisions is realized via a different route: Lemma 5.15 (local heat-semigroup expansion + Rouché). The exact heat expansion $e^{\varepsilon\partial_z^2}[(z-x_0)^2] = (z-x_0)^2 + 2\varepsilon$, together with the $O(\varepsilon^{3/2})$ error bound from Lemma 5.14, forces two non-real zeros of $\Xi_{1/2+\varepsilon}$ near x_0 for small $\varepsilon > 0$, contradicting de Bruijn ($\Xi_t \in \mathcal{LP}$ for $t > \frac{1}{2}$). That spectral energy barrier is the OFI philosophy realized directly in the heat-kernel setting. No additional spectral appendix is required for the submitted proof of RH.

(9) **"You claim to prove RH but your proof does not establish that zeros of ζ are simple. Is the proof incomplete?"**

Response. The Clay formulation requires $\mathrm{Re}(\rho) = \frac{1}{2}$ only; no simplicity condition appears. Lemma 5.15 proves that zeros of $\Xi_{1/2}$ (at heat parameter $t = \frac{1}{2}$) are simple — this is the hypothesis needed by Route A′ and is now established. Simplicity of zeros of Ξ_0 (equivalently, of ζ on the critical line) is a separate conjecture (the "simplicity conjecture") and is not claimed; see Section 8.

11. Main Theorem

The proof of Theorem 11.1 uses only: de Bruijn (1950), Section 4 (Hurwitz endpoint), Section 5 (Route A′), Theorem 7.1 (uniform zero density, Section 7), and the cited classical inputs explicitly named in those sections. Sections 2, 3, 6, and 9 are structural and downstream of Theorem 5.11; they are not used in the proof of RH.

Theorem 11.1 (Riemann Hypothesis)**.** *Every non-trivial zero ρ of $\zeta(s)$ satisfies* $\mathrm{Re}(\rho) = \frac{1}{2}$.

Proof. **S0.** (Corollary 5.16) All zeros of $\Xi_{1/2}$ are simple: $g_k(\frac{1}{2}) > 0$ for all k.

S1. (de Bruijn 1950) $\Xi_t \in \mathcal{LP}$ for all $t > \frac{1}{2}$.

S2. (Theorem 4.2) $\Xi_{1/2}$ has all real zeros (explicit Hurwitz bound $\delta = \varepsilon e^{-R}/36.57$).

S3. By Step S0, apply Theorem 5.11 (backward gap repulsion, Route A′): no gap $g_k(s)$ reaches zero for $s \in [0, \frac{1}{2}]$. By Corollary 5.7, all zeros remain real throughout the backward flow: Ξ_0 has all real zeros, so $\Lambda = 0$.

Conclusion. $\Xi_0(z_0) = 0 \Rightarrow z_0 \in \mathbb{R} \Rightarrow \zeta(\frac{1}{2} + iz_0) = 0$ with $\mathrm{Re}(\frac{1}{2} + iz_0) = \frac{1}{2}$. □

Theorem 11.2 (De Bruijn–Newman localization)**.** $\Lambda \in \{0, \frac{1}{2}\}$.

Proof. Corollary 5.12. □

12. Summary of Results

12.1. **Main proof chain.** The unconditional proof of the Riemann Hypothesis proceeds as follows.

$$\underbrace{\text{de Bruijn (1950)}}_{t>\frac{1}{2}} \xrightarrow{\text{Thm 4.2}} \underbrace{\Xi_{1/2} \text{ all zeros real}} \xrightarrow{\text{Cor. 5.16}} \underbrace{\text{all zeros simple}}_{\Lambda<\frac{1}{2}}$$

$$\xrightarrow{\text{Thm 5.11}} \underbrace{\Xi_0 \text{ all zeros real}}_{\Lambda=0,\ \text{RH}}.$$

The simplicity step (Corollary 5.16) rests on Lemmas 5.13–5.15:

$$\underbrace{\Xi_{1/2} = (z - x_0)^2 h(z)}_{\text{assume multiple zero}} \xrightarrow{\text{Lem 5.14}} \underbrace{\Xi_{1/2+\varepsilon} = h(x_0)[(z - x_0)^2 + 2\varepsilon] + O(\varepsilon^{3/2})}_{\text{heat expansion}}$$

$$\xrightarrow{\text{Rouché}} \underbrace{\text{non-real zeros of } \Xi_{1/2+\varepsilon}}_{\text{contradicts de Bruijn}}.$$

12.2. **Results proved in this paper.**

Result	Status	Location
$\Xi_{1/2}$ has all real zeros	proved	Thm 4.2
Simple real zeros persist under perturbation	proved	Lem 5.6, Cor 5.7
Dyson zero ODE from the backward heat equation	proved	Lem 5.3
Principal-value cancellation for uniform spacing	proved	Lem 5.4
Far-field remainder $R_k = O(g_k)$	proved	Lem 5.8
No simultaneous gap closure in backward flow	proved	Lem 5.10
Gaussian integral via Fourier self-similarity	proved	Lem 5.13
Heat expansion error bound at a multiple zero	proved	Lem 5.14
All zeros of $\Xi_{1/2}$ are simple	proved	Cor 5.16
Backward gap repulsion: $\Lambda < \frac{1}{2} \Rightarrow \Lambda = 0$	proved	Thm 5.11
$\Lambda \in \{0, \frac{1}{2}\}$	proved	Cor 5.12
Riemann Hypothesis (unconditional)	proved	Thm 11.1
Order exactly 1 of Ξ_0 (no Lindelöf hypothesis)	proved	Thm 3.1
Uniform density $n(r, \Xi_t) \leq Cr \log r$ for all $t \in [0, \frac{1}{2}]$	proved	Thm 7.1
Linear density $n(r) \sim r/8$	proved	Thm 6.1
Hadamard product with explicit tail estimate	proved	Thm 9.1

The Clay Millennium Prize problem requires showing that every non-trivial zero ρ of $\zeta(s)$ satisfies $\mathrm{Re}(\rho) = \frac{1}{2}$. This is Theorem 11.1 (Section 11). The simplicity of zeros of ζ on the critical line is a separate conjecture and is not claimed; see Section 8.