

Bell Nonlocality from Wigner Negativity in Qudit Stabilizer States

Uta Isabella Meyer, Frédéric Grosshans, Damian Markham

Sorbonne Université, CNRS, LIP6

December 14, 2023



Bell experiments

Measurement scenario for n parties

POVMs $\{(A_{m_i})_k\}_i$ with $\sum_i (A_{m_i})_k = \mathbb{1}_k$, $k = 1, \dots, n$

Experiment

Set of probabilities $\{p(\mathbf{a}|\mathbf{m}) = p(a_i, i = 1, \dots, n | m_i, i = 1, \dots, n)\}$

Quantum Model \rightarrow Born rule

$$p(\mathbf{a}|\mathbf{m}) = \text{Tr}(\Pi_{\mathbf{a}}^{\mathbf{m}} \rho) \quad (1)$$

Classical Model \rightarrow lhv model

$$p(\mathbf{a}|\mathbf{m}) = \sum_{\lambda} \mu(\lambda) \prod_{i=1}^n p_i(a_i | m_i, \lambda) \quad (2)$$

Bell Nonlocality

Bell operator

$$\mathcal{B} = \sum_{m_i} b_{\{m_i\}_i} \bigotimes_{k=1}^n (A_{m_i})_k \quad (3)$$

Bell Inequality

$$\langle \mathcal{B} \rangle_{lhv} = \sum_{m_i} b_{\{m_i\}_i} \sum_{\lambda} \mu(\lambda) \prod_i a_i p_i(a_i | m_i, \lambda) \leq C = \max_{lhv} \langle \mathcal{B} \rangle_{lhv} \quad (4)$$

Bell Violation

$$\text{tr}(\rho \mathcal{B}) = \sum_{m_i} b_{\{m_i\}_i} \left(\prod_i a_i \right) \text{tr} \left(\rho \bigotimes_{k=1}^n \Pi_{a_k}^{m_i} \right) > C \quad (5)$$

Qudits & Gross' Wigner function

Qubits \sim Qu2it	Qudit (d prime)
$X k\rangle = k+1\rangle, Z k\rangle = (-1)^k k\rangle$	$X k\rangle = k+1\rangle, Z k\rangle = \omega^k k\rangle, \omega = e^{2\pi i/d}$
$ZX + XZ = 0$	$ZX - \omega XZ = 0$
$\sigma_{(x,z)} = i^{xz} X^x Z^z,$ $x, z = 0, 1$	$D_{(x,z)} = \omega^{xz/2} X^x Z^z,$ $x, z = 0, \dots, d-1$
$\sigma^{-1} = \sigma = \sigma^\dagger$	$D_{(x,z)}^{-1} = D_{(-x,-z)} = D_{(x,z)}^\dagger$

Wigner function (quasi-probability distribution)

$$W_{(x,z)}(\rho) = \frac{1}{d} \sum_{q,p=0}^{d-1} \omega^{xp-zq} \text{Tr}(\rho D_{(q,p)}) \quad (6)$$

Pauli Stabilizer State & Measurements

Pauli Stabilizer State $|S\rangle$

$S_i|S\rangle = |S\rangle$ with $S_i = \omega^{t(\mathbf{x}_i, \mathbf{z}_i)} D_{(\mathbf{x}_i, \mathbf{z}_i)}$, $\{S_i\}_i$ abelian group.

Measurement description with Pauli operators

Operator Fourier transform

$$A_{(x,z)} = \sum_{q,p=0}^{d-1} \omega^{xp-zq} D_{(p,q)} \quad (7)$$

$\{A_{(x,z)}\}$ are orthonormal basis of operators $(A, B) = \text{tr}(A^\dagger B) / d^n$

Contextuality & Wigner Negativity

Contextuality, a generalization of nonlocality
(Kochen and Specker 1975)

Non-contextual value assignment (Delfosse et al. 2017)

~ Deterministic lhv model

$$D_{(\mathbf{x}, \mathbf{z})} \sim (\mathbf{a}_x, \mathbf{a}_z) \equiv \omega^{\mathbf{x}\mathbf{a}_z - \mathbf{z}\mathbf{a}_x}$$

& probability distributions $\{q\}$ such that

$$\text{tr}(\rho D_{(\mathbf{x}, \mathbf{z})}) = \sum_{\mathbf{a}_x, \mathbf{a}_z} q(\mathbf{a}_x, \mathbf{a}_z) \omega^{\mathbf{x}\mathbf{a}_z - \mathbf{z}\mathbf{a}_x}$$

Nonlocality \Rightarrow Contextuality \Leftrightarrow Wigner negativity (Delfosse et al. 2017)

Pauli Stabilizer States and Clifford operators are non-contextual (Gross 2006)

& efficiently classically simulatable

(Howard, Brennan, and Vala 2013; Howard et al. 2014)

Previous Works & This Work

Nonlocality in Singlet states, GHZ states, graph states

- **d even:** Cerf, Massar, and Pironio 2002, Tang, Yu, and Oh 2013
- **Qutrits:** Lawrence 2017, Chen et al. 2002, Kaszlikowski et al. 2002a; Kaszlikowski et al. 2002b, Acín et al. 2004, Gruca, Laskowski, and Żukowski 2012, Li and Chen 2011, Mackeprang et al. 2023
- Kaniewski et al. 2019, Collins et al. 2001, Ji et al. 2008; Liang, Lim, and Deng 2009

This Work

- Apply Wigner negativity straightaway
- Where to find Wigner negativity?
- No composite systems
- No slicing or subdivising of Qudit into $d' < d$ Qud'it
- Study of d -intrinsic character polynomials

Qudit $\pi/8$ -Gate

Clifford unitary operator (Appleby 2009):

$$C \sim D_{(x,z)} \sum_{j,k=0}^{d-1} \omega^{f_2(j,k)} |j\rangle\langle k| \quad (f_2 \in \text{poly}(\mathbb{Z}_d), \deg(f_2) = 2)$$

Qudit $\pi/8$ -Gate (Howard and Vala 2012)

$$U_\nu = \sum_{k=0}^{d-1} \omega^{\nu_k} |k\rangle\langle k| \quad (8)$$

maps Pauli operators to Clifford operators for $d > 3$ with

$$\nu \in \text{poly}(\mathbb{Z}_d), \deg(\nu) = 3$$

$$\nu_k = 12^{-1}k(\gamma + k(6z + (2k + 3)\gamma)) + \epsilon k \text{ for } z, \epsilon \in \mathbb{Z}_d \text{ and } \gamma \in \mathbb{Z}_d^*$$

Bell Inequalities with Qudit $\pi/8$ gates

Wigner function ($d > 3$)

of $|\Psi_\nu\rangle\langle\Psi_\nu| := (U_\nu \otimes \mathbb{1}) |\Psi\rangle\langle\Psi| (U_\nu^\dagger \otimes \mathbb{1})$, singlet state $|\Psi\rangle = \sum_{k=0}^{d-1} |k k\rangle / \sqrt{d}$:

$$W_{(u_1, u_2)}(|\Psi_\nu\rangle\langle\Psi_\nu|) = \frac{1}{d^3} \delta_{(u_1)_x, (u_2)_x} \sum_{k=0}^{d-1} \omega^{a_3 k^3 + a_1 k}, \quad (9)$$

$$a_3 = 24^{-1}\gamma, \quad a_1 = \epsilon + (u_1)_z + (u_2)_z + z(u_1)_x + 2^{-1}\gamma((u_1)_x^2 - (u_1)_x + 6^{-1})$$

$$\exists \gamma, \epsilon, \text{ s.t. } W_{(u_1, u_2)}(|\Psi_\nu\rangle\langle\Psi_\nu|) < 0$$

$$C(\nu) := W_{(0,0)}(|\Psi_\nu\rangle\langle\Psi_\nu|) < 0,$$

$$|C(\nu)| < 2\sqrt{d} \text{ (Weil's Theorem)}$$

Bell Inequalities with Qudit $\pi/8$ gates

$$D_{(s,r)} \otimes D_{(s,-r)} |\Psi\rangle = |\Psi\rangle : \mathcal{B}_2 = \frac{C(\nu)}{d^3} \sum_{s,r,q=0}^{d-1} U_\nu D_{(s,q-r)} U_\nu^\dagger \otimes D_{(s,r)}$$

$$D_{(\mathbf{s}, \Gamma \mathbf{s})} |G_\Gamma\rangle = |G_\Gamma\rangle : \mathcal{B}_G = \frac{C(\nu)}{d^{n+1}} \sum_{\mathbf{s} \in \mathbb{Z}_d^n, r \in \mathbb{Z}_d} U T_{s_1, (\Gamma \mathbf{s})_1 + r} U^\dagger \bigotimes_{i=2}^n T_{s_i, (\Gamma \mathbf{s})_i}$$

$$S|S\rangle = |S\rangle : \mathcal{B}_S = \frac{C(\nu)}{d^{n+1}} \sum_{\mathbf{u} \in V(S)} \left(\sum_{r=0}^{d-1} \omega^{-(u_1)_x r/2} U S_{u_1} Z_1^r U^\dagger \right) \bigotimes_{i=2}^n S_{u_i}$$

$$\langle \mathcal{B} \rangle_{\text{lhv}^*} = C(\nu) \delta_{(a_x)_1, 0} \delta_{\mathbf{a} \in V(S)} \leq 0, \quad \Rightarrow \quad \langle \mathcal{B} \rangle_{\text{lhv}} \leq 0$$

$$\text{Tr}(\mathcal{B}|S\rangle\langle S|) = C(\nu)^2/d^2 > 0.$$

Bell Inequalities Beyond Characters

$$V_\phi = \sum_{k \in \mathbb{Z}_d} \omega^{k\phi} |k\rangle\langle k|, \quad X_\phi = V_\phi X V_\phi^\dagger, \quad \phi \in \mathbb{Q} \setminus \mathbb{N}$$

Deterministic Bell violation

$$\mathcal{B}_\phi = \frac{1}{d} \sum_{k \in \mathbb{Z}_d} \omega^{-k[\phi]} X^k \otimes X_\phi^k, \quad \langle \Psi | \mathcal{B}_\phi | \Psi \rangle = 1, \quad |\langle \mathcal{B}_\phi \rangle_{\text{lhv}}| < 1$$

$$\phi = 1/2: |\langle \mathcal{B}_\phi \rangle_{\text{lhv}}| \approx 0.647(d=5), 0.642(d=7), 0.639(d=11)$$

Fixed number of measurement operators

$$2\mathcal{B}'_\phi = \omega^{[\phi]} X^\dagger \otimes X_\phi^\dagger + \omega^{[\phi]} X_\phi^\dagger \otimes X^\dagger + X \otimes X + \omega^{-2[\phi]} X_\phi \otimes X_\phi + h.c. \quad .$$
$$\langle \mathcal{B}'_\phi \rangle_{\text{lhv}^*} < 4, \quad \langle \Psi | \mathcal{B}'_\phi | \Psi \rangle = 4$$

$$\phi = 2/3: \langle \mathcal{B}'_\phi \rangle_{\text{lhv}} \approx 3.496(d=5), 3.737(d=7), 3.892(d=11)$$

Discussion

- * Higher degree polynomials $U = \sum_{k=0}^{d-1} \omega^{f_q(k)}$, $q = \deg(f) \leq d - 1$

$$\left\lceil \frac{d-1}{\deg(f)} \right\rceil + 1 \leq |V_f|$$










- * Self-testing mutually unbiased bases
- * Extension to continuous variables

Thank you very much for your attention!

References I

-  Acín, A. et al. (June 2004). “Coincidence Bell inequality for three three-dimensional systems”. In: *Physical Review Letters* 92 (25 I). ISSN: 00319007. DOI: 10.1103/PhysRevLett.92.250404.
-  Appleby, D. M. (2009). *Properties of the extended Clifford group with applications to SIC-POVMs and MUBs*. arXiv: 0909.5233 [quant-ph].
-  Appleby, D Marcus (2005). “Symmetric informationally complete-positive operator valued measures and the extended Clifford group”. In: *Journal of Mathematical Physics* 46.5.
-  Bell, B.A. et al. (2014). “Experimental demonstration of graph-state quantum secret sharing”. In: *Nature communications* 5.1, pp. 1–12. DOI: 10.1038/ncomms6480.
-  Campbell, Earl T., Hussain Anwar, and Dan E. Browne (2012). “Magic-state distillation in all prime dimensions using quantum reed-muller codes”. In: *Physical Review X* 2 (4). ISSN: 21603308. DOI: 10.1103/PhysRevX.2.041021.
-  Carlitz, L. (1979). “Explicit Evaluation of certain Exponential Sums”. In: *Math. Scand.* 44, pp. 5–16.
-  Case, William B. (Oct. 2008). “Wigner functions and Weyl transforms for pedestrians”. In: *American Journal of Physics* 76 (10), pp. 937–946. ISSN: 0002-9505. DOI: 10.1119/1.2957889.
-  Cerf, Nicolas J, Serge Massar, and Stefano Pironio (2002). “Greenberger-Horne-Zeilinger paradoxes for many qudits”. In: *Physical review letters* 89.8, p. 080402.
-  Chen, Jing-Ling et al. (2002). *Clauser-Horne-Bell inequality for three three-dimensional systems*.

References II

-  Chou, Wun Seng, Javier Gomez-Calderon, and Gary L. Mullen (1988). “Value sets of Dickson polynomials over finite fields”. In: *Journal of Number Theory* 30 (3). ISSN: 0022314X. DOI: 10.1016/0022-314X(88)90006-6.
-  Cohen, S. D. (1973). “The values of a polynomial over a finite field”. In: *Glasgow Mathematical Journal* 14 (2), pp. 205–208. ISSN: 1469509X. DOI: 10.1017/S0017089500001981.
-  Collins, Daniel et al. (2001). *Bell inequalities for arbitrarily high dimensional systems*.
-  Delfosse, Nicolas et al. (Dec. 2017). “Equivalence between contextuality and negativity of the Wigner function for qudits”. In: *New Journal of Physics* 19 (12). ISSN: 13672630. DOI: 10.1088/1367-2630/aa8fe3.
-  Durt, T., D. Kaszlikowski, and M. Zukowski (2001). “Violations of local realism with quantum systems described by N-dimensional Hilbert spaces up to [Formula Presented]”. In: *Physical Review A - Atomic, Molecular, and Optical Physics* 64 (2). ISSN: 10941622. DOI: 10.1103/PhysRevA.64.024101.
-  Fitzi, Matthias, Nicolas Gisin, and Ueli Maurer (Nov. 2001). “Quantum solution to the byzantine agreement problem”. In: *Physical Review Letters* 87 (21), pp. 217901–1–217901–4. ISSN: 10797114. DOI: 10.1103/PhysRevLett.87.217901.
-  Gross, D. (2006). “Hudson’s theorem for finite-dimensional quantum systems”. In: *Journal of Mathematical Physics* 47 (12). ISSN: 00222488. DOI: 10.1063/1.2393152.
-  Gruca, Jacek, Wiesław Laskowski, and Marek Żukowski (2012). “Nonclassicality of pure two-qutrit entangled states”. In: *Phys. Rev. A* 85 (2), p. 022118. DOI: 10.1103/PhysRevA.85.022118. URL: <https://link.aps.org/doi/10.1103/PhysRevA.85.022118>.
-  Howard, Mark, Eoin Brennan, and Jiri Vala (2013). “Quantum contextuality with stabilizer states”. In: *Entropy* 15 (6), pp. 2340–2362. ISSN: 10994300. DOI: 10.3390/e15062340.

References III



Howard, Mark and Jiri Vala (Aug. 2012). “Qudit versions of the qubit $\pi/8$ gate”. In: *Physical Review A - Atomic, Molecular, and Optical Physics* 86 (2). ISSN: 10502947. DOI: 10.1103/PhysRevA.86.022316.



Howard, Mark et al. (2014). “Contextuality supplies the ‘magic’ for quantum computation”. In: *Nature* 510 (7505). ISSN: 14764687. DOI: 10.1038/nature13460.



Ji, Se Wan et al. (Nov. 2008). “Multisetting Bell inequality for qudits”. In: *Physical Review A - Atomic, Molecular, and Optical Physics* 78 (5). ISSN: 10502947. DOI: 10.1103/PhysRevA.78.052103.



Kaniewski, Jędrzej et al. (Oct. 2019). “Maximal nonlocality from maximal entanglement and mutually unbiased bases, and self-testing of two-qutrit quantum systems”. In: *Quantum* 3, p. 198. ISSN: 2521-327X. DOI: 10.22331/q-2019-10-24-198. URL: <https://doi.org/10.22331/q-2019-10-24-198>.



Kaszlikowski, Dagomir et al. (2002a). “Clauser-Horne inequality for three-state systems”. In: *Physical Review A - Atomic, Molecular, and Optical Physics* 65 (3). ISSN: 10502947. DOI: 10.1103/PhysRevA.65.032118.










Kaszlikowski, Dagomir et al. (2002b). “Three-qutrit correlations violate local realism more strongly than those of three qubits”. In: *Physical Review A - Atomic, Molecular, and Optical Physics* 66 (3). ISSN: 10502947. DOI: 10.1103/PhysRevA.66.032103.



Keet, Adrian et al. (2010). “Quantum secret sharing with qudit graph states”. In: *Phys. Rev. A* 82 (6), p. 062315. DOI: 10.1103/PhysRevA.82.062315. URL: <https://link.aps.org/doi/10.1103/PhysRevA.82.062315>.

References IV

-  Kochen, Simon and E. P. Specker (1975). “The Problem of Hidden Variables in Quantum Mechanics”. In: *The Logico-Algebraic Approach to Quantum Mechanics: Volume I: Historical Evolution*. Dordrecht: Springer Netherlands, pp. 293–328. ISBN: 978-94-010-1795-4. DOI: 10.1007/978-94-010-1795-4_17. URL: https://doi.org/10.1007/978-94-010-1795-4_17.
-  Lawrence, Jay (2017). “Mermin inequalities for perfect correlations in many-qutrit systems”. In: *Phys. Rev. A* 95 (4), p. 042123. DOI: 10.1103/PhysRevA.95.042123. URL: <https://link.aps.org/doi/10.1103/PhysRevA.95.042123>.
-  Li, Hai Tao and Xiao Yu Chen (2011). “Entanglement of graph qutrit states”. In: *2011 International Conference on Intelligence Science and Information Engineering*, pp. 61–64. ISBN: 9780769544809. DOI: 10.1109/ISIE.2011.10.
-  Liang, Yeong-Cherng, Chu-Wee Lim, and Dong-Ling Deng (2009). “Reexamination of a multisetting Bell inequality for qudits”. In: *Phys. Rev. A* 80 (5), p. 052116. DOI: 10.1103/PhysRevA.80.052116. URL: <https://link.aps.org/doi/10.1103/PhysRevA.80.052116>.
-  Ma, Yuankui and Wenpeng Zhang (Jan. 2019). “On the two-term exponential sums and character sums of polynomials”. In: *Open Mathematics* 17 (1), pp. 1239–1248. ISSN: 23915455. DOI: 10.1515/math-2019-0107.
-  Mackeprang, Jelena et al. (2023). “The power of qutrits for non-adaptive measurement-based quantum computing”. In: *New Journal of Physics* 25.7, p. 073007. DOI: 10.1088/1367-2630/acdf77. URL: <https://dx.doi.org/10.1088/1367-2630/acdf77>.
-  Markham, Damian and Barry C. Sanders (2008). “Graph states for quantum secret sharing”. In: *Phys. Rev. A* 78 (4), p. 042309. DOI: 10.1103/PhysRevA.78.042309. URL: <https://link.aps.org/doi/10.1103/PhysRevA.78.042309>.

References V



Spekkens, Robert W. (2008). “Negativity and contextuality are equivalent notions of nonclassicality”. In: *Physical Review Letters* 101 (2). ISSN: 00319007. DOI: [10.1103/PhysRevLett.101.020401](https://doi.org/10.1103/PhysRevLett.101.020401).



Tang, Weidong, Sixia Yu, and C. H. Oh (2013). “Greenberger-Horne-Zeilinger paradoxes from Qudit graph states”. In: *Physical Review Letters* 110 (10). ISSN: 00319007. DOI: [10.1103/PhysRevLett.110.100403](https://doi.org/10.1103/PhysRevLett.110.100403).



Wan, Da Qing, Peter Jau-Shyong Shiue, and Ching Shyang Chen (1993). “Value sets of polynomials over finite fields”. In: *Proceedings of the American Mathematical Society* 119.3, pp. 711–717.