## bc-no-season: percent difference surface flux of SO2 – NH-sea surface concentration of BC – NH–sea surface concentration of SO4 – NH–sea surface flux of BC – NH–sea surface concentration of SO2 – NH–sea 0.3% 0% 0.2% 1.5% 3% -25% 0.1% Percent Percent Percent Percent Percent 1% -4% 2% 0% -0.1% 1% 0.5% -75% 0% -100% -0.3% 2001 2002 2003 2001 2002 2003 2001 2002 2003 2004 2001 2002 2003 2004 2001 2002 2003 Year Year Year Year Year upwelling longwave flux at TOA – NH-sea upwelling shortwave flux at TOA – NH-sea upwelling clear-sky longwav flux at TOA - NH-sea net radiative flux incident shortwave flux at TOA - NH-sea at TOA - NH-sea 0.2% 0.05% 0.2% 6e-08% 0.025% 0% 0.1% 0.1% 0% -08% Percent Percent Percent -0.01% 0% -0.025% 2e-08% -0.02% -0.05% -0.1% -0.1% 0% -0.03% -0.075% 2001 2003 2001 2003 2001 2003 2001 2002 2003 2004 2003 Year Year Year Year Year dry deposition rate of BC – NH–sea upwelling clear-sky shortway clear-sky net radiative wet deposition rate total deposition rate flux at TOA - NH-sea flux at TOA - NH-sea of BC - NH-sea of BC - NH-sea 5% 0.15% 0.15% 0.1% -5% 0% 0.1% Percent Percent Percent Percent 0.05% 0.05% -10% -6% -5% 0% 0% -15% -8% -0.05% -0.05 2003 2003 2001 2003 2001 2003 2001 2002 2001 2002 2001 2002 2003 Year Year Year Year Year dry deposition rate of SO4 – NH–sea dry deposition rate wet deposition rate wet deposition rate total deposition rate of SO2 - NH-sea of SO2 - NH-sea of SO4 - NH-sea of S - NH-sea 0.4% 0.1% 0% 3% 0% Percent Percent Percent Percent -0.2% 0% -0.2% 0% -0.25% -0.3% 2001 2002 2003 2001 2002 2003 2001 2002 2003 2004 2001 2002 2003 2004 2001 2002 2003 2004 Year Year Year ambient aerosol optical convective cloud cover total cloud cover percentage - NH-sea percentage - NH-sea 0% -20% 0.1% Percent Percent Percent 0.1% -40% 0% -60% -0.1% -80% 2002 2003 2002 2002 2003 2004 2003 2004 2001 2004 2001 2001 Year Year Year

CFSM1

CESM2

F3SM

**GEOS** 

**GFDI** 

**GISS** 

MIROC

NorESM2

OsloCTM3

**UKESM**