

Title

Global monthly sectoral water use for 2010-2100 at 0.5° resolution across alternative futures

Authors

Zarrar Khan¹, Isaac Thompson¹, Chris R. Vernon¹, Neal Graham¹, Thomas B. Wild¹, Min Chen²

Affiliations

1. Joint Global Change Research Institute, Pacific Northwest National Laboratory, 5825 University Research Ct., Suite 3500, College Park, MD, 20740, USA

2. Department of Forest and Wildlife Ecology, College of Agriculture & Life Sciences, University of Wisconsin – Madison, Russell Labs, 1630 Linden Drive, Madison, WI 53706

*corresponding author(s): Zarrar Khan (Zarrar.khan@pnnl.gov)

Abstract

Water usage is closely linked with societal goals that are both local and global in scale, such as sustainable development and economic growth. It is therefore of value, particularly for long-term planning, to understand how future sectoral water usage could evolve on a global scale at fine resolution. At the same time future water usage could be strongly shaped by global forces, such as socioeconomic and climate change, and the multi-sector dynamic interactions those forces create. We generate a novel global gridded monthly sectoral water withdrawal and consumption dataset at 0.5° resolution for 2010-2100 for a diverse range of 75 scenarios. Our scenario repository is harmonized with the five Shared Socioeconomic Pathways (SSPs) and four Representative Concentration Pathways (RCPs) scenarios to support its usage in studies evaluating the implications of uncertain human and earth system change for future global and regional dynamics. To generate the data, we couple the Global Change Analysis Model (GCAM) with a land use spatial downscaling model (Demeter), a global hydrologic framework (Xanthos), and a water withdrawal downscaling model (Tethys).

Background & Summary

This paper documents a global monthly gridded (0.5° resolution) sectoral water withdrawal and consumption dataset that contains conditional projections of water usage (from 2010 to 2100) across a range of future socio-economic and climate scenarios. We generated this dataset by linking together multiple models and datasets designed to explore the dynamic interactions among energy, water, and land systems at global scale and gridded resolution. Central to our modeling workflow is the Global Change Analysis Model (GCAM¹), an integrated tool for exploring the coarse regional dynamics of the coupled human-Earth system and the response of this system to global change, including human system and climate system changes into the future. Tethys² then spatially and temporally downscales outputs from GCAM to grid resolution. We enhance Tethys' projections of irrigation water usage by coupling it with Demeter³, a high-resolution downscaling model that uses GCAM outputs to calculate global gridded land-use change. With the combination of GCAM and Demeter, Tethys is able to project water withdrawal and consumption demands for 6 sectors (domestic, electricity generation, irrigation, livestock, industry and mining) with the irrigation sector further divided into 13 different crop types (biomass, corn, fiber crop, miscellaneous crops, oil crop, other grain, palm fruit, rice, root tuber, sugar crop, wheat, fodder herb, and fodder grass). Withdrawal refers to water that is extracted by a user and then returned to the system, while consumption refers to the part of water withdrawn that is consumed and not returned to the system. To capture a range of futures reflecting diverse global change across the human and Earth systems, we used 75 scenarios comprised of a combination of 4 Representative Concentration Pathways (RCPs)⁴, 5 Shared Socioeconomic Pathways (SSPs)⁵, and 5 Global Climate Models (GCMs) from the Inter-sectoral Impact Model Intercomparison Project (ISIMIP)⁶ protocol 2b. 15 viable combinations of the SSPs and RCPs were combined

with each of the 5 GCMs to arrive at the final 75 scenarios. Graham et al. 2020¹ provides the details on these original GCAM runs for the 75 scenarios. The entire workflow of data from the original scenarios through GCAM and Demeter to Tethys is shown in Figure 1.




Figure 1. Study workflow showing the 75 scenarios are a combination of 4 Representative Concentration Pathways (RCPs), 5 Shared Socioeconomic Pathways (SSPs) and 5 Global Climate Models (GCMs). 15 viable combinations of SSPs and RCPs were combined with each of the 5 GCMs to arrive at the final 75 scenarios which are then used to generate the corresponding GCAM scenarios which are then passed to onto Demeter and Tethys to generate the final results of this study.

This dataset is important because it quantifies the sources of demand-side pressures on scarce water resources globally under diverse future scenarios. Mekonnen & Hoekstra 2016⁷ (also cited in the UN World Water Development Report 2022⁸) estimated that roughly 71% (4.1 billion people) of the world's population was exposed to water scarcity at least one month in the year over the period from 1996 to 2005. In their more recent study, Van Vliet et al. 2021⁹ estimate global water scarcity over the period from 2000 to 2010 to range from 30% (without water quality considered) to 40% (when also including water quality). Global water scarcity is expected to increase across the globe with critical implications for sustainable development^{1,10–13}. Recent studies highlight that future water scarcity is primarily driven by human water demands rather than climate impacts on water availability^{1,14}. Additionally, irrigation water demands have been shown to have the largest relative impact on water scarcity^{10,11,15}. Furthermore, water access, availability and demands are highly localized, with large energy and economic costs associated with water transfers, and thus a regional understanding of water use is essential^{16,17}. This paper accounts for all of these key factors by providing a transparent and open-source dataset and accompanying methodology that captures the key drivers of future water scarcity (water use for human activities) at a fine spatio-temporal scale (0.5° resolution and monthly) and with added detail on irrigation water use by crop types.

Past studies^{18–20} that have evaluated global gridded water use at monthly resolution have been limited to historical analyses. Studies producing future projections²¹ have typically been conducted at coarser resolution, both temporally (i.e., annual time scale) and spatially (i.e., at aggregated country, basin or regional scales). Other studies producing future projections¹⁰ have also used different scenarios and modeling techniques than those we employ here. In

addition to offering a finer spatiotemporal resolution for future projections compared to previous studies, here we provide a broader suite of socioeconomic and climate forcing scenarios, and additional crop water demand resolution through the coupling of water demand model with a land allocation model. Table 1 compares the key features in this study as compared to a representative set of previous studies that have analysed global water use.

This study thus addresses the critical need for future projections of distributed water demand at a fine resolution so that scientists and water managers can start to explore and plan for future water needs. The dataset could also directly support the growing MultiSector dynamics research literature, particularly scenario-based studies of the future interactions between water and other sectors (e.g., energy and land) across scales in a global context^{22,23}. The diverse set of 75 scenarios we produce supports scenario-based water demand uncertainty analysis by varying key elements of human and earth system change. The entire dataset can be easily downloaded from a dataverse online repository (<https://doi.org/10.7910/DVN/VIQEAB>) and is accompanied by a meta-repository (https://jgcri.github.io/khan-etal_2022_tethysSSPRCP/) which provides detailed figures and workflows for interested readers.

Table 1 Comparison of selected global water use studies

| | Water Use Types | Sectors | Additional Sectors | Spatial Scope | Temporal Scope | Scenarios |
|--|--------------------------------|--|--|--|---|---|
| Khan et al. 2022 (This study) | - Withdrawals - Consumption | - Mining - Domestic - Electricity - Livestock - Industry - Irrigation | (13 Crops) Biomass, Corn, Fiber Crop, Misc Crop, Oil Crop, Other Grain, Palm Fruit, Rice, Root Tuber, Sugar Crop, Wheat, Fodder Herb, and Fodder Grass | - Global - 0.5 deg gridded | Historical - 2010 Future/Simulated - 2015 to 2100 - Monthly | Historical 2010 Future - SSPs 1 to 5 - RCP2.6, 4.5, 6.0, 8.5 - 5 CMIP5 GCMs (GFDL, HADGEM, IPSL, MIROC, NORESM) |
| Aqueduct (WRI) (2019, 2015) ^{21,24} | - Withdrawals - Consumption | - Domestic - Industry - Agriculture - Livestock | - | - Global - 0.083 deg (historical) - 0.5 deg (future) | Historical - 1990-2014 - Monthly Future/Simulated: - 2020, 2030, 2040 - Annual | Historical PCR-GLOBWB 2 Outputs Future - SSP2, SSP3 - RCP4.5, RCP8.5 - 6 CMIP5 GCMs (CCSM4, CNRM-CM5, GFDL-ESM2M, INMCM4, MPI-ESM-LR, MRI-CGCM3) |
| Huang et al. 2018 ¹⁸ | - Withdrawals - Consumption | - Mining - Domestic - Electricity - Livestock - Industry - Irrigation | - | - Global - 0.5 deg gridded | Historical - 1971-2010 - Monthly | Historical 4 GHMs: WaterGAP, H08, LPJml, PCR-GLOBWB) |
| Wada et al. 2014 ¹⁹ | - Withdrawals - Consumption | - Domestic - Livestock - Industry - Irrigation | - Paddy - Non-paddy | - Global - 0.5 deg gridded | Historical - 1979 - 2010 - Daily | Historical - 1979-2010 |
| Hanasaki et al. 2013 ¹⁰ | - Withdrawals | - Municipal - Industry - Irrigation | - | - Global - 0.5 deg gridded | Historical - 2000 to 2100 - Daily | Historical 2000 Future - SSPs 1 -5 - RCP2.6, 4.5, 6.0, 8.5 |

| | | | | | | |
|--|--------------------------------------|---------|---|-------------------------------|---|---|
| Mekonnen & Hoekstra 2011 ²⁰ | - Consumption (blue water footprint) | - Total | - Additional datasets available for crops, industrial products and livestock ²⁵⁻²⁷ | - Global - 0.5 deg gridded | <u>Historical</u> - 1996 - 2005 - Monthly | <u>Historical</u> Outputs of water balance model |
|--|--------------------------------------|---------|---|-------------------------------|---|---|

Methods

GCAM produces water withdrawal and consumption outputs for 32 regions for the domestic, mining, power generation, industry, and livestock sectors and for 434 region-basin intersections for the irrigation sector as shown in Figure 2. Tethys v1.3.1²⁸ was used to downscale the water withdrawals and consumption outputs from GCAM onto a 0.5° by 0.5° grid at the equator as shown in Figure 3. Of the 259,200 possible grid cells at this resolution (360 x 720), only the 67,420 cells categorized as land are considered. The Tethys outputs focus only on demand-side dynamics, so they make no distinctions regarding the water supply sources used to meet the demands (i.e., surface water, groundwater, desalinated water), though GCAM does make this distinction.




Figure 2. Water withdrawals and consumption from GCAM by a) 32 GCAM regions for domestic, mining, power generation, industry, and livestock sectors and b) 434 GCAM region and basin intersections for the irrigation sector.




Figure 3. Example outputs of Tethys spatial downscaling of 2010 water withdrawals by sector from GCAM regions and basins to 0.5° x 0.5° grids.

Spatial Downscaling – Non-Agriculture: Spatial downscaling for non-agricultural (domestic, electricity, manufacturing, and mining), water withdrawals and consumption in each grid cell are assumed to be proportional to that cell's population as compared to the larger GCAM region within which that grid cell is located. The population data set used for this paper is from "Gridded Population of the World" (SEDAC, 2016)²⁹. Tethys uses the nearest available year, which for this paper was 2010 in 2010, and 2015 in all other years. Each region's

population is determined by taking the sum of population over all cells belonging to that region. For each of these sectors, Tethys calculates the water withdrawals and consumption as shown in Equation (1) and Equation (2) for a given cell by:

$$\text{withdrawal}_{\text{cell}} = \text{withdrawal}_{\text{region}} \times \frac{\text{population}_{\text{cell}}}{\text{population}_{\text{region}}} \quad (1)$$

$$\text{consumption}_{\text{cell}} = \text{consumption}_{\text{region}} \times \frac{\text{population}_{\text{cell}}}{\text{population}_{\text{region}}} \quad (2)$$

Spatial Downscaling – Livestock: Spatial downscaling of livestock water use is calculated using gridded global maps from the FAO gridded livestock of the world (Wint and Robinson, 2007)³⁰ dataset for six types of livestock (cattle, buffalo, sheep, goats, pigs, and poultry). GCAM outputs are organized into five types (beef, dairy, pork, poultry, and “sheepgoat”), so these are first reorganized to match the six types from Wint and Robinson, 2007³⁰ using ratios for each region estimated from the dataset. The ratios are stored in two files that are used as inputs to Tethys: bfracFAO2005.csv (“buffalo fraction”) and gfracFAO2005.csv (“goat fraction”). The following formulas are used to map the five GCAM livestock types to the six livestock types from Wint and Robinson, 2007³⁰ for each region:

$$\text{buffalo} = (\text{beef} + \text{dairy}) \times \text{buffalo_fraction} \quad (3)$$

$$\text{cattle} = (\text{beef} + \text{dairy}) \times (1 - \text{buffalo_fraction}) \quad (4)$$

$$\text{goat} = (\text{sheepgoat}) \times \text{goat_fraction} \quad (5)$$

$$\text{sheep} = (\text{sheepgoat}) \times (1 - \text{goat_fraction}) \quad (6)$$

No adjustment is required for pork (pigs) or poultry. After this, downscaling for each livestock type is very similar to downscaling the nonagricultural sectors, except the respective livestock population (heads) is used as the proxy instead of human population.

$$\text{withdrawal}_{\text{animal,cell}} = \text{withdrawal}_{\text{animal,region}} \times \frac{\text{heads}_{\text{animal,cell}}}{\text{heads}_{\text{animal,region}}} \quad (7)$$

$$\text{consumption}_{\text{animal,cell}} = \text{consumption}_{\text{animal,region}} \times \frac{\text{heads}_{\text{animal,cell}}}{\text{heads}_{\text{animal,region}}} \quad (8)$$

The results for each of the six types are then added together to get the total livestock withdrawal and consumption for each cell:

$$\text{withdrawal}_{\text{livestock,cell}} = \begin{pmatrix} \text{withdrawal}_{\text{cattle,cell}} + \\ \text{withdrawal}_{\text{buffalo,cell}} + \\ \text{withdrawal}_{\text{sheep,cell}} + \\ \text{withdrawal}_{\text{goat,cell}} + \\ \text{withdrawal}_{\text{pigs,cell}} + \\ \text{withdrawal}_{\text{poultry,cell}} \end{pmatrix} \quad (9)$$

$$\text{consumption}_{\text{livestock,cell}} = \begin{pmatrix} \text{consumption}_{\text{cattle,cell}} + \\ \text{consumption}_{\text{buffalo,cell}} + \\ \text{consumption}_{\text{sheep,cell}} + \\ \text{consumption}_{\text{goat,cell}} + \\ \text{consumption}_{\text{pigs,cell}} + \\ \text{consumption}_{\text{poultry,cell}} \end{pmatrix} \quad (10)$$

Spatial Downscaling – Irrigation: GCAM irrigation water withdrawal and consumption outputs are organized by 13 crop types: Biomass, Corn, Fiber Crop, Miscellaneous Crop, Oil Crop, Other Grain, Palm Fruit, Rice, Root Tuber, Sugar Crop, Wheat, Fodder Herb, and Fodder Grass. By downscaling GCAM output, Demeter³ provides a spatial landcover breakdown by all crop types. In the current version of Tethys (v.1.3.1) biomass is downscaled uniformly within a region-basin intersection (with respect to land area), as given by:

$$\text{withdrawal}_{\text{biomass,cell}} = \text{withdrawal}_{\text{biomass,region}} \times \frac{\text{area}_{\text{cell}}}{\text{area}_{\text{region}}} \quad (11)$$

$$\text{consumption}_{\text{biomass,cell}} = \text{consumption}_{\text{biomass,region}} \times \frac{\text{area}_{\text{cell}}}{\text{area}_{\text{region}}} \quad (12)$$

When possible, the water withdrawal and consumption of the other 12 crops are downscaled in proportion to the crop land area maps from Demeter for each GCAM timestep, which have been reaggregated to the target resolution of 0.5 degrees. There are certain exceptions:

- If the GCAM withdrawal or consumption value for a crop in some region-basin is nonzero, but Demeter does not show any cells with that crop type in that region-basin, it will be downscaled uniformly, as is described above for biomass.
- Additionally, it is possible for GCAM and Demeter to have different total crop irrigation areas for a region-basin intersection, so applying the raw Demeter ratios to irrigation withdrawals or consumption (which are directly related to irrigation areas) could result in cell withdrawal values that imply larger irrigation area than total cell area. In order to avoid this situation, excess irrigation area in cells that are above capacity is assigned evenly among irrigated cells with capacity remaining if there are any, otherwise it is assigned evenly among the remaining cells in the region-basin. Should there still be excess after those cells have been filled, it would be dropped.

Using these adjusted irrigation area values for each crop, cell withdrawal values are given by:

$$\text{withdrawal}_{\text{crop,cell}} = \text{withdrawal}_{\text{biomass,region}} \times \frac{\text{area}_{\text{crop,cell}}}{\text{area}_{\text{crop,region}}} \quad (13)$$

$$\text{consumption}_{\text{crop,cell}} = \text{consumption}_{\text{biomass,region}} \times \frac{\text{area}_{\text{crop,cell}}}{\text{area}_{\text{crop,region}}} \quad (14)$$

The total irrigation sector value for a cell is the sum of that cell's values for all 13 crops.

Temporal Downscaling – Domestic:

Temporally downscaling domestic withdrawal and consumption uses the following formula from Wada et al., 2011³¹. The R parameter described below is from Huang et al. 2018¹⁸ and temperature data is from Weedon et al. 2014³². Withdrawals and consumption for each month of a year for each cell are given by the formula:

$$\text{withdrawal}_{\text{month}} = \frac{\text{withdrawal}_{\text{year}}}{12} \left[\left(\frac{\text{temp}_{\text{month}} - \text{temp}_{\text{min}}}{\text{temp}_{\text{max}} - \text{temp}_{\text{min}}} \right) R + 1 \right] \quad (15)$$

$$\text{consumption}_{\text{month}} = \frac{\text{consumption}_{\text{year}}}{12} \left[\left(\frac{\text{temp}_{\text{month}} - \text{temp}_{\text{min}}}{\text{temp}_{\text{max}} - \text{temp}_{\text{min}}} \right) R + 1 \right] \quad (16)$$

Where:

- temp_{month} = Average temperature for the month
- temp_{mean} = Mean monthly temperature for the year
- temp_{max} = Max monthly temperature for the year
- temp_{min} = Min monthly temperature for the year
- R = Parameter representing the relative difference of water use between the warmest and coolest months of the year

Temporal Downscaling – Electricity Generation:

Water withdrawal and consumption for electricity generation each month are assumed to be proportional to the amount of electricity generated, using the formula developed in Voisin et al., 2013³³:

$$\text{withdrawal}_{\text{month}} = \text{withdrawal}_{\text{year}} \left[\rho_b \left(\frac{\rho_h \frac{\text{HDD}_{\text{month}}}{\text{HDD}_{\text{year}}} + \frac{\rho_c \frac{\text{CDD}_{\text{month}}}{\text{CDD}_{\text{year}}} + \frac{1}{12}}{\rho_u \frac{1}{12}} \right) + \rho_{it} \frac{1}{12} \right] \quad (17)$$

$$\text{consumption}_{\text{month}} = \text{consumption}_{\text{year}} \left[\rho_b \left(\frac{\rho_h \frac{\text{HDD}_{\text{month}}}{\text{HDD}_{\text{year}}} + \frac{\rho_c \frac{\text{CDD}_{\text{month}}}{\text{CDD}_{\text{year}}} + \rho_u \frac{1}{12}}{\rho_c \frac{\text{CDD}_{\text{month}}}{\text{CDD}_{\text{year}}} + \rho_u \frac{1}{12}} \right) + \rho_{it} \frac{1}{12} \right] \quad (18)$$

219

220 Where:

221 ρ_b = Proportion of electricity used for buildings
 222 ρ_{it} = Proportion of electricity used for industry and transportation
 223 $\rho_b + \rho_{it} = 1$
 224 ρ_h = Proportion of electricity used for buildings heating
 225 ρ_c = Proportion of electricity used for buildings cooling
 226 ρ_u = Proportion of electricity used for buildings other
 227 $\rho_h + \rho_c + \rho_u = 1$
 228 HDD = Heating Degree Days
 229 CDD = Cooling Degree Days

230

231 Heating degree days (HDD) and cooling degree days (CDD) are indicators for the
 232 amount of electricity used to heat and cool buildings, and are calculated from mean
 233 daily outdoor air temperature. HDD for a month is the sum of $(18^\circ\text{C} - \text{temperature}_{\text{day}})$
 234 across all days where temperature is less than 18 degrees Celsius. CDD is the sum
 235 of $(\text{temperature}_{\text{day}} - 18^\circ\text{C})$ across all days where temperature is greater than 18.
 236 Annual HDD and CDD are the sum of their respective monthly values.

237

238 Tethys uses HDD, CDD, and pp values for each cell from the nearest available year in
 239 the input files listed at the end of this subsection, which is 2010 for this data set.

240

241 The formula is modified for cells with low annual HDD or CDD as described in Huang et
 242 al., 2018¹⁸, since these may not have heating or cooling services despite nonzero values
 243 of ρ_h or ρ_c .

244

245 When $\text{HDD}_{\text{year}} < 650$, the HDD term is removed and ρ_h is reallocated to the cooling
 246 proportion, giving:

247

$$\text{withdrawal}_{\text{month}} = \text{withdrawal}_{\text{year}} \left[\rho_b \left(\frac{(\rho_h + \rho_c) \frac{\text{CDD}_{\text{month}}}{\text{CDD}_{\text{year}}} + \rho_u \frac{1}{12}}{\rho_c \frac{\text{CDD}_{\text{month}}}{\text{CDD}_{\text{year}}} + \rho_u \frac{1}{12}} \right) + \rho_{it} \frac{1}{12} \right] \quad (19)$$

248

$$\text{consumption}_{\text{month}} = \text{consumption}_{\text{year}} \left[\rho_b \left(\frac{(\rho_h + \rho_c) \frac{\text{CDD}_{\text{month}}}{\text{CDD}_{\text{year}}} + \rho_u \frac{1}{12}}{\rho_c \frac{\text{CDD}_{\text{month}}}{\text{CDD}_{\text{year}}} + \rho_u \frac{1}{12}} \right) + \rho_{it} \frac{1}{12} \right] \quad (20)$$

249

When $CDD_{year} < 450$, the CDD term is removed and ρ_c is reallocated to the cooling proportion, giving:

$$withdrawal_{month} = withdrawal_{year} \left[\rho_b \left(\frac{(\rho_h + \rho_c) \frac{HDD_{month}}{HDD_{year}} + \frac{1}{\rho_u \frac{1}{12}}}{\rho_u \frac{1}{12}} \right) + \rho_{it} \frac{1}{12} \right] \quad (21)$$

$$consumption_{month} = consumption_{year} \left[\rho_b \left(\frac{(\rho_h + \rho_c) \frac{HDD_{month}}{HDD_{year}} + \frac{1}{\rho_u \frac{1}{12}}}{\rho_u \frac{1}{12}} \right) + \rho_{it} \frac{1}{12} \right] \quad (22)$$

When annual HDD and CDD are both below their respective thresholds, all sources of monthly variation vanish and the formula reduces to

$$withdrawal_{month} = \frac{withdrawal_{year}}{12} \quad (23)$$

$$consumption_{month} = \frac{consumption_{year}}{12} \quad (24)$$

Temporal Downscaling – Livestock, Manufacturing and Mining:

For livestock, manufacturing, and mining, a uniform distribution is applied. The withdrawal or consumption for the year is divided between months according to the number of days.

$$withdrawal_{month} = withdrawal_{year} \times \frac{days_{month}}{days_{year}} \quad (25)$$

$$consumption_{month} = consumption_{year} \times \frac{days_{month}}{days_{year}} \quad (26)$$

Temporal Downscaling – Irrigation:

Temporal downscaling for irrigation water withdrawal and consumption is based on weighted irrigation profiles for each of the 235 basins. Gridded monthly irrigation withdrawal values from the PCR-GLOBWB global hydrological (from Huang et al. 2018¹⁸, original data from ISIMIP³⁴) model are averaged across the years 1971-2010, then aggregated to the basin scale. The monthly irrigation withdrawal percentages for a basin are applied to all crops in each of its cells:

$$withdrawal_{month} = withdrawal_{year} \times percent_{basin,month} \quad (27)$$

$$consumption_{month} = consumption_{year} \times percent_{basin,month} \quad (28)$$

In the event that the model has no monthly data for a basin with nonzero irrigation, the profile of the nearest available basin is used.

Data Records

Data outputs from this experiment have been minted and are available in the repository indicated in Table 2. A meta-repository with detailed information on the workflows to produce the data is also available and shown in Table 2.

Table 2 Data records

| Record | Details | Location |
|----------------------------|--|---|
| Output Dataset | Data outputs from experiment | https://doi.org/10.7910/DVN/VIQEAB |
| Supporting Meta-repository | Meta-repository with detailed workflows for experiment | https://jgcri.github.io/khan-etal_2022_tethysSSPRCP/index.html |

The dataset contains separate files with names which start with a combination of the following SSP, RCP, GCM and water usage type:

- **SSP:** ssp1, ssp2, spp3, spp4, spp5
- **RCP:** rcp26, rcp45, rcp60, rcp85
- **GCM:** gfdl, hadgem, ipsl, miroc, noresm
- **Water use type:** consumption, withdrawals

Example 1: ssp1_rcp26_gfdl_consumption_XXX

Example 2: ssp1_rcp26_gfdl_withdrawal_XXX

The datasets files have been then divided into sub-sets to manage their size. The following list shows the file structure for one of the SSP, RCP, GCM combinations:

- ssp1_rcp26_gfdl_consumption_crops_annual.zip
- ssp1_rcp26_gfdl_consumption_crops_monthly_1.zip
- ssp1_rcp26_gfdl_consumption_crops_monthly_2.zip
- ssp1_rcp26_gfdl_consumption_sectors_annual.zip
- ssp1_rcp26_gfdl_consumption_sectors_monthly_1.zip
- ssp1_rcp26_gfdl_consumption_sectors_monthly_2.zip

The files with "_crops_" in their names include data for individual crops while the files with "_sectors_" in their name include data for other aggregated sectors. The following expanded list shows the individual files inside the zipped files for the example ssp1_rcp26_gfdl cases. **"cd" stands for "consumption downscaled"** and **"tcd" stands for "temporal consumption downscaled"**:

- ssp1_rcp26_gfdl_consumption_crops_annual.zip
 - crops_cdirr_biomass_km3peryr.csv
 - crops_cdirr_Corn_km3peryr.csv
 - crops_cdirr_FiberCrop_km3peryr.csv
 - crops_cdirr_FodderGrass_km3peryr.csv
 - crops_cdirr_FodderHerb_km3peryr.csv

- crops_cdirr_MiscCrop_km3peryr.csv
- crops_cdirr_OilCrop_km3peryr.csv
- crops_cdirr_OtherGrain_km3peryr.csv
- crops_cdirr_PalmFruit_km3peryr.csv
- crops_cdirr_Rice_km3peryr.csv
- crops_cdirr_Root_Tuber_km3peryr.csv
- crops_cdirr_SugarCrop_km3peryr.csv
- crops_cdirr_Wheat_km3peryr.csv
- ssp1_rcp26_gfdl_consumption_crops_monthly_1.zip
 - crops_tcdirr_biomass_km3peryr.csv
 - crops_tcdirr_Corn_km3peryr.csv
 - crops_tcdirr_FiberCrop_km3peryr.csv
 - crops_tcdirr_FodderGrass_km3peryr.csv
 - crops_tcdirr_FodderHerb_km3peryr.csv
 - crops_tcdirr_MiscCrop_km3peryr.csv
 - crops_tcdirr_OilCrop_km3peryr.csv
- ssp1_rcp26_gfdl_consumption_crops_monthly_2.zip
 - crops_tcdirr_OtherGrain_km3peryr.csv
 - crops_tcdirr_PalmFruit_km3peryr.csv
 - crops_tcdirr_Rice_km3peryr.csv
 - crops_tcdirr_Root_Tuber_km3peryr.csv
 - crops_tcdirr_SugarCrop_km3peryr.csv
 - crops_tcdirr_Wheat_km3peryr.csv
- ssp1_rcp26_gfdl_consumption_sectors_annual.zip
 - cddom_km3peryr.csv(Domestic)
 - cdelec_km3peryr.csv(Electricity Generation)
 - cdirr_km3peryr.csv(Irrigation)
 - cdliv_km3peryr.csv(Livestock)
 - cdmfg_km3peryr.csv(Industry & manufacturing)
 - cdmin_km3peryr.csv(Mining)
 - cdnonag_km3peryr.csv(Aggregated non-agriculture)
 - cdtotal_km3peryr.csv(Total)
- ssp1_rcp26_gfdl_consumption_sectors_monthly_1.zip
 - tcddom_km3peryr.csv(Domestic)
 - tcdelec_km3peryr.csv(Electricity Generation)
 - tcdirr_km3peryr.csv(Irrigation)
- ssp1_rcp26_gfdl_consumption_sectors_monthly_2.zip
 - tcdliv_km3peryr.csv(Livestock)
 - tcdmfg_km3peryr.csv(Industry & manufacturing)
 - tcadmin_km3peryr.csv(Mining)

Technical Validation

GCAM outputs are calibrated at a regional scale to matched observed data for base year values as described in Graham et al. 2020. In this study we the validation is limited to ensuring that the downscaling algorithms in Tethys are free of errors and there is no loss in values as a result of the temporal or spatial downscaling methodology. Results of the model were validated by re-aggregating spatial and temporal downscaled model outputs and comparing them to the original aggregated inputs. Figure 4a shows how the disaggregated water withdrawal values in km³ equal the original values both spatially for GCAM regions and temporally for annual values across sectors and crops. Figure 4b shows the same validation for how the disaggregated water consumption values in km³ equal back to the original values both spatially for GCAM regions and temporally for annual values across sectors and crops.





Figure 4 Validation of downscaled spatial and temporal Tethys water use. a) Water Withdrawals (km³) and b) Water Consumption (km³)

375 Additionally, Tethys outputs were also compared to results from two other studies: Huang et
376 al. 2018¹⁸ and Mekonnen, M.M. and Hoekstra, A.Y. 2011²⁰ as shown in Figure 5. Given the
377 larger number of variables and assumptions for future scenarios considered here, we limit
378 the validation with other studies to historical data. Since this work is primarily concerned
379 with the downscaling of existing projections to a gridded monthly scale, we look at how
380 spatial and temporal patterns in the year 2010 (for which all scenarios are identical) compare
381 to those of the chosen datasets.

382
383 Huang et al. 2018¹⁸, uses an earlier version of Tethys on historical data from 1971-2010. The
384 underlying data have more regions and different totals, but many of the downscaling
385 methods are identical, leading to similar results. For the non-agricultural sectors (domestic,
386 electricity, manufacturing, and mining), the same underlying population map is used to
387 downscale water use. For irrigation, Huang et al. 2018¹⁸ use USGS and FAO AQUASTAT
388 irrigation data, whereas the current version of Tethys uses crop landcover maps from
389 Demeter. Consumption and withdrawals generally showed similar spatial patterns, with
390 differences in assumptions regarding each region's and sector's consumption-to-withdrawal
391 ratios accounting for some differences. There are also some differences in accounting. For
392 example, in this study hydropower is included in the consumption for electricity generation
393 category, which by itself is several times greater than the entire water consumption for
394 electricity generation in Huang et al. 2018¹⁸.

395
396 The second data set we compared with is from Mekonnen, M.M. and Hoekstra, A.Y. 2011²⁰.
397 It contains monthly total blue water consumption values representing an average of years
398 1996-2005, which we compare to the base year values from 2010 from this study. The
399 sectoral breakdown is different between the two datasets, but the datasets are at the same
400 spatial resolution, so we compare monthly totals for each grid cell. We see some agreement
401 between the two data sets. As the largest sector, differences in irrigation downscaling are
402 likely responsible for the variation.

403
404 As seen in Figure 5 we see a general agreement in the sub-regional patterns across the data
405 sets. Figure 6 also shows similar sub-annual patterns across the dataset with some
406 differences in total values being attributed to underlying data and year of the study.
407 Additional details on differences between the datasets are discussed in the meta-repository
408 https://igcri.github.io/khan-etal_2022_tethysSSPRCP/index.html.




Figure 5 Spatial distribution of water withdrawals and consumption across this study, Huang et al. 2018¹⁸ and Mekonnen, M.M. and Hoekstra, A.Y. 2011²⁰




Figure 6 Temporal distribution of water withdrawals and consumption across this study, Huang et al. 2018¹⁸ and Mekonnen, M.M. and Hoekstra, A.Y. 2011²⁰

Usage Notes

Users are encouraged to explore the accompanying meta-repository (https://jgcri.github.io/khan-etal_2022_tethysSSPRCP/index.html), which provides detailed visualization across the various scenarios, sectors and time periods. Users can then download specific datasets for water withdrawal or consumption for relevant sectors, crops and desired SSP, RCP or GCM from the accompanying dataset repository (<https://doi.org/10.7910/DVN/VIQEAB>) to analyze the raw data. Some example figures from the meta-repository are presented in this section.

Figure 7a shows the total annual water withdrawals by sector for each of the 75 SSP-RCP-GCM combinations from 2010 to 2100. Similar figures are available for consumption as well as by crop. Figure 7b shows the sub-annual temporal distribution across the same set of scenarios for 2010 and for 2100. Patterns such as an increase in summer water withdrawals can be seen in such figures.

The meta-repository also includes details on three selected basins: the Indus, Nile and Upper Colorado River Basin (U.S.). These are used to show how the data can be used to explore trends and patterns at this finer resolution. Figure 8a and Figure 8b are examples showing how land-use change impacts which type of crop becomes the dominant water user in the Indus basin over time for the SSP1-RCP2.6-GFDL scenario. Figure 8c and Figure 8d show the accompanying distribution of total water withdrawals both spatially and temporally. Similar figures are provided in the meta-repository for water consumption as well as for other sectors, crops and scenarios.






Figure 7 Global water withdrawals for the 75 SSP-RCP-GCM combinations by sector. a) Annual water withdrawals by sector from 2010 to 2100. b) Monthly water withdrawals for 2010 and 2100


a) Indus basin: ssp1_rcp26_gfdl maximum crop water withdrawals



b) Indus basin: ssp1_rcp26_gfdl annual crop water withdrawals



c) Indus basin: ssp1_rcp26_gfdl total water withdrawals by grid cell



d) Indus basin: ssp1_rcp26_gfdl total water withdrawals by month and year




Figure 8 Indus Basin water withdrawals (km³) by crop for scenario SSP 1, RCP 2.6, GCM GFDL. a) Showing which crop has the maximum water withdrawals (km³) in each grid cell for years 2025, 2050, 2075 and 2100. b) Aggregated water withdrawals (km³) by crop in the Indus Basin from 2015 to 2100. c) Showing total water withdrawals (km³) in each grid cell for years 2025, 2050, 2075 and 2100. d) Aggregated total water withdrawals (km³) in the Indus Basin from 2015 to 2100.

We highlight that several developments have been planned in the next release of Tethys to improve the methodologies used to downscale water use for the dataset in this paper. Some of the key planned developments include:

1. Improving the spatial distribution of powerplant water use based on actual and projected powerplant location instead of based on population.
2. Updating the output resolution to 1/8th degrees from the existing 1/2 degree resolution.

3. Including future population projections to improve on the current methodology which uses a static base year population map even for future years.
4. Improving the downscaling of biomass water use which is currently distributed equally within each region.
5. Making Tethys compatible with GCAM-USA³⁵, which allow use of more accurate state-level water use data instead of using national data as inputs to Tethys.

Code Availability

The following table provides links to all models, data, versions and DOI's used to generate this dataset.

| Type | Details | Model Version | Data DOI | Model DOI |
|----------------|---|---------------|---|---|
| Tethys | Used to generate the data presented in this paper | v1.3.0 | https://doi.org/10.7910/DVN/VIQEAB | https://doi.org/10.5281/zenodo.6399488 |
| GCAM | Water use data used as inputs for Tethys | v4.3.chen | https://data.pnnl.gov/dataset/13224 | http://doi.org/10.5281/zenodo.3713432 |
| Demeter | Landuse change data used as input for Tethys | v1.chen | https://data.pnnl.gov/dataset/13192 | http://doi.org/10.5281/zenodo.3713378 |

Acknowledgements

This research was supported by the U.S. Department of Energy, Office of Science, as part of research in MultiSector Dynamics, Earth and Environmental System Modeling Program. The Pacific Northwest National Laboratory is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RL01830. The views and opinions expressed in this paper are those of the authors alone.

Author contributions

Z.K., I.T., P.P., C.R.V., N.G., T.W., and M.C., designed the research.
 Z.K. and I.T. ran Tethys to produce the outputs, prepared the figures and the data repository.
 N.G. produced the GCAM data used as inputs for Tethys.
 M.C. produced the Demeter data used as inputs for Tethys.
 Z.K., I.T., P.P., C.R.V., N.G., T.W. all contributed to writing and reviewing the paper.

Competing interests

The authors declare no competing interests.

References

1. Graham, N. T. *et al.* Humans drive future water scarcity changes across all Shared Socioeconomic Pathways. *Environ. Res. Lett.* **15**, 014007 (2020).
2. Li, X. *et al.* Tethys – A Python Package for Spatial and Temporal Downscaling of Global Water Withdrawals. *Journal of Open Research Software* **6**, (2018).
3. Vernon, C. R. *et al.* Demeter – A Land Use and Land Cover Change Disaggregation Model. *Journal of Open Research Software* **6**, 15 (2018).
4. van Vuuren, D. P. *et al.* The representative concentration pathways: an overview. *Climatic Change* **109**, 5 (2011).
5. O'Neill, B. C. *et al.* The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century. *Global Environmental Change* **42**, 169–180 (2017).
6. ISIMIP. Inter Sectoral Impact Model Intercomparison (ISIMIP) - Input Data and Bias Correction. (2019).
7. Mekonnen, M. M. & Hoekstra, A. Y. Four billion people facing severe water scarcity. *Science Advances* **2**, e1500323 (2016).
8. UNESCO. *The United Nations World Water Development Report 2022: Groundwater: Making the invisible visible*. <https://unesdoc.unesco.org/ark:/48223/pf0000380721> (2022).
9. Vliet, M. T. H. van *et al.* Global water scarcity including surface water quality and expansions of clean water technologies. *Environ. Res. Lett.* **16**, 024020 (2021).
10. Hanasaki, N. *et al.* A global water scarcity assessment under Shared Socio-economic Pathways – Part 1: Water use. *Hydrol. Earth Syst. Sci.* **17**, 2375–2391 (2013).
11. Hanasaki, N. *et al.* A global water scarcity assessment under Shared Socio-economic Pathways – Part 2: Water availability and scarcity. *Hydrol. Earth Syst. Sci.* **17**, 2393–2413 (2013).

- 520 12. Hejazi, M. I. *et al.* Integrated assessment of global water scarcity over the 21st
521 century under multiple climate change mitigation policies. *Hydrology and Earth System*
522 *Sciences* **18**, 2859–2883 (2014).
- 523 13. Wada, Y. & Bierkens, M. F. P. Sustainability of global water use: past reconstruction
524 and future projections. *Environ. Res. Lett.* **9**, 104003 (2014).
- 525 14. Wada, Y., Beek, L. P. H. van, Wanders, N. & Bierkens, M. F. P. Human water
526 consumption intensifies hydrological drought worldwide. *Environ. Res. Lett.* **8**, 034036
527 (2013).
- 528 15. Yoshikawa, S. *et al.* An assessment of global net irrigation water requirements from
529 various water supply sources to sustain irrigation: rivers and reservoirs (1960–2000 and
530 2050). <https://hess.copernicus.org/preprints/10/1251/2013/hessd-10-1251-2013.pdf>
531 (2013) doi:10.5194/hessd-10-1251-2013.
- 532 16. Veldkamp, T. I. E. Water scarcity at the global and regional scales: unravelling its
533 dominant drivers in historical and future time periods. (2017).
- 534 17. Wada, Y., de Graaf, I. E. M. & van Beek, L. P. H. High-resolution modeling of human
535 and climate impacts on global water resources. *Journal of Advances in Modeling Earth*
536 *Systems* **8**, 735–763 (2016).
- 537 18. Huang, Z. *et al.* Reconstruction of global gridded monthly sectoral water withdrawals
538 for 1971–2010 and analysis of their spatiotemporal patterns. *Hydrology and Earth System*
539 *Sciences* **22**, 2117–2133 (2018).
- 540 19. Wada, Y., Wisser, D. & Bierkens, M. F. P. Global modeling of withdrawal, allocation
541 and consumptive use of surface water and groundwater resources. *Earth System*
542 *Dynamics* **5**, 15–40 (2014).
- 543 20. Mekonnen, M. M. & Hoekstra, A. Y. *Total monthly blue water footprints of*
544 *production at a 30 × 30 arc minute grid resolution (1996-2005).*

- 545 <https://waterfootprint.org/en/resources/waterstat/monthly-gridded-blue-water->
546 [footprint-statistics/](https://waterfootprint.org/en/resources/waterstat/monthly-gridded-blue-water-footprint-statistics/) (2011).
- 547 21. Hofste, R. W. *et al.* Aqueduct 3.0: Updated decision-relevant global water risk
548 indicators. *World Resources Institute: Washington, DC, USA* (2019).
- 549 22. Wild, T. B. *et al.* The Implications of Global Change for the Co-Evolution of
550 Argentina's Integrated Energy-Water-Land Systems. *Earth's Future* **9**, e2020EF001970
551 (2021).
- 552 23. Reed, P. M. *et al.* Multisector Dynamics: Advancing the Science of Complex Adaptive
553 Human-Earth Systems. *Earth's Future* **10**, e2021EF002621 (2022).
- 554 24. WRI. WRI Aqueduct. (2021).
- 555 25. Mekonnen, M. & Hoekstra, A. National water footprint accounts: The green, blue
556 and grey water footprint of production and consumption. Volume 1: Main Report.
557 *Daugherty Water for Food Global Institute: Faculty Publications* (2011).
- 558 26. Mekonnen, M. M. & Hoekstra, A. Y. The green, blue and grey water footprint of
559 crops and derived crops products. (2010).
- 560 27. Mekonnen, M. M. & Hoekstra, A. Y. A Global Assessment of the Water Footprint of
561 Farm Animal Products. *Ecosystems* **15**, 401–415 (2012).
- 562 28. Khan, Z., Thompson, I. & Vernon, C. R. *Tethys v1.3.1*. (2022).
- 563 29. SEDAC. Gridded Population of the World, Version 4 (GPWv4): Population Density,
564 Revision 11. *Center for International Earth Science Information Network - CIESIN -*
565 *Columbia University. 2018. Palisades, NY: NASA Socioeconomic Data and Applications*
566 *Center (SEDAC)* <https://doi.org/10.7927/H49C6VHW> (2018).
- 567 30. Wint, W. & Robinson, T. *Gridded livestock of the world 2007*. (FAO, Roma (Italia),
568 2007).
- 569 31. Wada, Y. *et al.* Global monthly water stress: 2. Water demand and severity of water
570 stress. *Water Resources Research* **47**, (2011).

- 571 32. Weedon, G. P. *et al.* The WFDEI meteorological forcing data set: WATCH Forcing Data
572 methodology applied to ERA-Interim reanalysis data. *Water Resources Research* **50**, 7505–
573 7514 (2014).
- 574 33. Voisin, N. *et al.* One-way coupling of an integrated assessment model and a water
575 resources model: evaluation and implications of future changes over the US Midwest.
576 *Hydrology and Earth System Sciences* **17**, 4555–4575 (2013).
- 577 34. Warszawski, L. *et al.* The Inter-Sectoral Impact Model Intercomparison Project (ISI–
578 MIP): Project framework. *PNAS* **111**, 3228–3232 (2014).
- 579 35. Binsted, M. *et al.* GCAM-USA v5.3_water_dispatch: integrated modeling of
580 subnational US energy, water, and land systems within a global framework. *Geoscientific*
581 *Model Development* **15**, 2533–2559 (2022).
- 582
- 583