

Algebra I

September 28, 2023

Grade Weights

50% Homework + 50% Final

Participation matters for pass/fail.

Office Hours

Tuesday / Thursday 11:25 - 12:00

Or by appointment (jusuh@ucsc.edu)

Recommended Text

Abstract Algebra (3e) - Dummit and Foote

Finite Groups: An Introduction (2nd revised) - Jean-Pierre Serre

Robert Boltje's Lecture Notes - (<https://boltje.math.ucsc.edu/courses/f17/f17m200notes.pdf>)

Definition: Binary Operation

Let S be a set. A binary operation on S is a function $f : S \times S \rightarrow S$. We will almost never use f for the binary operation ($f(s, t)$).

The usual notation for binary operations is $s * t$.

Example

1. $S = \mathbb{R}^3$, define $f : S \times S \rightarrow S$ as $(\vec{x}, \vec{y}) \rightsquigarrow \vec{x} + \vec{y}$.

2. $S = \mathbb{R}^3$, define $S \times S \xrightarrow{f} S$ as $(\vec{x}, \vec{y}) \rightsquigarrow \vec{x} + \vec{y}$.

- Note that $(\vec{x}, \vec{y}) \rightsquigarrow \vec{x} \cdot \vec{y}$ is not a binary operation.

3. $S = \mathbb{Z}$ as $(m, n) \mapsto m \cdot n$.

4. $S = \mathbb{R}^3$ as $(\vec{x}, \vec{y}) \rightsquigarrow \frac{\vec{x} + \vec{y}}{2}$



5. Let $n \geq 1$ be an integer and $S = M_{n \times n}(\mathbb{R}) = \{n \times n \text{ real matrices}\}$. Then $(A, B) \rightsquigarrow AB$.

Observations

Examples 1,3,5 are associative; examples 2,4 are not.

Examples 1-4 are commutative; example 5 commutes only when $n = 1$.

$\vec{0}$ for example 1, 1 for example 3, and I_n for example 5.

Q: What is a Group?

A group is a set equipped with a binary operation which satisfies three axioms.

Let $*$ be a binary operation on a set S .

1. Say $*$ is associative if $\forall a, b, c \in S, (a * b) * c = a * (b * c)$.
2. Say $*$ is commutative if $\forall a, b \in S, a * b = b * a$.
3. An element $e \in S$ is a neutral element (with respect to $*$) if $\forall a \in S, a * e = a = e * a$.
 - If there exists a neutral element, then it is unique.
4. Suppose $(S, *)$ has a neutral element e . Let $a \in S$. Then $b \in S$ is called an inverse of a (with respect to $*$) if $a * b = e = b * a$.

Definition: Group

A group is a set G equipped with a binary operation $*$ such that

1. $*$ is associative.
2. $*$ has a neutral element e .
3. Every $g \in G$ has an inverse.

If, in addition, $*$ is commutative, we say $(G, *)$ is an abelian or commutative group.

Examples

$(\mathbb{R}^3, +)$ is a commutative group.

(\mathbb{R}^3, \times) has no neutral element.

(\mathbb{Z}, \cdot) has no inverse (except ± 1).

$(\mathbb{R}^3, \text{mid})$ is not associative. (the midpoint)

$(M_{n \times n}(\mathbb{R}), \cdot)$ has no inverse of $0_{n \times n}$.

For $n \geq 1$, $(\mathbb{R}^n, +)$ and $(\mathbb{C}^n, +)$ are abelian groups.

Proof that the Neutral Element is unique.

Let e, e' be neutral elements. Then $e' = e * e' = e$. ■

Proof that the Inverse is unique.

Left to the reader.

Definition: Subgroup

Let G be a group, and let H be a subset of G . We say that H is a subgroup of G if

1. $\forall h_1, h_2 \in H, h_1 * h_2 \in H$.
2. $e \in H$.
3. $\forall h \in H, h^{-1} \in H$.

Examples

$\mathbb{Z}^n \subseteq \mathbb{R}^n$ is a subgroup ($*$ = +).

$G = \{A \in M_{n \times n} : \det(A) \neq 0\}$. Then (G, \cdot) is a group.

- This is the General Linear Group on \mathbb{R} : $\text{GL}_n(\mathbb{R})$.
- Recall $A^{-1} = \frac{1}{\det(A)} \left((-1)^{ij} \det(M_{\alpha_i}) \right)$.

Definition: General Linear Subgroups

$S = \{A \in \text{GL}_n(\mathbb{R}) : a_{ij} \in \mathbb{Z}, \forall 1 \leq i, j \leq n\}$.

S is closed under \cdot and $I_n \in S$, but for example

$$\begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}^{-1} = \frac{1}{1 \cdot 4 - 2 \cdot 3} \begin{pmatrix} 4 & -2 \\ -3 & 1 \end{pmatrix} = \begin{pmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{pmatrix}$$

so S is not a subgroup.

However, $T = \{A \in S : \det(A) = \pm 1\} \subseteq \text{GL}_n(\mathbb{R})$.

- Note that if $AA' = I_n$ then $\det(A)\det(A') = 1$.

Definition: Additive Groups

For groups like \mathbb{Z}^n , \mathbb{R}^n and \mathbb{C}^n , we will use $+$ for the binary operation and say that they are additive groups. The Neutral Element is denoted as 0.

The inverse is denoted as $-g$.

For $m \geq 1$ and $g \in G$, $mg = g + \dots + g$ and $(-m)g = -(mg)$.

Definition: Multiplicative Groups

For groups like $\text{GL}_n(\mathbb{C})$ or $\text{GL}_n(\mathbb{Z})$, we say that the group is multiplicative.

Denote the neutral element as 1.

Denote the inverse of g as g^{-1} .

For $m \geq 1$, $g^m = \underbrace{g \cdots g}_m$.

$g^0 = 1$.

$g^{-m} = (g^m)^{-1}$.

Definition: Group Element Order

Let G be a group, $g \in G$, and $m \geq 1$.

Say g has order m if $g^m = 1$ and $g^k \neq 1$, $\forall k$ such that $1 \leq k \leq m$.

An element has infinite order if $g^m \neq 1$, $\forall m \in \mathbb{Z}^+$.

Examples

In D_{10} , I_2 has order 1, rotations have order 5 and reflections have order 2.

Groups from Geometry

Pentagon

Consider the regular pentagon P .



$$H = \{T \in \text{GL}_2(\mathbb{R}) : T(P) = P\}.$$

This is the symmetry group of P or D_{10} (sometimes D_5)

$$H \leq \text{GL}_2(\mathbb{R}).$$

- Proof of closure.

Suppose $T_1, T_2 \in H$. Then $T_1(P) = P$, $T_2(P) = P$ and $(T_1 \circ T_2)(P) = T_1(T_2(P)) = T_1(P) = P$.

Therefore H is closed under \circ .

- Proof of identity.

$\text{Id}_{\text{GL}_2} = I_2$ does satisfy $I_2(P)$.

- Proof of inverse.

If $T \in H$ (i.e. $T \in \text{GL}_2(\mathbb{R})$ and $T(P) = P$, apply T^{-1} and get $T^{-1}(T(P)) = T^{-1}(P)$. Therefore $P = T^{-1}(P)$.

Tetrahedron

Let X be the regular tetrahedron and $A = \{\text{rotational symmetries of } X\}$.



Then A contains

- The identity: 1.
- $2 \cdot 4 = 8$ rotations by 120° .
- 3 rotations of 180° .

So we have a bijection $r : \{B, P, W, Y\} \rightarrow \{B, P, W, Y\}$ where



Definition: Symmetric Group

Let S be a set (e.g. $E = \{B, P, W, Y\}$). The Symmetric Group $\text{Sym}(E)$ is the set of bijections $f : E \rightarrow E$ equipped with the binary operation \circ (composition).

October 3, 2023

Homework

First homework should be released this Thursday, October 5th.

Next lecture will be on group actions.

Propositions: Symmetric Group

Let X be a set.

When $|X| = n$ denote the elements $\{1, 2, \dots, n\}$.

$\text{Sym}(X) = \{f : X \rightarrow X \mid f \text{ is bijective}\}$.

With \circ (composition of functions) as a binary operation, $\text{Sym}(X)$ is a group.

Symmetric Group Order

If $|X| = n$, then $|\text{Sym}(X)| = n!$

- Proof

Let $X = \{1, 2, \dots, n\}$. A bijection f consists of $f(1), f(2), \dots, f(n)$.

For $f(1)$, we have n choices; for $f(2)$ we have $n - 1$ choices. This continues until only 1 choice remains for $f(n)$.

Therefore the choices are $(n)(n - 1) \cdots (1) = n!$

Example

For the symmetric group on four letters $\{a, b, c, d\}$, $|\text{Sym}(4)| = 4! = 24$

Definition: Cycles

Let $x = \{1, \dots, n\}$, $m \geq 1$ be an integer and a_1, a_2, \dots, a_m distinct elements in X .

Then the m -cycle denoted by $(a_1 a_2 \dots a_m)$ is the element of $\text{Sym}(X)$ which maps a_1 to a_2 , a_2 to a_3, \dots, a_{m-1} to a_m , and a_m to a_1 .

Example

Let $n = 7$ and $m = 4$. Then $(2 \ 7 \ 1 \ 3)$ is a bijection.



Degenerate Case

$m = 1$ gives Id_X .

First Non-Degenerate Case

A transposition is, by definition a 2-cycle: $(a_1 a_2)$.

Proposition: Symmetric Group as Cycle Composition

Every element in $\text{Sym}(X)$ is the product (using \circ) of m -cycles, where m can vary.

- Proof

Consider $\text{Sym}(6)$.



$6 \longrightarrow 6$ This gives a bijection $\pi = (1 \ 4 \ 2)(3 \ 5)(6)$ which is the composition of cycles.

We say that this π has cycle type $3 + 2 + 1$.

- Cycle Type

If instead $\pi = (1 \ 4 \ 2)(3 \ 5 \ 6)$ then the cycle type is given as $3 + 3$.

Finite Symmetric Groups

For $n = 2$, $\text{Sym}(X) = \{\text{Id}, (1\ 2)\}$.

This gives cycle types $1 + 1$ and 2 .

For $n = 3$, $\text{Sym}(X) = \{\text{Id}, (1\ 2), (1\ 3), (2\ 3), (1\ 2\ 3), (1\ 3\ 2)\}$.

This gives cycle types $1 + 1$, $2 + 1$ and 3 .

Symmetric Group for Tetrahedron

For $n = 4$ let $X = \{B, P, W, Y\}$.

Partitions of $n = 4$ are

$4 = 4$	6	$(B\ P\ W\ Y)$	\dots					$\text{sign} = -1$
$= 3 + 1$	$4 \cdot 2 = 8$	$(P\ W\ Y)$	\dots					$\text{sign} = +1$
$= 2 + 2$	$\frac{\binom{4}{2}}{2} = 3$	$(B\ P)(W\ Y)$	$(B\ W)(P\ Y)$	$(B\ Y)(P\ W)$				$\text{sign} = +1$
$= 2 + 1 + 1$	$\binom{4}{2} = 6$	$(B\ P)$	$(B\ W)$	$(B\ Y)$	$(P\ W)$	$(P\ Y)$	$(W\ Y)$	$\text{sign} = -1$
$= 1 + 1 + 1 + 1$	1	Id_X						$\text{sign} = +1$

Rotation Group for Tetrahedron

$$\begin{aligned}
 A &= \{\text{Rotational Symmetries}\} \\
 &= \{\text{Id}_X, 8\ 3\text{-cycles}, 3\ \text{of type } 2+2\}
 \end{aligned}$$

Note, from the sign, that $A \leq \text{Sym}(4)$.

Symmetries Not in Rotation

Why, for example, is $(B\ P)$ not in the rotation group?

If it were, it should be possible to swap vertices and then undo the switch with only rotation.

However, the two tetrahedra are mirror images across a plane.

Observe that the right hand rule with respect to P , W and Y will give opposite, orthogonal vectors.

Rotation as a Subgroup of Symmetry

Q: Is A a subgroup of $\text{Sym}(4)$?

Following the definition, it would be necessary to verify

- $\text{Id} \in A$
- A is closed under inverse.
- A is closed under composition.

Group Homomorphism

Let G and H be groups (whose binary operations are denoted by $g_1 \cdot g_2$).
A (group) homomorphism from G to H is a function $\phi : G \rightarrow H$ such that

- $\phi(g_1 \cdot_G g_2) = \phi(g_1) \cdot_H \phi(g_2)$

Properties of Group Homomorphism

1. $\phi(1_G) = 1_H$

2. $\phi(g^{-1}) = [\phi(g)]^{-1}, \forall g \in G$

- Proof

By definition, $\phi(1_G \cdot 1_G) = \phi(1_G) \cdot \phi(1_G)$.

Letting $e = \phi(1_G)$, we get $e = e \cdot e$.

By multiplying both sides by e^{-1} , we get $1_H = e$.

Part two is left as an exercise.

Example 1

Let $n \geq 1$ and $G = \text{GL}_n(\mathbb{R}) = \{A \in M_{n \times n}(\mathbb{R}) \mid \det(A) \neq 0\}$.

In particular, when $n = 1$, $\text{GL}_1(\mathbb{R}) = \mathbb{R}^* = \{r \in \mathbb{R} \mid r \neq 0\}$ (with multiplication as the binary operation).

Then $\det : G \rightarrow H$ is a group homomorphism.

That is $\det(AB) = \det(A)\det(B)$ (as learned in MATH 21).

Example 2

Let $n \geq 1$, $G = \text{Sym}(n)$, $H = \text{GL}_n(\mathbb{R})$.

Construct a group homomorphism $\rho : G \rightarrow H$.

Recall that a linear transformation $A \in H$ is completely determined by Ae_1, Ae_2, \dots, Ae_n

where $e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \dots, e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$.

For $\pi \in G = \text{Sym}(n)$, $\rho(\pi)$ is the linear transformation that maps e_i to e_j whenever π maps i to j .

This is a surjective linear transformation on a vector space and, therefore, invertible.

- Example

For $n = 4$ and $\pi = (2 \ 3 \ 4)$

$$\begin{array}{ccc} 1 & \longrightarrow & 1 \\ 2 & \searrow & 2 \\ 3 & \nearrow & 3 \\ 4 & \searrow & 4 \end{array}$$

$\rho(\pi)$

$$\begin{array}{ccc}
e_1 & \longrightarrow & e_1 \\
e_2 & \searrow & e_2 \\
e_3 & \nearrow & e_3 \\
e_4 & \searrow & e_4
\end{array}$$

Therefore

$$\rho(\pi) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

– Is this a group homomorphism?

Let $\pi_1, \pi_2 \in G$ be arbitrary elements.

Need to show: $\rho(\pi_1 \circ \pi_2) = \rho(\pi_1) \circ \rho(\pi_2)$.

Both sides are linear transformations and, hence, determined by their actions on e_i for $i = 1, \dots, n$.

$$\begin{aligned}
\rho(\pi_1 \circ \pi_2)e_i &= e_{\pi(i)} \\
&= e_{\pi_1(\pi_2(i))} \\
\rho(\pi_1)(\rho(\pi_2)e_i) &= \rho(\pi_1)(e_{\pi_2(i)})
\end{aligned}$$

Composition of Group Homomorphisms

Let G, H and K be groups and $G \xrightarrow{\phi} H$ and $H \xrightarrow{\psi} K$ be homomorphisms.

Then the composite $\psi \circ \phi : G \rightarrow K$ is a group homomorphism.

Proof

Let $g_1, g_2 \in G$ be arbitrary.

$$\begin{aligned}
(\psi \circ \phi)(g_1 g_2) &= \psi(\phi(g_1 g_2)) && \text{by definition of } \circ \\
&= \psi(\phi(g_1) \phi(g_2)) && \text{since } \phi \text{ is a group homomorphism} \\
&= \psi(\phi(g_1)) \psi(\phi(g_2)) && \text{since } \psi \text{ is a group homomorphism} \\
&= (\psi \circ \phi)(g_1) \circ (\psi \circ \phi)(g_2) && \text{by definition of } \circ
\end{aligned}$$

Definition: Sign Homomorphism

Let $n \geq 1$ and $G = \text{Sym}(n)$.

This sign homomorphism is the composition $\text{sign}: G \xrightarrow{\rho} \text{GL}_n(\mathbb{R}) \xrightarrow{\det} \mathbb{R}^*$

Sign of Symmetric Group

$$\text{sign}(\text{sym}(n)) \subseteq \{1, -1\} \leq \mathbb{R}^*$$

- Lemma

Let a_1, \dots, a_m be distinct numbers between 1 and n .

Then $(a_1 \cdots a_m)$ is equal to $(a_1 \cdots a_{m-1})(a_{m-1} a_m)$.

This will be proven on homework.

- Corollary

Any m cycle is the composition of $m - 1$ transpositions.

Namely, $(a_1, \dots, a_m) = (a_1 a_2)(a_2 a_3) \cdots (a_{m-1} a_m)$.

Easily check: $\text{sign}((a_i a_{i+1})) = -1$.

Now any $g \in \text{Sym}(n)$ allows a cycle decomposition.

Definition: Kernel of a Homomorphism

Let $G \xrightarrow{\phi} H$ be a group homomorphism.

The kernel of ϕ is $\ker(\phi) := \{g \in G \mid \phi(g) = 1_H\}$.

The Kernel is a Subgroup

Let $g_1, g_2 \in \ker(\phi)$. Then

$$\begin{aligned}\phi(g_1 g_2) &= \phi(g_1) \phi(g_2) \\ &= 1_H 1_H \\ &= 1_H\end{aligned}$$

ϕ is a homomorphism

$$g_1, g_2 \in \ker(\phi)$$

$$g_1, g_2 \in \ker(\phi)$$

Similarly, $1_G \in \ker(\phi)$ and $g^{-1} \in \ker(\phi)$ if $g \in \ker(\phi)$. ■

Definition: Alternating Group

Let X be a set, $|X| = n \leq \infty$.

The alternating group on X is the $\text{Alt}(X) = \ker(\text{sign} : \text{Sym}(X) \rightarrow \{\pm 1\})$.

October 5, 2023

Definition: Group Action

Let G be a group and X a set.

A (left) action of G on X is a function $\alpha : G \times X \rightarrow X$ which satisfies two conditions:

1. $\alpha(1_G, x) = x$ for all $x \in X$.
2. $\alpha(g_1, \alpha(g_2, x)) = \alpha(g_1 g_2, x)$ for all $g_1, g_2 \in G$ and $x \in X$.

Notation

Write $\alpha(g, x) = g * x = g \cdot x = gx$.

Example A

Let X be any set, and let $G = \text{Sym}(X) = \{f : X \rightarrow X \text{ bijections}\}$ where the group operation \circ is the composition of functions.

Then G acts (on the left) on X by $f * x \stackrel{\text{def}}{=} f(x)$.

Then the features

1. $\text{Id}_X(x) = x, \forall x \in X$
 2. $g_1 * (g_2 * x) = (g_1 \circ g_2) * x, \forall g_1, g_2 \in G, \forall x \in X$
- Or $g_1(g_2(x)) = (g_1 \circ g_2)(x)$

are satisfied.

Example B

Let $G = \text{Sym}(\{B, P, W, Y\})$ which acts on $X = \{B, P, W, Y\}$.


If $H \leq G$, then H acts on X as well, define $h * x = \dot{h} * x$ (where \dot{h} is regarded as in the alternating group of G).

In particular, $\text{Alt}(\{B, P, W, Y\})$ acts on X by rotations.

Example C*

This example is not required for this class.

From complex Analysis we have the Riemann sphere $\mathbb{P}^1(\mathbb{C}) = \mathbb{C} \cup \{\infty\}$.



Let $G = \text{SL}_2(\mathbb{C})$.

Define G -action on $X = \mathbb{P}^1(\mathbb{C})$ by

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} z := \frac{\alpha z + \beta}{\gamma z + \delta} \quad (\infty \text{ if } \gamma z + \delta = 0)$$

This is called the Möbius group action on $\mathbb{P}^1(\mathbb{C})$.

Exercise: show that 1. and 2. are satisfied.

Definitions

Let G act on X . (Say X is a (left) G -set)

Stabilizer

Let $x \in X$. The stabilizer of x in G is $\text{Stab}_G(x) = \{g \in G \mid g * x = x\} \subseteq G$.

- Example 1

Let G be any group and X a G -set.

Then for any $x \in X$, $\text{Stab}_G(x) \leq G$.

– Proof

1. $1_G \in \text{Stab}_G(x)$ since, by definition, $1_G * x = x$.

Therefore the identity is present.

2. If $g_1, g_2 \in \text{Stab}_G(x)$ are such that $g_1 * x = x$ and $g_2 * x = x$, then $(g_1 g_2) * x \stackrel{\text{2nd Axiom}}{=} g_1 * (g_2 * x) = g_1 * x = x$.

Therefore the stabilizer is closed under composition.

3. Say $g \in \text{Stab}_G(x)$ and $g * x = x$. Apply g^{-1} to both sides to get

$$x \stackrel{\text{1st Axiom}}{=} 1_G * x = (g^{-1} g) * x \stackrel{\text{2nd Axiom}}{=} g^{-1} * (g * x) = g^{-1} * x$$

Therefore the stabilizer is closed under inverse.

- Example 2

Let $G = \text{Alt}(\{B, P, W, Y\})$ and consider $H = \text{Stab}_G(W) = \{\text{Id}, (B P Y), (B Y P)\}$.

Fact: H does not act transitively on X , since W is fixed and no element $g \in H$ satisfies $g * W = B$.

Orbit

Let $x \in X$. The G -orbit of x in X is $G \cdot x = \{g * x \mid g \in G\} \subseteq X$.

Let G act on X and $x, y \in X$. Either $G \cdot x = G \cdot y$ or $G \cdot x \cap G \cdot y = \emptyset$.

So X is the disjoint union of G -orbits.

e.g. $\{B, P, W, Y\} = \{W\} \sqcup \{B, P, Y\}$ gives the $\text{Stab}_G(W)$ -orbits.

- Example 1

When $G = \text{Alt}(X)$, for $X = \{B, P, W, Y\}$, there is only one orbit since $\forall x \in X, G \cdot x = X$.

- Example 2

When $G = \text{Stab}_G(W)$, for $X = \{B, P, W, Y\}$, then $G \cdot W = \{W\}$ while

$$\begin{aligned} G \cdot B &= \{\text{Id}(B), (B P Y)(B), (B Y P)(B)\} = \{B, P, Y\} \\ &= G \cdot P = \{\text{Id}(P), (B P Y)(P), (B Y P)(P)\} = \{P, Y, B\} \\ &= G \cdot Y \end{aligned}$$

Transitivity

Say G acts transitively on X (or the action is transitive) if, for any pair $x, y \in X$, there exists $g \in G$ (depending on x and y) such that $g * x = y$.

- Example

$G = \text{Alt}(\{B, P, W, Y\}) \subset \text{Sym}(\{B, P, W, Y\})$ is transitive.

- Proof

Let $x, y \in X$ be arbitrary.

If $x = y$, then take $g = \text{Id}_X$ and we have $g * x = y$.

Suppose $x \neq y$, then write $X = \{x, y, z, w\}$ and take $g = (x\ y)(z\ w)$. We have $g * x = y$.

e.g. $x = P, y = Y, z = B$ and $w = W$ gives $g = (P\ Y)(B\ W)$.

- Exercise *

This exercise is not required for the course.

Prove that $\text{SL}_2(\mathbb{C})$ acts transitively on $\mathbb{P}^1(\mathbb{C})$.

Say $\mathbb{P}^1(\mathbb{C})$ is a homogeneous space under $\text{SL}_2(\mathbb{C})$.

Proposition: Group Action Gives Group Homomorphisms

(\longrightarrow) Let G act on X . Then

1. For any $g \in G$, the function $\pi_g : X \rightarrow X$ defined by $\pi_g(x) = g * x$ is a bijection of X , hence $\pi_g \in \text{Sym}(X)$.
2. The function $G \xrightarrow{\phi} \text{Sym}(X)$ given by $\phi(g) = \pi_g$ is a group homomorphism.

Proof of 1

Need to show that π_g is injective and surjective.

(Inj) Let $x, y \in X$ and assume $\pi_g(x) = \pi_g(y)$ (i.e. $g * x = g * y$).

Apply $g^{-1} *$ on both sides, such that $x = g^{-1} * (g * x) = g^{-1} * (g * y) = y$.

(Sur) Let $x \in X$ be arbitrary. Need to find $y \in X$ such that $\pi_g(y) = x$.

Take $y = g^{-1} * x$, and $\pi_g(y) = g * (g^{-1} * x) = x$.

Proof of 2

Need to show that $\forall g_1, g_2 \in G, \phi(g_1 g_2) = \phi(g_1) \phi(g_2)$.

$\phi(g_1 g_2) \in \text{Sym}(X)$ is characterized by $[\phi(g_1 g_2)](x) = \pi_{g_1 g_2}(x) = (g_1 g_2) * x$.

On the other hand, $\phi(g_1) \phi(g_2) \in \text{Sym}(X)$ is characterized by $[\phi(g_1) \phi(g_2)](x) = \phi(g_1)[\phi(g_2)(x)] = g_1 * (g_2 * x)$.

By the second group action axiom, these must be the same.

Proposition: Group Homomorphism Admits Group Action

(\longleftarrow) Let $G \xrightarrow{\rho} \text{Sym}(X)$ be a group homomorphism.

Then, by letting $g * x = \rho(g)(x) \in X$ we get a left G -action on X .

Proof

1. $1_G * x = \rho(1_G)(x) = \text{Id}_X(x) = x$.
2. Let $g_1, g_2 \in G$ and $x \in X$. Then $(g_1 g_2) * x = [\rho(g_1 g_2)](x) = [\rho(g_1) \circ \rho(g_2)](x) = \rho(g_1)[\rho(g_2)(x)] = g_1 * (g_2 * x)$.

Definition: Right Group Actions

Let G be a group and X be a set. A right G -action on X is a function $\beta : X \times G \rightarrow X$ such that

1. $\beta(x, 1_G) = x, \forall x \in X$.
2. $\beta(x, g_1 g_2) = \beta(\beta(x, g_1), g_2), \forall g_1, g_2 \in G, \forall x \in X$.

Notation

$$\beta(x, g) = x * g = x \cdot g = xg$$

Remark

If $\alpha : G \times X \rightarrow X$ is a left action, we get a right action $\beta : X \times G \rightarrow X$ by $\beta(x, g) = \alpha(g^{-1}, x)$ and vice versa. That is $x * g = g^{-1} * x$.
Proof recommended as an exercise.

Analogues

Stability, orbit and transitivity all have analogues which can be demonstrated by converting to left actions.

Definition: Cosets

Let $H \leq G$, and let $X = G$.

We have left action $H \times X \rightarrow X$ and $h * x = hx$ (taken in G).

As well as right action $X \times H \rightarrow X$ where $x * h = xh$.

A (left) H -coset is an orbit xH for some $x \in X$.

A (right) H -coset is an orbit Hx for some $x \in X$.

Example

Let $G = \text{Alt}(4)$, $H = \text{Stab}_G(W) = \{\text{Id}, (B P Y), (B Y P)\}$.

1. Take any $x \in H$, $xH = H$.
2. Take $x = (B P)(W Y)$, and $xH = \{(B P)(W Y), (B P)(W Y)(B P Y) = (P W Y), (B P)(W Y)(B Y P) = (B W Y)\}$.

3. There are two more; what are they?

October 10, 2023

Cosets Revisited

Let G be a group, $H \leq G$. Then a (left) H -coset in G is a set of the form

$$gH = \{gh | h \in H\}$$

, where $g \in G$

Coset Space

G/H is the set of H -cosets.

- Example

For $G = \text{Alt}(4)$, given $C_1 = H = \text{Stab}_G(B) = \{1, (P W Y), (P Y W)\}$, we have
 $C_2 = (B P W)H = \{(B P W), (B P)(W Y), (B P Y)\}$
 $(B P W) \circ (P W Y) = (B P)(W Y)$

P \leftarrow B \leftarrow B
 B \leftarrow W \leftarrow P
 Y \leftarrow Y \leftarrow W
 W \leftarrow P \leftarrow Y

$$(B P W) \circ (P Y W) = (B P Y)$$

P \leftarrow B \leftarrow B
 Y \leftarrow Y \leftarrow P
 W \leftarrow P \leftarrow W
 B \leftarrow W \leftarrow Y

$$C_3 = (B W P)H = \{(B W P), (B W Y), (B W)(P Y)\}$$

$$C_4 = (B Y P)H = \{(B Y P), (B Y)(P W), (B Y W)\}$$

Then $G/H = \{C_1, C_2, C_3, C_4\}$.

- Q: What do the 3 elements in C_3 have in common in geometric terms?
 C_3 sends B to W .
 Similarly, the cosets send B to all other vertices (including to itself).

Definition: Transporter

Let G be a group and X a G -set.

For two points, $x, y \in X$, the transporter $\text{Trsp}_G(x, y) = \{g \in G | gx = y\}$.

Example

$$G/H = \{\text{Trsp}_G(B, B), \text{Trsp}_G(B, P), \text{Trsp}_G(B, W), \text{Trsp}_G(B, Y)\}$$

Note

When $x = y$, we recover $\text{Trsp}_G(x, x) = \text{Stab}_G(x)$.

For general G and H , there may not be a nice geometric action associated with it.

But G/H is still a G -set since $g'(gH) = (g'g)H$.

Proposition (B)

Let $H \leq G$ be a subgroup and let $g \in G$.

Then the map $H \xrightarrow{f} gH$ defined by $h \mapsto f(h) = gh$ is a bijection.

Proof

(Surjective) Any element x in gH is, by definition, of the form gh for some $h \in H$. So $x = f(h)$.

(Injective) Say $h_1, h_2 \in H$ satisfy $f(h_1) = f(h_2)$. That is $gh_1 = gh_2$. Multiplying g^{-1} on the left, we get $h_1 = h_2$.

Proposition (C)

Let G act on X , $x \in X$, and $g \in G$.

Take $y := gx$ and $H = \text{Stab}_G(x)$. Then $gH = \text{Trsp}_G(x, y)$.

Proof

(\subseteq) Let $gh \in gH$ be arbitrary. Then

$$(gh) * x \underset{\text{Axiom 2}}{=} g * (h * x) \underset{h \in \text{Stab}_G(x)}{=} g * x \underset{y = g * x}{=} y$$

Therefore $gh \in \text{Trsp}_G(x, y)$.

(\supseteq) Suppose $g' \in \text{Trsp}_G(x, y)$. Consider $g^{-1}g'$. Then

$$(g^{-1}g') * x = g^{-1} * (g' * x) \underset{g' \in \text{Trsp}_G(x, y)}{=} g^{-1}(y) = x$$

Therefore $(g^{-1}g') \in \text{Stab}_G(x)$. Setting $g^{-1}g' := h$, so $g' = gh \in gH$.

Theorem: Orbit-Stabilizer Theorem (OST)

Let G act transitively on a set X (so that there is only one orbit in X , namely X itself).

If $|G| < \infty$, then for any $x \in X$ we have

$$|X| \cdot |\text{Stab}_G(x)| = |G|$$

Proof

Let us count $|G|$ by partitioning G into transporters.

$$G = \bigsqcup_{y \in X} \text{Trsp}_G(x, y)$$

Therefore

$$|G| = \sum_{y \in X} |\text{Trsp}_G(x, y)| \stackrel{B+C}{=} \sum_{y \in X} |\text{Stab}_G(x)| = |X| |\text{Stab}_G(x)| \quad \blacksquare$$

Theorem: Lagrange's Theorem

If G is a finite group and $H \leq G$, then $|G| = |H| \cdot |G/H|$.

Proof (Sketch)

Apply the Orbit-Stabilizer Theorem to $X = G/H$.

This action is transitive as $g(1H) = gH$.

Note $gH = H \iff g \in H$ and $g1 \in H$.

Therefore $\text{Stab}_G(1H) = \{g \in G \mid g(1H) = 1H\} = H$.

Corollary

If $H \leq G$ and $|G| < \infty$, then $|H| \mid |G|$.

The converse is not true. No subgroup of order 6 in $\text{Alt}(4)$ (where $|\text{Alt}(4)| = 12$).

Definition: Conjugate

Let G be a group, $H \leq G$, $g \in G$.

1. For $x \in G$ the g -conjugate of x is $gxg^{-1} = {}^g x$.
2. The g -conjugate of H is $gHg^{-1} = {}^g H = \{gxg^{-1} \mid x \in H\}$.

Example

Let $G = \text{Alt}(4)$ and $H = \text{Stab}_G(B) = \{1, (P W Y), (P Y W)\}$. Then, for $g = (B Y P)$

$$gHg^{-1} = \{1, (B W P), (B P W)\} = \text{Stab}_G(Y)$$

$$(B Y P)1(B P Y) = 1$$

$$(B Y P)(P W Y)(B P Y) = (B W P)$$

$$\begin{array}{l}
W \leftarrow W \leftarrow P \leftarrow B \\
B \leftarrow P \leftarrow Y \leftarrow P \\
P \leftarrow Y \leftarrow W \leftarrow W \\
Y \leftarrow B \leftarrow B \leftarrow Y
\end{array}$$

- Note: Shortcut

$$(gxg^{-1})^{-1} = (g^{-1})^{-1}x^{-1}g^{-1} = gx^{-1}g^{-1}.$$

Applying this to $g = (B \ Y \ P)$ with $x = (P \ W \ Y)$

Therefore, from the previous calculation, $gx^{-1}g^{-1} = (gxg^{-1})^{-1} = (B \ P \ W)$.

Proposition: Geometric Meaning of Conjugate

Let G act on a set X , $x \in X$, $g \in G$, and define $y := g * x$.

Then for $H = \text{Stab}_G(x)$, we have

$$gHg^{-1} = \text{Stab}_G(y)$$

That is, the conjugate of a stabilizer is a stabilizer.

Proof

(\subseteq) Let $ghg^{-1} \in gHg^{-1}$ be arbitrary.

Then

$$(ghg^{-1}) * y = g * (h * (g^{-1} * y)) = g * (h * x) = g * x = y$$

Therefore $ghg^{-1} \in \text{Stab}_G(y)$.

(\supseteq) Let $g' \in \text{Stab}_G(y)$ be arbitrary.

Consider $g^{-1}g'g$. Then

$$(g^{-1}g'g) * x = g^{-1} * (g' * (g * x)) = g^{-1} * (g' * y) = g^{-1} * y = x$$

Therefore $h := g^{-1}g'g \in H$. Then by multiplying g on the left and g^{-1} on the right, we get

$$g' = ghg^{-1} \in gHg^{-1}$$

Orbit-Stabilizer Theorem and Lagrange

1. If G acts transitively on X , then all the stabilizers have the same cardinality because they are all conjugates.

So the Orbit-Stabilizer Theorem is consistent.

2. If $X = G/H$, then $\text{Stab}_G(1H) = H$. What about $\text{Stab}_G(gH) = gHg^{-1}$?

October 12, 2023

Recall: Conjugate

$h \in G$, $H \leq G$, $g \in G$

The conjugates $ghg^{-1} = {}^g h$, $gHg^{-1} = \{{}^g h \mid h \in H\}$.

Meaning

If G acts on X , $x \in X$, $g \in G$, $g * x =: y$, $H = \text{Stab}_G(x)$, then $gHg^{-1} = \text{Stab}_G(y)$.

- Example

$G = \text{Alt}(4)$, $x = B$, $H = \{1, (P W Y), (P Y W)\}$, $g = (B P)(Y W)$
gives $gHg^{-1} = \text{Stab}_G(P) = \{1, (B W Y), (B Y W)\}$.

Definition: Cyclic Subgroup

The cyclic subgroup generated by g is given as $\langle g \rangle = \{g^n \mid n \in \mathbb{Z}\} = \{\dots, g^{-2}, g^{-1}, 1, g, g^1, g^2, \dots\}$.

Definition: Normal Subgroup

Let $H \leq G$.

Say H is normal in G and write $H \trianglelefteq G$ if for all $g \in G$, $gHg^{-1} = H$.

Equivalently, $gH = Hg$ for all $g \in G$.

Trivial Example

If $H = \{1\}$ or $H = G$, then $H \trianglelefteq G$.

Example 1: Non-Normal Subgroup

$G = \text{Alt}(4)$, $H = \text{Stab}_G(B)$ is not normal.

Example 2: Alternating Group

Consider $G = \text{Alt}(4)$, $X = \{x_1, x_2, x_3\}$, $H = \text{Stab}_G(x_1)$.



Then we have the following facts.

- First: Transitivity
 G acts transitively.

- Second: Order of H
By the Orbit-Stabilizer Theorem,

$$|H||X| = |G| \implies |H| \cdot 3 = 12 \implies |H| = 4$$

- Third: Description of G

We know

$$G = \left\{ 1, \begin{pmatrix} 8 \\ 3\text{-cycles} \end{pmatrix}, \begin{pmatrix} 3 \\ 2 + 2\text{-cycles} \end{pmatrix} \right\}$$

- Q: Can a 3-cycle belong to H ?

A: No. Lagrange's Theorem.

Suppose a 3-cycle $g \in H$. Then the cyclic subgroup $\langle g \rangle = \{1, g, g^2\}$, but $\langle g \rangle \leq H$ and $3 \nmid 4$.

- Fourth: Description of H

By Lagrange's Theorem, $H \subseteq \left\{ 1, \begin{pmatrix} 3 \\ 2+2\text{-cycles} \end{pmatrix} \right\} \implies H = \left\{ 1, \begin{pmatrix} 3 \\ 2+2\text{-cycles} \end{pmatrix} \right\}$.

- Fifth: H is a Normal Subgroup

Run the argument with x_1 replaced with x_2 , then $\text{Stab}_G(x_2) = H = \text{Stab}_G(x_3)$.

Therefore, $H \trianglelefteq G$.

Example 3: Non-Normal

From Homework 1, $\mathbb{P}_1(1)$, $H = \{A \in \text{SL}_2\mathbb{C} \mid A^\dagger A = I_2\}$ is not normal.

$\mathbb{P}_1(2)$, $S = \{A = A^\dagger\} \not\trianglelefteq G$.

Example 4: SL As Kernel

$$\text{SL}_n(\mathbb{R}) = \ker(\det \mid \text{GL}_n(\mathbb{R}) \rightarrow \mathbb{R}^\times) \trianglelefteq \text{GL}_n(\mathbb{R})$$

Example 5: Alternating Group As Kernel

$$\text{Alt}(n) = \ker(\text{sign} \mid \text{Sym}(n) \rightarrow \{\pm 1\}) \trianglelefteq \text{Sym}(n)$$

- Proof

Let $g \in G$ and $k \in K$ be arbitrary.

We need to show that $gkg^{-1} \in K$.

$$\phi(gkg^{-1}) \underset{\phi \text{ homomorphism}}{=} \phi(g)\phi(k)\phi(g^{-1}) \underset{k \in K}{=} \phi(g)[\phi(g)]^{-1} = 1_H.$$

Exercise 1

$$\text{Stab}_{\text{Alt}(4)}(1 \text{ of the 6 edges}) \stackrel{?}{\trianglelefteq} \text{Alt}(4).$$

Exercise 2

Can you have a subset closed under conjugation?

On Homework 2 we will examine conjugate classes.

Theorem: Quotient Group

Let $N \trianglelefteq G$. Then on G/N , we have a structure of a group, the quotient group, with the binary operation $g_1N * g_2N := (g_1g_2)N$.

The identity element $1N = N$.

The inverse $(gN)^{-1} = g^{-1}N$.

Proof

The main difficulty is in demonstrating that $*$ is well-defined.

That is, if g'_1, g'_2 are other elements such that (1) $g'_1N = g_1N$ and (2) $g'_2N = g_2N$, then we need to show that $(g_1g_2)N = (g'_1g'_2)N$.

But (1) means that $g'_1 = g_1n_1$ for some $n_1 \in N$, while (2) similarly means $g'_2 = g_2n_2$ for $n_2 \in N$.

It follows that $g'_1g'_2 = g_1n_1g_2n_2$.

Recall that $N \trianglelefteq G$ implies $Ng_2 = g_2N$, so $n_1g_2 = g_2n_3$ for some $n_3 \in N$.

Therefore $g_1n_1g_2n_2 = g_1g_2n_3n_2 \in N$.

Finally, multiplying N on the right side gives $(g_1g_2)N = (g'_1g'_2)N$.

- Associativity

Proof of associativity is left as an exercise.

Remark

If $H \leq G$ and $g_1H * g_2H \stackrel{\text{def}}{=} g_1g_2H$ defines (well) a group, then $H \trianglelefteq G$.

Proposition: Kernel Is Normal

Let $G \xrightarrow{\phi} H$ be a group homomorphism, then $K := \ker(\phi) = \{g \in G \mid \phi(g) = 1_H\}$ is a normal subgroup of G .

Proposition: Abelian Subgroups are Normal

If G is abelian, and $H \leq G$, then $H \trianglelefteq G$.

Proof

$$ghg^{-1} = h$$

Proposition: Subgroup of Index 2

Let $H \leq G$ of index 2 (i.e. $[G : H] = |G/H| = 2$), then $H \trianglelefteq G$.

Proof

Let $g \in G$ be arbitrary.

Need to show that $ogHg^{-1} = H$ or, equivalently, $gH = Hg$.

If $g \in H$, there is nothing to prove. So assume $g \notin H$.

But both gH and Hg are the (set) complement of H in G .



- Example

$$\text{Alt}(n) \trianglelefteq \text{Sym}(n)$$

If $n = 1$, there is nothing to prove.

$$\text{If } n \geq 2, \text{ then } |\text{Alt}(n)| = \frac{n!}{2} = \frac{|\text{Sym}(n)|}{2}.$$

Definition: Simple Group

A group G is simple if G has exactly 2 normal subgroups, namely $\{1\} \neq G$.

Non-Example

$$G = \text{Alt}(4) \supseteq H = \text{Stab}_G(x_1), \quad H \neq \{1\} \text{ or } G$$

So G is not simple.

Proposition: Group of Prime Order

Let p be a prime number.

Any group G of order p is both cyclic and simple.

$$\mathbb{Z}/p\mathbb{Z} = \{\bar{0}, \bar{1}, \bar{2}, \dots, \overline{p-1}\} \text{ with } +.$$

- Proof

Suppose $g \in G$ is not the identity.

$$\text{Form } \langle g \rangle = \{g^n \mid n \in \mathbb{Z}\} =: H \leq G.$$

By Lagrange's Theorem, $|H||G| = p$. Since the only positive divisors of p are 1 and p , $\langle g \rangle = H = G$ for any $g \neq 1_G$.

If $H \leq G$ is any normal subgroup, $|H| \mid |G|$ so $H = \{1\}$ or $H = G$.

Remarks*

For all $n \geq 5$, $\text{Alt}(n)$ is a simple group.

This may be asked as a homework exercise.

If $p \geq 5$ is a prime, then $\text{SL}_2(\mathbb{F}_p)/\{\pm I_2\}$ is a simple group.

Relatedly, if $n \geq 4$ then $\text{SL}_n(\mathbb{F}_p)/Z(\text{SL}_n(\mathbb{F}_p))$ is simple.

The point is that there are infinitely many finite simple groups.

Circa 1980, classification of finite simple groups was announced.

Requires $\sim 10^4$ pages (not everyone has been convinced).

Definition: Center

Let G be a group.

The center $Z(G) = \{z \in G \mid zg = gz, \forall g \in G\}$ is a normal subgroup ($gzg^{-1} = z, \forall g, z$).

More Group Action Terminology

Let G act on a set X .

Recall that this corresponds to a group homomorphism $G \xrightarrow{\rho} \text{Sym}(X)$.

Definition: Faithful

Say the action is faithful if $\ker(\rho) = \{1_G\}$.

$$\forall g \in G : gx = x, \forall x \in X \implies g = 1_G$$

Definition: Free

Say the action is free if $\forall g \in G, \forall x \in X, gx = x \implies g = 1_G$.

$$\text{Stab}_G(x) = \{1_G\}, \forall x \in X$$

Example 1

Let G act on $X = G$ by left multiplication. Then this is free.

Example 2

$$G = \text{Alt}(4)$$

$X_1 = \{B, P, W, Y\}$ is faithful but not free.

$X_2 = \{6 \text{ edges}\}$ is faithful but not free.

$X_3 = \{3 \text{ strings}\}$ is not faithful.

October 17, 2023

Recall: Normal Subgroup

A subgroup $N \leq G$ is normal & $N \trianglelefteq G$, if $gNg^{-1} = N$ for all $g \in G$.

Notation

$$\begin{aligned} G &\xrightarrow{\pi} G/N \\ g &\mapsto gN \end{aligned}$$

Recall: Quotient Group

We have constructed, for $N \trianglelefteq G$, the quotient group G/N , with $g_1N * g_2N = g_1g_2N$.

Definition: Group Homomorphisms

For groups G, K , $\text{Hom}(G, K) = \{G \rightarrow K \mid \text{group homomorphisms}\}$.

Theorem: Universal Mapping Property of the Quotient Group

Let H be any group.

Part 1

For any group homomorphism $\bar{f} : G/N \rightarrow H$, by composing, we get a homomorphism

$$\bar{f} \circ \pi = f : G \xrightarrow{\pi} G/N \xrightarrow{\bar{f}} H$$

such that $f(N) = \{1_H\}$.

- Proof

Let $\bar{f} \in \text{Hom}(G/N, H)$ and $f := \bar{f} \circ \pi \in \text{Hom}(G, H)$.

Then

$$f(n) = \bar{f}(\pi(n)) = \bar{f}(nN) = \bar{f}(N) \underset{N \in G/N \text{ is the identity}}{=} 1_H$$

Part 2

Conversely, for any group homomorphism $f : G \rightarrow H$ such that $f(N) = \{1_H\}$ we get a unique homomorphism $\bar{f} : G/N \rightarrow H$ such that $f = \bar{f} \circ \pi$.

- Proof

- Well Defined

Let $f \in \text{Hom}(G, H)$ satisfy $f(N) = \{1_H\}$.

Define $\bar{f}(gN) = f(g)$ for every $g \in G$.

Need to Show: If $g_1N = g_2N$, then $f(g_1) = f(g_2)$.

But $g_1N = g_2N$ implies, since $g_1 1_G \in g_1N$, that $g_1 = g_2n$ for some $n \in N$.

Since f is a group homomorphism, we have

$$f(g_1) = f(g_2)f(n) \underset{f(N)=\{1_H\}}{=} f(g_2)1_H = f(g_2)$$

- Homomorphism

\bar{f} is a homomorphism since for any $g_1N, g_2N \in G/N$, we have

$$\bar{f}(g_1N * g_2N) \underset{\text{definition of } *}{=} \bar{f}(g_1g_2N) \underset{\text{definition of } \bar{f}}{=} f(g_1g_2) \underset{f \text{ is homomorphism}}{=} f(g_1)f(g_2) \underset{\text{definition of } \bar{f}}{=} \bar{f}(g_1N)\bar{f}(g_2N)$$

- Uniqueness

Suppose $l \in \text{Hom}(G/N, H)$ is any element satisfying $f = l \circ \pi$.

Then for any $g \in G$, we have

$$\bar{f}(gN) = f(g) = l(\pi(g)) = l(gN)$$

Therefore $\bar{f} = l$.

Rephrase: Universal Mapping Property of Quotient Group

$$\text{Hom}(G/N, H) \xrightarrow{1} \{f \in \text{Hom}(G, H) \mid f(N) = \{1_H\}\}$$

Equivalently, $f(N) = \{1_H\} \iff N \leq \ker(f)$.

Note: $f(N) = \{1_H\} \iff N \subseteq f^{-1}(\{1_H\}) = \ker(f)$

Loosely: Universal Mapping Property of Quotient Group

Giving a homomorphism $G/N \rightarrow H$ is the same as giving a homomorphism $G \rightarrow H$ that kills N .

Definition: Group Generators

Let G be a group and let $S \subseteq G$.

Definition: Word

A word in S of length $l \geq 1$ is an expression $x_1 x_2 \cdots x_l$ where $x_i \in \{s, s^{-1}\}$ for some $s \in S$ and $i = 1, \dots, l$.

The word of length zero is, by convention, 1_G .

Definition: Generated Subgroup

The subgroup generated by S , $\langle S \rangle$ is the subset of G consisting of all the words in S of all possible lengths.

Fact: $\langle S \rangle \leq G$.

- Example

$$S = \{A, C, I, T, L\}.$$

One word in S of length 3 is $CA^{-1}T$.

Inverse: $(CA^{-1}T) = T^{-1}AC^{-1}$.

Composition: $CA^{-1}T * T^{-1}A^{-1}IL = CA^{-1}TT^{-1}A^{-1}IL = CA^{-2}IL$. (Note that, strictly, $A^{-2} \notin S$ but rather $A^{-2} \equiv A^{-1}A^{-1}$).

Common Usage

We usually do not use the bare definition to describe what $\langle S \rangle$.

- Example

Let $G = \text{Sym}(n)$ and let $S = \{\text{transpositions } (a\ b) \mid 1 \leq a < b \leq n\}$.

Then $\langle S \rangle = G$.

- Proof

Step 1: Any $\pi \in G$ has a cycle decomposition. So $\pi = \gamma_1 \gamma_2 \cdots \gamma_k$, where γ_i is an l_i -cycle for some $l_i \geq 1$.

Step 2: If γ is an l -cycle, $l \geq 2$, say $\gamma = (a_1\ a_2\ \cdots\ a_l)$, then $\gamma \stackrel{\text{HW1}}{=} (a_1\ a_2)(a_2\ a_3) \cdots (a_{l-1}\ a_l)$.

Definition: Commutator

A commutator in G is an element of the form $[g, h] := ghg^{-1}h^{-1}$ for some $g, h \in G$.

Definition: Commutator Subgroup

The commutator subgroup (or derived subgroup) $D(G)$ (sometimes $[G, G]$) is the subgroup of G generated by all the commutators.

That is, a typical element in $D(G)$ is $C_1 C_2 \cdots C_l$ where $C_i = [g_i, h_i]$ for some $g_i, h_i \in G, \forall i$.

Note that $[g, h]^{-1} = (ghg^{-1}h^{-1})^{-1} = hgh^{-1}g^{-1} = [h, g]$.

Proposition: Commutator Subgroup Is Normal

$$D(G) \trianglelefteq G.$$

Lemma:

Let $g_0 \in G$ be fixed.

The map $G \rightarrow G$ defined as $g \mapsto g_0 g g_0^{-1}$ – called the inner homomorphism and denoted $\text{Int}(g_0)$ – is a group homomorphism.

- Proof

$$\text{Int}(g_0)(g_1 g_2) = g_0 g_1 g_2 g_0^{-1} = g_0 g_1 g_0^{-1} g_0 g_2 g_0^{-1} = [\text{Int}(g_0)(g_1)][\text{Int}(g_0)(g_2)]$$

Proof

Let $g_0 \in G$ and $C_1 \cdots C_l \in D(G)$ be arbitrary.

$$g_0(C_1 \cdots C_l)g_0^{-1} = \text{Int}(g_0)(C_1 \cdots C_l) = \underset{\text{lemma}}{=} [\text{Int}(g_0)(C_1)] \cdots [\text{Int}(g_0)(C_l)]$$

It suffices to prove that $\text{Int}(g_0)(C)$ is a commutator for any commutator C .

$$\text{Int}(g_0)(C) = \text{Int}(g_0)(ghg^{-1}h^{-1}) = [\text{Int}(g_0)(g), \text{Int}(g_0)(h)] \quad \blacksquare$$

Definition: Abelianization

The quotient group $G/D(G)$ is called the Abelianization G^{ab} of G .

Property: Commutativity

G^{ab} is commutative.

- Proof

Let $D = D(G)$ and let $gD, hD \in G^{\text{ab}}$ be arbitrary.

Then

$$[gD, hD] = (gD)(hD)(gD)^{-1}(hD)^{-1} = ghg^{-1}h^{-1}D = D = 1_{G^{\text{ab}}}$$

Therefore, by multiplying on the right by $(hD)(gD)$, we get

$$(gD)(hD) = (hD)(gD)$$

So G^{ab} is Abelian.

Note: Abelian Groups

G is Abelian if and only if $D(G) = \{1_G\}$.

Proposition: Minimal Normal Subgroup with Commutative Quotient

$D(G)$ is the smallest normal subgroup $N \trianglelefteq G$ such that G/N is commutative.

Proof

Suppose $N \trianglelefteq G$ has Abelian G/N .

Need to Show: $D(G) \leq N$.

So let $g, h \in G$ be arbitrary and let $C = ghg^{-1}h^{-1}$.

Then in G/N , $[gN, hN] = 1_{G/N} = N$.

Therefore $[g, h] \in N$ and $[G, G] \leq N$. ■

Remark*: Homology in Algebraic Topology

Let X be a connected manifold, say

IMAGE HERE - DOUBLE TORUS WITH CHUNK BEING REMOVED

In Algebraic Topology, we study loops up to homotopy $G = \pi_1(X, x_0)$.

Then $G^{\text{ab}} = \underset{\text{Hurewicz}}{H_1(X; \mathbb{Z})}$ is a homology.

Theorem: Universal Mapping Property of Abelianization

Let G be any group and H be any Abelian group.

Then we have a bijection $\text{Hom}(G, H) \cong \text{Hom}(G^{\text{ab}}, H)$.

Proof

Since H is Abelian, any homomorphism $f : G \rightarrow H$ will satisfy $f([G, G]) = \{1_H\}$.

So use UMP for quotient G/N . ■

October 19, 2023

Review: Commutator

Let G be a group.

A commutator is an element of the form $[g, h] = ghg^{-1}h^{-1}$.

The derived (or commutator) subgroup $DG = [G, G]$ is the subgroup generated by all the commutators.

Properties of Commutators

$DG \trianglelefteq G$

$G^{\text{ab}} := G/DG$ is abelian.

G^{ab} satisfies the universal mapping property. That is, for all abelian H .

$$\text{Hom}(G, H) \cong \text{Hom}(G^{\text{ab}}, H)$$

Example: G Abelian

If G is abelian, then $D(G) = \{1_G\}$ and $G \xrightarrow{\sim} G^{\text{ab}} = G/\{1_G\}$ (usually written $G = G^{\text{ab}}$).
 Conversely, $DG = \{1_G\}$ implies G is abelian.

Example: G Simple and Non-abelian

If G is simple and non-abelian, then $DG = G$.

Definition: Perfect Group

A group G is called perfect if $D(G) = G$.

cf. Poincaré Conjecture (Perelman circa 2002: Every closed 3-manifold M with $\pi_1(M) = \{1\}$ is homeomorphic to S^3).

Example: G Symmetric Group 3

Let $G = \text{Sym}(3)$.

By Lagrange, if $H \leq G$, then $|H| \mid |G| = 3! = 6$. Consider

$$|H| = 1, 2, 3, \text{ or } 6$$

Then



Which one is DG ?

It cannot be $\{1\}$ since $G \neq G^{\text{ab}}$.

It cannot be $\langle \tau \rangle$, $\tau^2 = 1$, since they are not normal.

$G/\langle (1\ 2\ 3) \rangle$ is a group of order 2 and, therefore, abelian.

Recall

DG is the smallest normal subgroup N of G such that G/N is abelian.

Therefore, $DG = D(\text{Sym}(3)) = \langle (1\ 2\ 3) \rangle$.

Note

For $G = \text{Sym}(3)$,

$$G \supsetneq_{\neq} D(G) = \langle (1\ 2\ 3) \rangle \supsetneq_{\neq} D(D(G)) = \{1\}$$

Definition: Solvable Group

Suppose G is a group such that

$$D(D(D(\cdots D(G)\cdots)) = \{1\}$$

Then G is solvable (or soluble).

Definition: Conjugacy Class

The conjugacy class of $\bar{x} \in G$ is the set of elements in G that are conjugate to \bar{x} .

Let G act on (the set) $G = X$ by conjugation: $g * x = gxg^{-1}$.

Then the conjugacy class of x is the G -orbit of $x \in X$.

Definition: Centralizer

The centralizer $x \in G$ is the stabilizer of x in this conjugation action.

$$C_G(x) = \{g \in G \mid gxg^{-1} = x\} \iff gx = xg$$

Example: G Alternating Group 4

Let $G = \text{Alt}(4)$.

Then, by Lagrange, $H \leq G \implies |H| \mid |G| = 12$.

So $|H| = 1, 2, 3, 4, 6$, or 12



Note: $\text{Stab}(\text{String})$ refers to the stabilizer of x_1 as depicted in Example 2 on page 19.

Proposition: No Order 6 Subgroup

No subgroup H of order 6 in $\text{AAlt}(4)$.

- Proof

Suppose $H \leq G$ has order six.

Then $[G : H] = \frac{12}{6} = 2$ and, therefore, $H \trianglelefteq G$.

So H must be a union of certain conjugacy classes.

From Homework 1, G acts transitively on $\{6 \text{ edges}\}$, so $\text{Stab}(\overline{BP})$, $\text{Stab}(\overline{BW})$, and $\text{Stab}(\overline{BY})$ are conjugate to each other (HW2 P1(b)).

Therefore the 3 elements of order 2, $T = \{(B P)(W Y), (B W)(P Y), (B Y)(P W)\}$ form a conjugacy class in G

It follows that $G = \{1\} \sqcup T \sqcup \{8 \text{ elements of order } 3\}$.

By the following lemma, this resolves to

$$G = \{1\} \sqcup T \sqcup \{p\}^\# \sqcup \{p^2\}^\#$$

- Lemma (Key)

Every element p of order 3 in $G = \text{Alt}(4)$ has conjugacy class of size 4.

– Proof

$|(\text{conjugacy class of } p)| \geq 4$ since G acts transitively on $\{B, P, W, Y\}$ and $\exists p \in \text{Stab}(P)$ such that, without loss of generality, $gpg^{-1} = \text{Stab}(B)$.

Since $p = 1$ implies $gpg^{-1} = 1$, each g sends non-trivial p to either $(P W Y)$ or $(P Y W)$ in $\text{Stab}(B)$. Then, also

$$|(\text{conjugacy class of } p)| = |\text{orbit of } p \text{ under the conjugation action}| = |G|/|C_G(p)| \leq 12/3 = 4$$

Therefore no normal group of order 6 is in $G = \text{Alt}(4)$.

Example Continued: 6 Alternating Group 4

$\{1\}$ is non-abelian.

$\text{Stab}(\overline{BP})$ inhabits a conjugate class.

Because $|G/\text{Stab}(\text{String})| = 3$ it is cyclic and $DG = \{1\} \coprod T$

Observe that DG is abelian.

Proof

What to Show: For $t = (B P)(W Y)$ and every $t \in T$, $C_G(t) \supseteq DG$.

Then

$$|C_G(T)| \stackrel{OST}{=} \frac{|G|}{|T|} = \frac{12}{3} = 4$$

Therefore, $C_G(t) = DG$.

Remark: Enumerability of Subgroups

There are very few groups G such that we can enumerate all the subgroups of G .

Theorem: Product of Integers is a Subgroup

Let $G = (\mathbb{Z}, +)$.

1. For any $a \in \mathbb{Z}$, the subset $a\mathbb{Z} = \{an \mid n \in \mathbb{Z}\}$ is a subgroup of G .

2. Conversely, any subgroup $H \leq \mathbb{Z}$ is of the form $H = a\mathbb{Z}$ for some $a \in G$.

Proof of 1

Need to Show: $a\mathbb{Z}$ contains 0, is closed under $+$ and is closed under (additive) inverse.

But, $0 = a0$, and $an + am = a(n + m)$ and $-(an) = a(-n)$, for $n, m \in \mathbb{Z}$.

Proof of 2

Suppose $H \leq (\mathbb{Z}, +)$.

Since $+$ is commutative, all subgroups will be normal.

If $H = \{0\}$, then $h = 0\mathbb{Z}$, and we're done.

If not, say $H \ni x \neq 0$, then $h \ni -x$.

Therefore $S = \{x \in H \mid x > 0\}$ must be non-empty.

Let a be the smallest element in S .

Then we claim $H = a\mathbb{Z}$.

- Proof of the Claim

(\supseteq) Since $H \ni a$ and $H \leq G$.

(\subseteq) Let $x \in H$ be arbitrary. Apply Division Algorithm and get

$$x = aq + r$$

where $q, r \in \mathbb{Z}$ and $0 \leq r < a$.

If $r > 0$, then

$$r = x - aq \in H$$

so $r \in S$.

However, since $r < a$, this contradicts the very choice of a as the smallest element.

So $r = 0$.

But then $x = aq \in a\mathbb{Z}$.

Therefore $H \subseteq a\mathbb{Z}$. ■

Homework 2 Question

If $H \leq G$, then $gHg^{-1} \leq G$.

Consider the set X_G of all the subgroups H of G .

The group G acts on X_G by conjugation: $g * H := gHg^{-1}$.

Definition: Normalizer

The normalizer of H in G is $N_G(H) = \{g \in G \mid gHg^{-1} = H\}$.

October 24, 2023

Isomorphism Theorems

Definition: Isomorphism

An isomorphism between two groups, G and H , is a group homomorphism $f : G \rightarrow H$ that is bijective.

Two groups are said to be isomorphic if there is an isomorphism between them.

Example

Let G be any group, $N = \{1_G\}$, and let $\pi : G \rightarrow G/N$ be the canonical projection.

Then $gN = \{g\}, \forall g \in G$.

So π is an isomorphism.

e.g. for $G = \{1, a, b, c\}$

$G/\{1_G\} = \{\{1\}, \{a\}, \{b\}, \{c\}\}$

1 - $\{1\}$

a - $\{a\}$

b - $\{b\}$

c - $\{c\}$

Here we will write $G = G/\{1_G\}$.

Lemma:

Let $G \xrightarrow{\phi} H$ be a group homomorphism.

Suppose ϕ is surjective. Form $K = \ker(\phi) \leq G$.

Then we have an isomorphism, induced by ϕ :

$$\bar{\phi} : G/K \rightarrow H$$

Proof

Giving $\bar{\phi}$ is the same as giving a homomorphism $G \xrightarrow{\psi} H$ such that $\psi(K) = \{1_H\}$ by the Universal Mapping Property.

So take $\psi = \phi$.

So have a group homomorphism $\bar{\phi}(gK) = \phi(g), \forall g \in G$.

$$\begin{array}{ccc} G & \xrightarrow{\phi} & H \\ & \searrow \pi & \nearrow \bar{\phi} \\ & K & \end{array}$$

Set Theory

ϕ surjective $\implies \bar{\phi}$ surjective.

Remains to show: $\bar{\phi}$ is injective.

Let $g_1K, g_2K \in G/K$ such that $\bar{\phi}(g_1K) = \bar{\phi}(g_2K)$.

Then, by definition of $\bar{\phi}$,

$$\phi(g_1) = \phi(g_2)$$

So

$$\phi(g_1g_2^{-1}) \underset{\phi \text{ homomorphism}}{=} \phi(g_1\phi(g_2^{-1})) = 1_H.$$

Therefore $g_1g_2^{-1} \in \ker(\phi) = K$ and $g_1K = g_2K$. ■

Definition: Normalizer (Revisited)

Let $H \leq G$. The normalizer $N_G(H)$ is $\{g \in G \mid gHg^{-1} = H\}$.

Note: Normalizer is a Stabilizer

It is the stabilizer of H in the conjugation action of G on the set of all subgroups of G .

Since $\forall h \in H$, we have $hHh^{-1} = H$, we have $H \leq N_G(H)$.

Since $N_G(H)$ is a stabilizer, $N_G(H) \leq G$.

Therefore $H \leq N_G(H) \leq G$.

$H \leq N_G(H) \leq G$.

Note: H is Normal If the Normalizer of H is G

$H \trianglelefteq G \iff N_G(H) = G$.

Example 1

Let $G = \text{Alt}(4)$, $K_1 = \text{Stab}_G(B) = \langle (P \ W \ Y) \rangle$, and $H_1 = \langle (B \ P)(W \ Y) \rangle$.

Question: What are $N_G(H_1)$ and $N_G(K_1)$?

- H1

H_1 is not normal in G , so $N_G(H_1) \neq G$.

Since $H_1 \leq N_G(H_1) \leq G$.

Recall that $H_1 \leq \text{Stab}_G(\text{String}) \leq G$ and $\text{Stab}_G(\text{String})$ is abelian.

Therefore $N_G(H_1) = \text{Stab}_G(\text{String})$.

Note also: $N_G(\text{Stab}_G(\text{String})) = G$.

- K1

From homework, $\forall g \in G$, $gK_1g^{-1} = \text{Stab}_G(gB)$ and, if $gB \neq B$, then $\text{Stab}_G(gB) \neq K_1$.

So $K_1 \leq N_G(K_1) \subseteq \text{Stab}_G(B) = K_1$.

That is, $N_G(K_1) = K_1$. It is a “self-normalizer.”

Example 2

Let $G = \text{GL}_2(\mathbb{R})$, $H = \left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{R} \right\}$.

What is $N_G(H)$?

Say $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ belongs to $N_G(H)$.

Then $g \begin{pmatrix} 1 & 3 \\ 0 & -1 \end{pmatrix} g^{-1} \in H$. Say

$$g \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix} g^{-1} = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$$

for some $x \in \mathbb{R}$. Then

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$\begin{pmatrix} a & b+3a \\ c & d+3c \end{pmatrix} = \begin{pmatrix} a+xc & b+xd \\ c & d \end{pmatrix}$$

Therefore, $c = 0$ (and $x = \frac{3a}{d}$). So

$$N_G(H) \leq \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \mid a, d \in \mathbb{R}^+, b \in \mathbb{R} \right\}$$

Are they equal?

$$\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}^{-1} = \begin{pmatrix} a & b+ax \\ 0 & d \end{pmatrix} \frac{1}{ad} \begin{pmatrix} d & -b \\ 0 & a \end{pmatrix} = \begin{pmatrix} 1 & ? \\ 0 & 1 \end{pmatrix} \in H$$

Therefore, for $g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}$, $gHg^{-1} \in H$.

Now, apply the same argument to $g^{-1} = \begin{pmatrix} 1/a & -b/ad \\ 0 & 1/d \end{pmatrix}$ and get $g^{-1}Hg \subseteq H$.

That is to say, $H \subseteq gHg^{-1}$.

It follows that $(A) + (B) \implies g \in N_G(H)$, and

$$N_{\text{GL}_2(\mathbb{R})} \left(\left\{ \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \mid x \in \mathbb{R} \right\} \right) = \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \mid a, d \in \mathbb{R}^+, b \in \mathbb{R} \right\}$$

Definition: Product

Let $H, K \leq G$.

Then $HK := \{hk \mid h \in H, k \in K\}$

HK need not be a subgroup, and may differ from KH .

Example

Let $G = \text{Alt}(4)$, $H = \langle (B P)(W Y) \rangle$, and $K = \langle (P W Y) \rangle$.

Note $|H| = 2$ and $|K| = 3$. Then

$$HK = \{1 \cdot 1, 1(P W Y), 1(P Y W), (B P)(W Y)1, (B P)(W Y)(P W Y), (B P)(W Y)(P Y W)\}$$

Fact

If $H, K \leq G$ such that $H \cap K = \{1_G\}$ and $h_1, h_2 \in H$ and $k_1, k_2 \in K$, then

$$h_1 k_1 = h_2 k_2 \implies h_1 = h_2 \text{ and } k_1 = k_2$$

Proof

Multiply h_2^{-1} on the left and k_1^{-1} on the right. Then

$$H \ni h_2^{-1} h_1 = k_2 k_1^{-1} \in H \quad \blacksquare$$

Therefore $|HK| = 6$.

Since G contains no subgroup of order 6, $HK \not\leq G$.

Theorem: First Isomorphism Theorem

Note: Dummit and Foote lists this as the Second Isomorphism Theorem.
Let $H, K \leq G$ and assume that $H \leq N_G(K)$,

1. $HK = KH$ is a subgroup of G ,
2. $K \trianglelefteq HK$ and $H \cap K \trianglelefteq H$, and
3. $HK/K \xrightarrow{\sim} H/H \cap K$ is an isomorphism of (quotient) groups.

Proof of 1

- Equality

Need to prove $HK \subseteq KH$ and $KH \subseteq HK$.

(\implies) Let $h \in H$ and $k \in K$. We want $hk \in KH$.

Since $h \in N_G(K)$, $hKh^{-1} = K$.

So, in particular, $hkh^{-1} \in K$ for some $k_2 \in K$. Multiplying by h on the right,

$$hk = k_2h \in KH$$

(\impliedby) Let $kh \in KH$.

Since $H \in N_G(K)$ and $N_G(K)$ is a subgroup, $h^{-1} \in N_G(K)$.

So $h^{-1}K(h^{-1})^{-1} = K$.

In particular, $h^{-1}kh = k_3$ for some $k_3 \in K$. So

$$kh = h_{k_3} \in HK$$

- Subgroup

Identity: $1 = 1 \cdot 1 \in HK$

Product: Let h_1k_1 and h_2k_2 be arbitrary. Then for some $h_4 \in H, k_4 \in K$

$$(h_1k_1)(h_2k_2) = h_1(k_1h_2)k_2 \stackrel{KH=HK}{=} h_1(h_4k_4)k_2 = \overset{\in H}{(h_1k_4)}\overset{\in K}{(k_4k_2)} \in HK$$

So HK is closed under the group operation.

Inverse: $(hk)^{-1} = k^{-1}h^{-1} \in KH = HK$.

Proof of 2

- K is Normal in HK

Let $hk \in HK$ be arbitrary.

Since $h \in H \leq N_G(K)$, $k \in K \leq N_G(K)$, and $N_G(K)$ is a subgroup,

$$hk \in N_G(K)$$

Therefore $HK \leq N_G(K)$ and K is normal in HK .

- H Meet K is Normal in H

For $h \in H$, $hKh^{-1} = K$ by assumption, and $hHh^{-1} = H$.

Therefore $h(H \cap K)h^{-1} = H \cap K$, and $K \cap H$ is normal in H .

Proof of 3

Let us start with the inclusion homomorphism

$$\begin{array}{ccccc} h & \longmapsto & h \cdot 1 & \longmapsto & hK \\ H & \longrightarrow & HK & \longrightarrow & HK/K \\ & \searrow \phi & & \nearrow & \end{array}$$

ϕ is surjective.

A typical element in HK/K is $hK = hK$ which is the image of $h \in H$ under ϕ .

Then $\ker(\phi) = \{h \in H \mid hK = 1_K\} \iff h \in K$. This is $H \cap K$.

Therefore, by lemma, $H/H \cap K \xrightarrow[\sim]{\bar{\phi}} HK/K$.

Example

Let $G = \text{GL}_3(\mathbb{R})$ and $H = \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mid x, y, z \in \mathbb{R} \right\} . H \leq G$ called the Heisenberg group of dimension .

October 26, 2023

Heisenberg Groups

$$G = \text{Heis}_3(\mathbb{R}) = \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} \mid x, y, z \in \mathbb{R} \right\} \leq \text{GL}_3(\mathbb{R})$$

Subgroup

1. $I_3 \in G$ since we can take $x = y = z = 0$.

2. (Empty elements are read as zeros.)

$$\underbrace{\begin{pmatrix} 1 & x & z \\ & 1 & y \\ & & 1 \end{pmatrix}}_{=g} \underbrace{\begin{pmatrix} 1 & a & c \\ & 1 & b \\ & & 1 \end{pmatrix}}_{=h} \begin{pmatrix} 1 & x+a & z+c+xb \\ & 1 & y+b \\ & & 1 \end{pmatrix}$$

So closed under matrix multiplication.

Note: (the $(1,2)$ -component of gh) = (the $(1,2)$ -component of g) + (the $(1,2)$ -component of h).

Ditto for the $(2,3)$ -component.

1. G is closed under inverse.

Let $a = -x$, $b = -y$ and $c = -z + xy$. Then

$$\begin{pmatrix} 1 & x & z \\ & 1 & y \\ & & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & -x & xy - x \\ & 1 & -y \\ & & 1 \end{pmatrix} \in G$$

Example

Identify $\mathbb{R}^3 = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} \right\}$ with G and let G act on $\mathbb{R}^3 = X$ by left multiplication. i.e.

$$g = \begin{pmatrix} 1 & x & z \\ & 1 & y \\ & & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x \\ y \\ z \end{pmatrix} \rightsquigarrow g * \vec{v} = \begin{pmatrix} 1 & a & c \\ & 1 & b \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & x & z \\ & 1 & y \\ & & 1 \end{pmatrix} = \dots$$

For example, if

$$\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \leftrightarrow I_3 \quad \text{and} \quad g = \begin{pmatrix} 1 & 0 & 0 \\ & 1 & z \\ & & 1 \end{pmatrix}$$

then g maps $\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$ to $\begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}$ while

$$h = \begin{pmatrix} 1 & 3 & 0 \\ & 1 & 0 \\ & & 1 \end{pmatrix} \quad \text{maps} \quad \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad \text{to} \quad \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}.$$

However, since this is a noncommutative group

$$\begin{aligned} gh &= \begin{pmatrix} 1 & 3 & 0 \\ & 1 & 2 \\ & & 1 \end{pmatrix} \quad \text{maps} \quad \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad \text{to} \quad \begin{pmatrix} 3 \\ 2 \\ 0 \end{pmatrix} \\ hg &= \begin{pmatrix} 1 & 3 & 0 \\ & 1 & 0 \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ & 1 & 2 \\ & & 1 \end{pmatrix} = \begin{pmatrix} 1 & 3 & 6 \\ & 1 & 2 \\ & & 1 \end{pmatrix} \end{aligned}$$

So hg maps $\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$ to $\begin{pmatrix} 3 \\ 2 \\ 6 \end{pmatrix}$.

So G acts on \mathbb{R}^3 transitively and freely.

This is one of the eight “model geometries” of William Thurston.

Heisenberg Group Center and Derived Group

Recall

The center

$$Z(G) = \{g \in G \mid gh = hg, \forall h \in G\}$$

and the derived group

$$D(G) = \langle \{[g, h] \mid g, h \in G\} \rangle.$$

Center of Heisenberg Group

If

$$g = \begin{pmatrix} 1 & a & c \\ & 1 & b \\ & & 1 \end{pmatrix} \in Z(G),$$

then for any $x, y, z \in \mathbb{R}$, we have

$$\begin{aligned} \begin{pmatrix} 1 & a & c \\ & 1 & b \\ & & 1 \end{pmatrix} \overbrace{\begin{pmatrix} 1 & x & z \\ & 1 & y \\ & & 1 \end{pmatrix}}^{=h} &= \begin{pmatrix} 1 & x & z \\ & 1 & y \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & a & c \\ & 1 & b \\ & & 1 \end{pmatrix} \\ \begin{pmatrix} 1 & a+x & c+z+ay \\ & 1 & b+y \\ & & 1 \end{pmatrix} &= \begin{pmatrix} 1 & x+a & z+c+xb \\ & 1 & y+b \\ & & 1 \end{pmatrix} \end{aligned}$$

Therefore $ay = xb$ for all $x, y, z \in \mathbb{R}$.

Take $(x, y) = (1, 0)$, and we get $0 = b$.

Take $(x, y) = (0, 1)$, and we get $a = 0$.

It follows that

$$Z(G) = \left\{ \begin{pmatrix} 1 & 0 & c \\ & 1 & 0 \\ & & 1 \end{pmatrix} \mid c \in \mathbb{R} \right\}$$

Derived Group of Heisenberg Group

$[g, h] = ghg^{-1}h^{-1}$ has

$$(1, 2) - \text{component} = a + x + (-a) + (-x) = 0$$

$$(2, 3) - \text{component} = b + y + (-b) + (-y) = 0$$

So $D(G) \leq Z(G)$ for this G .

- Q: Is the center a subgroup of the derived group?

Form

$$\begin{aligned} G/Z(G) &\xrightarrow[\phi]{\sim} (R^2, +) \\ \begin{pmatrix} 1 & a & 0 \\ & 1 & b \\ & & 1 \end{pmatrix} Z(G) &\longmapsto (a, b) \end{aligned}$$

ϕ is a group homomorphism since the $(1,2)$ -component and $(2,3)$ component of gh are the sum of those components in g and h .

ϕ is surjective, since

$$\begin{pmatrix} 1 & a & c \\ & 1 & b \\ & & 1 \end{pmatrix} Z(G) = \begin{pmatrix} 1 & a & 0 \\ & 1 & b \\ & & 1 \end{pmatrix} Z(G)$$

where the matrix on either side differ by only an element in $Z(G)$.

ϕ is injective, since if

$$\phi(a, b) = \begin{pmatrix} 1 & a & 0 \\ & 1 & b \\ & & 1 \end{pmatrix} \in Z(G)$$

, then $a = b = 0$. So $(a, b) = 0$ in \mathbb{R}^2 .

This is a cool isomorphism, but it only demonstrates $D(G) \leq Z(G)$.

- Q: Is the center a subgroup of the derived group? (Take Two)

$$\begin{aligned} \begin{pmatrix} 1 & 1 & \\ & 1 & \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & \\ & 1 & c \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 & \\ & 1 & \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & \\ & 1 & \\ & & -c \end{pmatrix} \\ = \begin{pmatrix} 1 & 1 & c \\ & 1 & c \\ & & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 & c \\ & 1 & -c \\ & & 1 \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 & c \\ & 1 & 0 \\ & & 1 \end{pmatrix} \end{aligned}$$

Therefore $Z(G) = D(G)$.

Theorem: Correspondence + 2nd Isomorphism Theorem

Let $N \trianglelefteq G$, $E_1 = \{H \leq G \mid H \geq N\}$ (subgroup of G containing N), $E_2 = \{K \leq G/N\}$ (subgroups of the quotient group).

Further, let $\pi : G \rightarrow G/N$ be the canonical projection homomorphism.

Then E_1 and E_2 are in bijection by

$$\begin{aligned} \Phi : E_1 &\rightarrow E_2 \text{ given by } \Phi(H) = \pi(H) \\ \Psi : E_2 &\rightarrow E_1 \text{ given by } \Psi(K) = \pi^{-1}(K) \end{aligned}$$

1. Φ and Ψ preserve inclusion (i.e. $H_1 \leq H_2$ in E_1 implies $H_1/N \leq H_2/N$, and vice versa).
2. Φ and Ψ preserve normality (i.e. $H \trianglelefteq G$ in E_1 implies $H/N \trianglelefteq G/N$).
3. For any $H \in E_1$ that is normal in G , $G/H \cong (G/N)/(H/N)$ as groups.

General Fact

If $\phi : G_1 \rightarrow G_2$ is any homomorphism of groups, then
$$\begin{cases} H_1 \leq G_1 \implies \phi(H_1) \leq G_2 \\ H_2 \leq G_2 \implies \phi^{-1}(H_2) \leq G_1 \end{cases}.$$

So Φ and Ψ are well-defined.

• Proof of Second Fact

1. $1_{G_1} \in \phi^{-1}(H_2)$, since $\phi(1_{G_1}) = 1_{G_2} \in H$.
2. If $g_1, g_1' \in \phi^{-1}(H_2)$, then $\phi(g_1 g_1') = \phi(g_1) \phi(g_1') \in H_2$ since H_2 is closed under product.
3. If $g_1 \in \phi^{-1}(H_2)$, then $g_1^{-1} \in \phi^{-1}(H_2)$ similarly.

Proof of 1

Say $H_1, H_2 \in E_1$ such that $H_1 \leq H_2$.

Then $\Phi(H_1) = \{h_1 N \mid h_1 \in H_1\} \subseteq \{h_2 N \mid h_2 \in H_2\} = \Phi(H_2)$.

Similarly, $K_1 \leq K_2 \in H_2$ implies $\Psi(K_1) \leq \Psi(K_2)$.

Proof of 2

Suppose $H \in E_1$ and $H \trianglelefteq G$. We want to prove that $\Phi(H) = H/N$ is normal in G/N .

So let gN be an arbitrary element in G/N , then $\forall h \in H$ we have

$$(gN)(hN)(gN)^{-1} = ghg^{-1}N \leq H/N = \Phi(H)$$

Let $h \in H$ vary, and we get

$$(gN)(H/N)(gN)^{-1} \subseteq H/N$$

Run the argument with g replaced by g^{-1} ,

$$(g^{-1}N)(H/N)(g^{-1}N)^{-1} = (gN)^{-1}(H/N)(gN) \subseteq H/N$$

hence

$$H/N \subseteq (gN)(H/N)(gN)^{-1}.$$

Therefore, gN normalizes $H/N = \Phi(H)$.

Proof of 3


1.

Start with $G \xrightarrow{\pi'} G/H$. As a canonical projection, π' is surjective.

That is $g \mapsto \pi'(g) = gH$, and $\pi'(N) = \{1_{G/H}\}$ since for all $n \in N$, $n \in H$ so $nH = H$.

1.

By the Universal Mapping Property of G/N , we get a unique homomorphism



$$gN \longmapsto gH \quad \text{where } \pi'' \text{ is surjective.}$$

- Recall: 0th Isomorphism Theorem

If $\phi : G_1 \rightarrow G_2$ is a surjective homomorphism of groups, then $\bar{\phi} : G_1 / \ker(\phi) \xrightarrow{\sim} G_2$ is an isomorphism. Apply this to $\pi'' = \phi$.

$$\begin{aligned} \ker(\pi'') &= \{gN \in G/N \mid gH = H\} \iff g \in H \\ &= \{hN \mid h \in H\} \\ &= H/N \end{aligned}$$


Therefore $(G/N)/(H/N) \xrightarrow{\sim} G/H$.

Example 1


Let $G = \text{Heis}_3(\mathbb{R})$ and $N = Z(G) = D(G)$. Some interesting subgroups of $(\mathbb{R}^2, +)$ are

1. $\{0\}$

2. $\vec{v} \neq \vec{0} \rightsquigarrow \mathbb{R}\vec{v}$ and $\mathbb{Z}\vec{v}$ are subgroups.



3. Say \vec{v}, \vec{w} are linear independent. $\mathbb{Z}\vec{v} + \mathbb{R}\vec{w}$.



4. IMAGE HERE - LEGIT NO IDEA. Collection of vertical lines in \mathbb{R}^3 that form inverse of example 2?

October 31, 2023

Definition: Subnormal Series

Let G be a group.

A subnormal series (or filtration) of G is a series

$$G = G_0 \supseteq G_1 \supseteq G_2 \supseteq \cdots \supseteq G_l = \{1\}$$

of subgroups of G , such that $G_i \supseteq G_{i+1}$ for all $i = 0, \dots, l-1$.

The length of the filtration is l .

Definition: Normal Series

If each G_i is normal in G , we say the series is normal.

Definition: Factor Group

The quotient group G_i/G_{i+1} is called the i th factor group of the filtration.

This is often denoted $\text{gr}_i(G.)$ meaning “graded.”

Definition: Composition Series

If, for each $i = 0, \dots, l-1$, $\text{gr}_i(G.)$ is simple, then we say the filtration is a composition series (or a Jordan-Hölder series).

Example 1

Let $G = \text{Alt}(4)$.

Take $G_1 = D(G) = \text{Stab}_G(\text{String})$. (Recall $|G_1| = 4$.)

Take $G_2 = \{1\}$.

Then $G = G_0 = G_1 \supseteq G_2$ is a filtration.

It is a normal series.

G_0/G_1 is a cyclic group of order 3 and, hence, simple.

$G_1/G_2 = G_1$ (strictly isomorphic; this is an abuse of notation) has order 4, is abelian, and contains $\langle (B\ P)(W\ Y) \rangle$ as a proper subgroup. So $G.$ is not a JH series (composition series).

Example 2

Let $G = \text{Alt}(4)$, $H_1 = G_1 = D(G)$, $H_2 = \langle (B\ P)(W\ Y) \rangle$ and $H_3 = \{1\}$.

Then,

$$G = H_0 \supseteq H_1 \supseteq H_2 \supseteq H_3$$

is a subnormal series.

It is not a normal series, since H_2 is not normal in G .

It is a JH series, since H_0/H_1 , H_1/H_2 , and H_2/H_3 have orders 3, 2 and 2 respectively and, therefore, are all simple.

Proposition:

Every finite group G has at least one Jordan-Hölder Series.

Proof

By induction on $|G|$.

If $|G| = 1$, then, by convention, (G) is a JH series of length 0.

Suppose $|G| > 1$.

If G is simple (i.e. G has exactly 2 normal subgroups, namely $\{1\}$ and G), then

$$(G = G_0 \supseteq \{1\} = G_1)$$

is a JH series of length 1.

Suppose G is not simple.

Then G contains a normal subgroup N , $N \neq \{1\}$ and $G \neq N$.

Among all such nontrivial proper normal $N \trianglelefteq G$, choose the one with maximal $|N|$.

Then $Q := G/N$ is simple since, by the 2nd isomorphism theorem, if Q were not simple, it would contain a nontrivial proper normal subgroup $Q_1 \triangleleft Q$, which would correspond to a normal subgroup $N \triangleleft N_1 \triangleleft G$, but then $|N_1| > |N|$, which would contradict the choice of N .

By induction hypothesis, $|N| < |G|$, we have a JH series for N , say

$$N = N_0 \supseteq N_1 \supseteq \cdots \supseteq N_l = \{1\}$$

Then take $G_0 = G$ and $G_{i+1} = N_i$ for $i = 0, 1, 2, \dots, l$ to get


$$G = \underbrace{G_0 \supseteq N_0}_{Q} \supseteq N_1 \supseteq \cdots \supseteq N_l = \{1\}$$

which have simple factors.

Therefore, N is a JH series. ■

Example 1

For $G = \text{Alt}(4)$,



has 3 JH series.


Example 2

For $G = \mathbb{Z}/12\mathbb{Z}$.

Fact: by the 2nd isomorphism theorem, subgroups of G are exactly $a\mathbb{Z}/12\mathbb{Z}$ where $a \mid 12$.

(\iff subgroup $a\mathbb{Z}$ of \mathbb{Z} containing $12\mathbb{Z}$)

12 has factors 1,2,3,4,6,12. Then



there are 3 JH series.

Definition: Product Group

Let G and H be groups.

The product group $G \times H$ is the cartesian product equipped with the component wise group operation.

$$(g_1, h_1) * (g_2, h_2) = (g_1 * g_2, h_1 * h_2)$$

Example

Let p be a prime number.

Take $G = H = \mathbb{Z}/p\mathbb{Z}$. Then

$$(\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/p\mathbb{Z}) \underset{\text{notation}}{=} (\mathbb{Z}/p\mathbb{Z})^2$$

where $|(\mathbb{Z}/p\mathbb{Z})^2| = p^2$ and

$$(a, b) + (a', b') = (a + a', b + b')$$

for $a, a', b, b' \in \mathbb{Z}$.

A subgroup K of $\tilde{G} = (\mathbb{Z}/p\mathbb{Z})^2$, by Lagrange's Theorem, must have order 1, p or p^2 .

We know $|\{1\}| = 1$ and $|\tilde{G}| = p^2$.

If $|K| = p$, then K is cyclic and generated by (a, b) .

If $k \in \{1, 2, \dots, p-1\}$, then note that for $(a, b) \neq (0, 0)$

$$\langle (a, b) \rangle = \langle (ka, kb) \rangle$$

Because both have order p .

Conversely, if $(a, b) \neq (0, 0)$ and $(a', b') \neq (0, 0)$ satisfy

$$\langle (a, b) \rangle = \langle (a', b') \rangle$$


then for some $k \in \{1, 2, \dots, p-1\}$ we have $(a', b') = (ka, kb)$.

- Upshot

There are exactly $\frac{p^2-1}{p-1} = p+1$ subgroups K of order p in \tilde{G} .

– Example

For $p = 3$,



there are 4 JH series.

Definition: Automorphism

Let G be a group.

A (group) automorphism of G is a bijective group homomorphism

$$\phi : G \rightarrow G$$

$\text{Aut}(G) = \{\phi : G \rightarrow G \mid \text{automorphisms}\}.$

Fact: $\text{Aut}(G)$ equipped with composition (of functions) is a group.

Justification: $\phi \circ \psi \in \text{Aut}(G)$ if $\phi, \psi \in \text{Aut}(G)$; $\phi^{-1} = \phi^{-1}$ and $1_{\text{Aut}(G)} = \text{Id}_G$.

Example 1

Let $G = (\mathbb{Z}, +)$.

A homomorphism $G \xrightarrow{\phi} G$ is completely determined by $\phi(1) = a_\phi$.

Since $\phi(n) \underset{\phi \text{ homomorphism}}{=} n\phi(1) = na_\phi$.

Thus $\{\text{group homomorphisms } \phi : G \rightarrow G\} \leftrightarrow (\mathbb{Z}, \times)$ and $\phi \leftrightarrow a_\phi$.

If $(\phi(1) = a_\phi$ and $\psi(1) = a_\psi$, then note that

$$(\phi \circ \psi)(1) = \phi(\psi(1)) = \phi(a_\psi) = a_\phi a_\psi$$

So if $\phi \in \text{Aut}(G)$, then $a_\phi \in \mathbb{Z}$ must have a multiplicative inverse in \mathbb{Z} .

Hence $a_\phi = 1$ or $a_\phi = -1$.

Conversely $\begin{cases} \phi_1(n) = n \\ \phi_{-1}(n) = -n \end{cases}, \forall n \in \mathbb{Z}$ are in $\text{Aut}(G)$.

Therefore $\text{Aut}((\mathbb{Z}, +)) = \{\pm 1, x \setminus\}$.

Example 2

Let p be a prime and $G = (\mathbb{Z}/p\mathbb{Z}, +)$.

A group homomorphism $\phi : G \rightarrow G$ is (again) determined by $\phi(1) = a_\phi$.

So $\{\text{group homomorphisms } \phi : G \rightarrow G\} \leftrightarrow \mathbb{Z}/p\mathbb{Z}$, $\phi \leftrightarrow a_\phi$ and $\phi \circ \psi \leftrightarrow a_\phi a_\psi$.

So if ϕ is bijective, then a_ϕ must have a multiplicative inverse in $\mathbb{Z}/p\mathbb{Z}$.

Fact: Any nonzero element $a \in \mathbb{Z}/p\mathbb{Z}$ has a multiplicative inverse.

• Proof

Look at $\langle a \rangle = \{ka \mid k \in \mathbb{Z}\}$.

Since $\mathbb{Z}/p\mathbb{Z}$ is simple, $\langle a \rangle = \mathbb{Z}/p\mathbb{Z}$.

Hence $\langle a \rangle \ni 1$, so $\exists k \in \mathbb{Z}$ such that $ka = 1$ in $\mathbb{Z}/p\mathbb{Z}$.

Take $b := k \pmod{p}$, and we have $ba = 1$. ■

Conclusion

$$\text{Aut}((\mathbb{Z}/p\mathbb{Z}, +)) = (\mathbb{Z}/p\mathbb{Z})^x = ((\mathbb{Z}/p\mathbb{Z}) \setminus \{0\}, \cdot)$$

Example

Take $G = (\mathbb{Z}/p\mathbb{Z})^2$.

Compute $|\text{Aut}(G)|$.

$\text{Aut}(G)$ is also called $\text{GL}_2(\mathbb{F}_p)$.

Definition: Semidirect Product

Let H and N be groups and

$$\psi : H \rightarrow \text{Aut}(N)$$

be a group homomorphism.

The semidirect product $H \ltimes_{\psi} N$, as a set is (simply) the cartesian product $H \times N$ equipped with the binary operation

$$(h_1, n_1) * (h_2, n_2) = \left(h_1 h_2, n_1 \overbrace{(\underbrace{\psi(h_1)}_{\in \text{Aut}(N)} n_2)}^{\in N} \right)$$

This is a group.

$(1_H, 1_N)$ is the identity.