

# Random Matrix Theory

April 1, 2025

## Preliminaries

Let  $\xi_{ij}, \eta_{ij}$  be normal random variables (i.e. Gaussian, mean 0, variance 1).

e.g.  $\mathbb{P}(\xi_{11} < s) = \int_{-\infty}^s \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$ .

$\int_{-\infty}^{\infty} x^2 \cdot \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$  is the variance.

$\frac{1}{\sqrt{2\pi}} e^{-x^2/2}$  is the Probability Density Function (PDF).

$\frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$  is the probability measure on our probability space (i.e. totally finite measure space).

We build matrices

$$\begin{bmatrix} \xi_{11} & \frac{\xi_{12} + i\eta_{12}}{\sqrt{2}} & \frac{\xi_{13} + i\eta_{13}}{\sqrt{2}} & \dots \\ \frac{\xi_{21} + i\eta_{21}}{\sqrt{2}} & \xi_{22} & \frac{\xi_{22} + i\eta_{22}}{\sqrt{2}} & \\ \frac{\xi_{31} + i\eta_{31}}{\sqrt{2}} & \frac{\xi_{32} + i\eta_{32}}{\sqrt{2}} & \xi_{33} & \\ \vdots & & & \ddots \end{bmatrix}$$

## Homework

Is the PDF of  $\frac{a+b}{2}$  the same as  $\frac{\xi_{12}}{\sqrt{2}}$  for normal RVs  $a, b, \xi_{12}$ ?

i.e.  $\mathbb{P}\left(\frac{a+b}{2} < s\right) \stackrel{?}{=} \left(\mathbb{P}\frac{\xi_{12}}{\sqrt{2}}\right)$

## 2x2 Random Matrix

Our matrix  $L$  corresponds to eigenvalues  $\lambda_1, \lambda_2$  which are random variables determined by  $\{\xi_{ij}, \eta_{ij}\}$ .

Then the number of evaluations in the interval  $B$  is given by  $\sum_{j=1}^2 \chi_B(\lambda_j)$ . We may take the average by

$$\int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \sum_{j=1}^2 \chi_B(\lambda_j) \frac{1}{\sqrt{2\pi}} e^{-\xi_{11}^2} \cdot \frac{1}{\sqrt{2\pi}} e^{-\xi_{22}^2} \cdot \frac{1}{\sqrt{2\pi}} e^{-\xi_{12}^2} \cdot \frac{1}{\sqrt{2\pi}} e^{-\eta_{12}^2} d\xi_{11} d\xi_{22} d\xi_{12} d\eta_{12}.$$

## Expected Evaluations

We have that the expectation of the number of evaluations in the interval  $(a, b)$  is given by  $\int_a^b G(s) ds$  where

$$G(s) = e^{-\frac{s^2}{2}} \sum_{\ell=0}^2 P_{\ell}(s)^2$$

and  $P_{\ell}(s)$  is the Hermite polynomial of degree  $d$ .