

# Algebra III

April 1, 2024

## Chapter 0: Review

### Definition: Category

A category  $\mathcal{C}$  consists of the following data:

1. A class of objects,  $\text{Obj}(\mathcal{C})$ .
2. For any pair of objects  $X, Y \in \text{Obj}(\mathcal{C})$ , a set of morphisms  $\text{Mor}_{\mathcal{C}}(X, Y)$ ,  $\text{Hom}_{\mathcal{C}}(X, Y)$  or  $\mathcal{C}(X, Y)$ .
3. For any triple of objects  $X, Y, Z \in \text{Obj}(\mathcal{C})$ , a map

$$\begin{aligned}\text{Hom}_{\mathcal{C}}(Y, Z) \times \text{Hom}_{\mathcal{C}}(X, Y) &\rightarrow \text{Hom}_{\mathcal{C}}(X, Z) \\ (g, f) &\mapsto g \circ f\end{aligned}$$

called compositions subject to the following axioms:

1. Associativity:  $f \circ (g \circ h) = (f \circ g) \circ h$  whenever this makes sense.
2. For every object  $X \in \text{Obj}(\mathcal{C})$ , there exists a morphism  $\text{id}_X \in \text{Hom}_{\mathcal{C}}(X, X)$  such that

$$\text{id}_X \circ f = f \quad \text{and} \quad g \circ \text{id}_X = g, \quad \forall f \in \text{Hom}_{\mathcal{C}}(W, X), g \in \text{Hom}_{\mathcal{C}}(X, W)$$

### Example 1

Let  $E$  be a set (or a class).

Define  $\mathcal{C}$  by taking  $\text{Obj}(\mathcal{C}) = E$  and  $\text{Hom}_{\mathcal{C}}(X, Y) = \begin{cases} \emptyset & \text{if } x \neq y \\ \{\text{id}_X\} & \text{if } x = y \end{cases}$ .

### Example 2

Let  $\mathcal{C} = \text{Set}$  the category of all sets with set functions acting as morphisms.

Let  $\mathcal{C} = \text{Grp}$  the category of all groups with group homomorphisms acting as morphisms.

Abelian Rings: Ab, Rings: Ring, Commutative Rings: CRing, Vector Spaces over  $F$ :  $\text{Vect}_F$ , Topological Spaces: Top, etc.

### Example 3

Let  $G$  be a group (or more generally a monoid).

Define  $\text{Obj}(\mathcal{C}) = \{*\}$ ,  $\text{Hom}_{\mathcal{C}}(*, *) = G$  and

$$\text{Hom}_{\mathcal{C}}(*, *) \times \text{Hom}_{\mathcal{C}}(*, *) \rightarrow \text{Hom}_{\mathcal{C}}(*, *)$$

the group operator.

#### Example 4

Let  $(E, \leq)$  be a preordered set (i.e. reflexive and transitive).  
Define  $\mathcal{C}$  by  $\text{Obj}(\mathcal{C}) = E$ ,

$$\text{Hom}_{\mathcal{C}}(x, y) = \begin{cases} \emptyset & \text{if } x \not\leq y \\ \{f_{xy}\} & \text{if } x \leq y \end{cases}$$

#### Notation

If  $f \in \text{Hom}_{\mathcal{C}}(X, Y)$  we write  $X \xrightarrow{f} Y$  in  $\mathcal{C}$ .

#### Definition: Isomorphism

A morphism  $f : X \rightarrow Y$  in  $\mathcal{C}$  is an isomorphism if  $\exists g : Y \rightarrow X$  such that  $g \circ f = \text{id}_X$  and  $f \circ g = \text{id}_Y$ .

#### Definition: Endomorphism

A morphism on  $X$  with  $f : X \rightarrow X$ .

#### Definition: Automorphism

An automorphism on  $X$  is just an isomorphism  $f : X \xrightarrow{\sim} X$  from  $X$  to itself.  
Note that  $\text{Aut}_{\mathcal{C}}(X) \subseteq \text{End}_{\mathcal{C}}(X) = \text{Hom}_{\mathcal{C}}(X, X)$ .

#### Remark:

The collection of all endomorphisms on  $X$  form a monoid.  
The collection of all automorphisms on  $X$  forms a group called the automorphism group of  $X$ .

#### Example 1

Let  $\mathcal{C} = \text{Set}$ ,  $X = \{1, \dots, n\}$ . Then  $\text{Aut}_{\text{Set}}(\{1, \dots, n\}) = \text{Perm}(X) = S_n$ .

#### Example 2

Let  $\mathcal{C} = \text{Vect}_F$ ,  $X = F^n$ . Then  $\text{Aut}_{\text{Vect}_F}(F^n) = \text{GL}_n(F)$ .

#### Definition: Functors

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories.  
A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  from  $\mathcal{C}$  to  $\mathcal{D}$  consists of the following data

1. For each object  $X \in \text{Obj}(\mathcal{C})$ , a chosen object  $F(X) \in \text{Obj}(\mathcal{D})$ .
2. For each pair of objects  $X, Y \in \text{Obj}(\mathcal{C})$ , a function

$$\begin{aligned} \text{Hom}_{\mathcal{C}}(X, Y) &\rightarrow \text{Hom}_{\mathcal{D}}(F(X), F(Y)) \\ f &\mapsto F(f) \end{aligned}$$

such that

1. For any two composable morphisms  $X \xrightarrow{f} Y \xrightarrow{g} Z$  in  $\mathcal{C}$ , we have  $F(g \circ f) = F(g) \circ F(f)$ .
2. For each object  $X \in \text{Obj}(\mathcal{C})$ ,  $F(\text{id}_X) = \text{id}_{F(X)}$ .

### Example 1

For  $\mathcal{D} := \mathcal{C}$ ,  $\text{Id} : \mathcal{C} \rightarrow \mathcal{C}$ ,  $X \mapsto X$ ,  $f \mapsto f$ .

### Example 2: Forgetful Functors

$\mathcal{U} : \text{Grp} \rightarrow \text{Set}$  given as  $(G, \cdot) \mapsto G$ .

$\text{Ring} \rightarrow \text{Ab}$  given as  $(R, +, \cdot) \mapsto (R, +)$ .

### Example 3: Tensors

Let  $R$  be a commutative ring,  $M \in \text{Mod}_R$ .

Then  $\otimes_R M : \text{Mod}_R \rightarrow \text{Mod}_R$  and  $\text{Hom}_R(M, -) : \text{Mod}_R \rightarrow \text{Mod}_R$ .

### Definition:

Let  $X$  be an object in a category  $\mathcal{C}$  and  $G$  a group.

An action of  $G$  on  $X$  is a group homomorphism  $G \rightarrow \text{Aut}_{\mathcal{C}}(X)$ .

### Example 1

Let  $\mathcal{C} = \text{Set}$ .

A  $G$ -set is a set  $X \in \text{Set}$  equipped with a group homomorphism

$$G \rightarrow \text{Perm}(X) = \text{Aut}_{\text{Set}}(X)$$

### Exercise 1

A  $G$ -set is the same thing as a functor  $G \rightarrow \text{Set}$ ,  $* \mapsto X$ ,  $\text{Hom}_{\mathcal{C}}(*, *) \rightarrow \text{Hom}_{\text{Set}}(X, X)$  ( $G \rightarrow \text{Aut}_{\text{Set}}(X)$ ).

### Definition: Adjunctions

Let  $\mathcal{C}$  and  $\mathcal{D}$  be categories and  $F : \mathcal{C} \rightarrow \mathcal{D}$  and  $G : \mathcal{D} \rightarrow \mathcal{C}$  be functors.

We say that  $F$  is left adjoint to  $G$  (and that  $G$  is right adjoint to  $F$ , and that we have a pair of adjoint functors) if for each object  $X \in \text{Obj}(\mathcal{C})$  and  $Y \in \text{Obj}(\mathcal{D})$ , we have a bijection

$$\text{Hom}_{\mathcal{D}}(F(X), Y) \xrightarrow{\sim} \text{Hom}_{\mathcal{C}}(X, G(Y))$$

which is “natural in  $X$  and  $Y$ ”:

For any  $f : X \rightarrow X'$  in  $\mathcal{C}$ ,

$$\begin{array}{ccc} \text{Hom}_{\mathcal{D}}(F(X'), Y) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{C}}(X', G(Y)) \\ \downarrow - \circ F(f) & & \downarrow - \circ f \\ \text{Hom}_{\mathcal{D}}(F(X), Y) & \xrightarrow{\sim} & \text{Hom}_{\mathcal{C}}(X, G(Y)) \end{array}$$

and for every  $g : Y \rightarrow Y'$  in  $\mathcal{D}$

$$\begin{array}{ccc}
\mathrm{Hom}_{\mathcal{D}}(F(X), Y) & \xrightarrow{\sim} & \mathrm{Hom}_{\mathcal{C}}(X, G(Y)) \\
\downarrow -\circ F(f) & & \downarrow -\circ f \\
\mathrm{Hom}_{\mathcal{D}}(F(X), Y') & \xrightarrow{\sim} & \mathrm{Hom}_{\mathcal{C}}(X, G(Y'))
\end{array}$$

We write

$$\begin{array}{ccc}
& \mathcal{C} & \\
F \downarrow & \uparrow & G \\
& \mathcal{D} &
\end{array}$$

### Example 1

For  $M \in \mathrm{Mod}_R$  we have

$$\begin{array}{ccc}
& \mathrm{Mod}_R & \\
-\otimes_R M \downarrow & \uparrow & \mathrm{Hom}_R(M, -) \\
& \mathrm{Mod}_R &
\end{array}$$

where

$$\begin{aligned}
\mathrm{Hom}_R(M_1 \otimes M_2, N) &\cong \mathrm{Hom}_R(M_1, \mathrm{Hom}_R(M, \mathrm{Hom}_R(M_2, N))) \\
f &\mapsto (x \mapsto (y \mapsto f(x \otimes y)))
\end{aligned}$$

### Example 2

Let  $R \xrightarrow{\phi} S$  be a ring homomorphism.

We can regard an  $S$ -module  $N$  as an  $R$ -module via

$$r \cdot x := \phi(r)x, \quad \forall r \in R, x \in N$$

This defines a functor  $\mathrm{Mod}_S \rightarrow \mathrm{Mod}_R$  called a “restriction of scalars”, which has a left adjoint called “extension of scalars.”

$$\begin{array}{ccc}
& \mathrm{Mod}_R & \\
S \otimes_R - \downarrow & \uparrow & \\
& \mathrm{Mod}_R &
\end{array}$$

### Recall

For commutative ring  $R$ ,  $\sim \mathrm{Mod}_R$ .

e.g.  $R = F$  a field,  $\mathrm{Mod}_R \equiv \mathrm{Vect}_F$ ;  $R = \mathbb{Z}$ ,  $\mathrm{Mod}_R \equiv \mathrm{Ab}$ .

### Definition: R-Algebra

An  $R$ -algebra is an Abelian group  $(A, +)$  that has both the structure of

1. an  $R$ -module and
2. a ring

which are compatible in that

$$r(ab) = (ra)b = a(rb), \quad \forall r \in R, a, b \in A$$

### Example 1

The polynomial ring  $R[x]$  is an  $R$ -algebra.

### Example 2

The ring of  $n \times n$  matrices  $M_n(R)$  is an  $R$ -algebra.

### Example 3

If  $R \xrightarrow{\phi} S$  is a homomorphism of commutative rings, then  $S$  is an  $R$ -algebra via  $r := \phi(r)a$ ,  $\forall r \in R, a \in S$ .

### Example 4

$\mathbb{R} \hookrightarrow \mathbb{C}$ . So  $\mathbb{C}$  is an  $\mathbb{R}$ -algebra.

$R \hookrightarrow R[x]$ .

More generally,  $R[x_1, x_2, \dots, x_n]$  is an  $R$ -algebra.

### Commutative R-Algebras

An  $R$ -algebra is commutative if it is commutative as a ring.

$\text{CAlg}_R \subset \text{Alg}_R$ .

### Question: Why are polynomials important?

An algebraic perspective: they are the “free commutative algebras.”

### Recall

For  $R$  a commutative ring, we have the notion of a free  $R$ -module – one that admits a basis.

Categorically, we have an adjunction.

$$\begin{array}{ccc}
& \text{Set} & \\
& \downarrow f & \uparrow \mathcal{U} \\
& \text{Mod}_R &
\end{array}$$

The left adjoint of the forgetful functor sends a set  $I$  to the free  $R$ -module with basis  $I$ .

$$F(I) = R^{(I)} = \bigoplus_{i \in I} R$$

The adjunction says that for any set  $I$  and  $R$ -module  $M$ ,

$$\begin{aligned}
& \text{Hom}_{\text{Mod}_R}(R^{(I)}, M) \xrightarrow{\sim} \text{Hom}_{\text{Set}}(I, M) \\
& \exists! R\text{-linear map } f: R^{(I)} \rightarrow M \quad \leftarrow \{x_i\}_{i \in I} \\
& \quad e_i \mapsto x_i
\end{aligned}$$

Similarly, the forgetful functor  $\mathcal{U} : \text{CAlg}_R \rightarrow \text{Set}$  has a left adjoint

$$\begin{array}{ccc}
& \text{Set} & \\
& \downarrow f & \uparrow \mathcal{U} \\
& \text{CAlg}_R &
\end{array}$$

which sends a set  $I$  to the “free commutative  $R$ -algebra on  $I$ .”

Explicitly,  $F(I) = R[\{x_i\}_{i \in I}]$  the polynomial algebra with an indeterminate  $x_i$  for each  $i \in I$ .

## Example 1

$$I = \{*\} \rightsquigarrow F(\{*\}) = R[x].$$

$$I = \{1, \dots, n\} \rightsquigarrow F(\{1, \dots, n\}) = R[x_1, \dots, x_n].$$

$$I = \mathbb{N} \rightsquigarrow F(\mathbb{N}) = R[x_1, x_2, \dots].$$

## Adjunction

For any set  $I$  and commutative  $R$ -algebra  $A \in \mathbf{CAlg}_R$ , we have a bijection

$$\begin{aligned} \text{Hom}_{\mathbf{CAlg}_R}(R[\{x_i\}_{i \in I}], A) &\cong \text{Hom}_{\mathbf{Set}}(I, A) \\ \exists! R\text{-algebra homomorphism } R[\{x_i\}_{i \in I}] &\xrightarrow{x_i \mapsto a_i} A \leftarrow \{a_i\}_{i \in I} \end{aligned}$$

## Example 1

Let  $A$  be a commutative  $R$ -algebra.

For any  $a \in A$ , there exists a unique  $R$ -algebra homomorphism  $R[x] \rightarrow A$  which sends  $X \mapsto a$ .

Explicitly,  $f(x) \mapsto f(a)$ .

## Corollary

Let  $R \xrightarrow{\phi} S$  be a homomorphism of commutative rings.

For any  $a \in S$ , there is a unique ring  $R[x] \xrightarrow{\bar{\phi}} S$  such that  $\bar{\phi}|_R = \phi$  and  $\bar{\phi}(X) = a$ .

## Example 1

Let  $R \subseteq S$  be a subring.

For each  $a \in S$ , there is a unique ring homomorphism  $R[x] \xrightarrow{\phi} S$  such that  $\phi|_R = \text{id}$  and  $\phi'(X) = a$ .

We call this the “evaluation at  $a$ .”

$$\begin{aligned} R[x] &\xrightarrow{\text{ev}_a} S \\ f &\mapsto f(a) \end{aligned}$$

## Definition: Subalgebra

Let  $A$  be a commutative  $R$ -algebra, and let  $S \subset A$  be a subset.

The subalgebra of  $A$  generated by  $S$ , denoted  $R[S]$ , is the intersection of all subalgebras of  $A$  which contain  $S$ .

Explicitly,

$$R[S] = \{a \in A : \exists n \geq 1, s_1, \dots, s_n \in S, f \in R[x_1, \dots, x_n], a = f(s_1, \dots, s_n)\}$$

## Example 1

Let  $A = R[x]$ . Then  $A = R[x]$ . That is,  $A$  is generated by  $\{x\}$  as an algebra.

Similarly,  $R[x_1, \dots, x_n]$  is generated as an algebra by  $\{x_1, \dots, x_n\}$ .

## Example 2

If  $R[x]/I$  with  $I \subset R[x]$  an ideal, and  $x := \bar{X} \in A$ , then  $A = R[x]$ . That is,  $A$  is generated by  $x = \bar{X}$  as an algebra.

More generally, if  $I \subset R[x_1, \dots, x_n]$  an ideal, then  $R[x_1, \dots, x_n]/I$  is generated by  $\{\bar{x}_1, \dots, \bar{x}_n\}$ .

## Proposition

If  $A \in \text{CAlg}_R$  is a finitely generated, commutative  $R$ -algebra, then  $A \cong R[x_1, \dots, x_n]/I$  for some  $n \geq 1$  and ideal  $I \subset R[x_1, \dots, x_n]$ .

**April 3, 2024**

## Definition: Symmetric Polynomials

Let  $R$  be a commutative ring.

A polynomial  $f \in R[x_1, \dots, x_n]$  is symmetric if  $f(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = f(x_1, \dots, x_n)$  for all  $\sigma \in S_n$ .

In more detail: the symmetric group  $S_n$  acts on  $R[x_1, \dots, x_n]$  by  $R$ -algebra homomorphism.

$\sigma \in S_n \curvearrowright R[x_1, \dots, x_n] \rightarrow R[x_1, \dots, x_n]$  given by  $x_i \mapsto x_{\sigma(i)}$ .

The canonical action of  $S_n$  on  $\{1, \dots, n\}$  is

$$\begin{aligned} S_n &\rightarrow \text{Set} \xrightarrow{F} \text{CAlg}_R \\ * &\mapsto \{1, \dots, n\} \mapsto R[x_1, \dots, x_n] \end{aligned}$$

### Exercise 1

The symmetric polynomials form a subalgebra of  $R[x_1, \dots, x_n]$ .

### Example 1

Consider the polynomial

$$(*) \quad (t - x_1)(t - x_2) \cdots (t - x_n) \in R[x_1, \dots, x_n][t]$$

Write

$$t^n - s_1 t^{n-1} + s_2 t^{n-2} + \cdots + (-1)^n s_n$$

where  $s_1, \dots, s_n \in R[x_1, \dots, x_n]$ .

### Examples

Let  $n = 2$ .

$$(t - x_1)(t - x_2) = t^2 - \underbrace{(x_1 + x_2)}_{s_1} t + \underbrace{x_1 x_2}_{s_2}$$

Let  $n = 3$ .

$$(t - x_1)(t - x_2)(t - x_3) = t^3 - \underbrace{(x_1 + x_2 + x_3)}_{s_1} t^2 + \underbrace{(x_1 x_2 + x_2 x_3 + x_1 x_3)}_{s_2} t - \underbrace{x_1 x_2 x_3}_{s_3}$$

### Exercise 2

Show that the polynomials  $s_1, \dots, s_n \in R[x_1, \dots, x_n]$  are symmetric using the fact that  $(*)$  is unchanged by permuting the  $x_i$ s.

## Definition: Elementary Symmetric Polynomials

The polynomials  $s_1, \dots, s_n \in R[x_1, \dots, x_n]$  are the elementary symmetric polynomials in  $n$  variables. Explicitly,

$$\begin{aligned} s_1 &= x_1 + x_2 + \dots + x_n \\ s_2 &= \sum_{1 \leq i < j \leq n} x_i x_j \\ &\vdots \\ s_k &= \sum_{1 \leq i_1 < \dots < i_k \leq n} x_{i_1} x_{i_2} \dots x_{i_k} \\ &\vdots \\ s_n &= x_1 x_2 \dots x_n \end{aligned}$$

## Theorem: Fundamental Theorem on Symmetric Polynomials

Every symmetric polynomial  $f \in R[x_1, \dots, x_n]$  can be expressed in a unique way as a polynomial in the elementary symmetric polynomials.

In particular,  $R[s_1, \dots, s_n] \subseteq R[x_1, \dots, x_n]$  is the subalgebra of symmetric polynomials.

## Recall: Group of Units

If  $R$  is a ring, then  $U(R) = R^\times = \{a \in R : a \text{ is invertible}\}$ .

This is the multiplicative group of units in  $R$ .

### Exercise 3

This determines a functor  $\text{Ring} \rightarrow \text{Grp}$ .

## Definition: Field

A field is a nonzero commutative ring  $F$  in which every nonzero element is invertible (i.e.  $F^\times = F \setminus \{0\}$ ).

### Remarks:

A field has no nontrivial ideals.

A commutative ring  $R$  is a field if and only if  $(0)$  is a maximal ideal.

If  $I \subset R$  is an ideal in a commutative ring then  $R/I$  is a field if and only if  $I$  is a maximal ideal.

## Definition: Domain

A (integral) domain is a nonzero commutative ring  $R$  such that  $\forall a, b \in R, ab = 0 \implies a = 0 \text{ or } b = 0$ .

### Remarks:

A commutative ring  $R$  is a domain if and only if  $(0)$  is a prime ideal.

If  $I \subset R$  is an ideal in a commutative ring, then  $R/I$  is a domain if and only if  $I$  is a prime ideal.

## Example 1

Every field is a domain.



In fact, every subring of a field is a domain.  
Conversely, domains can be characterized as the subrings of fields.

## Definition: Field of Fractions

Let  $R$  be a domain.  
Its field of fractions,  $\text{Frac}(R)$ , is the set of all “formal fractions”

$$\text{Frac}(R) = \left\{ \frac{a}{b} : a, b \in R, b \neq 0 \right\}$$

More precisely,  $\text{Frac}(R) = (R \times (R \setminus \{0\})) / \sim$  where

$$(a_1, b_1) \sim (a_2, b_2) \iff a_1 b_2 = a_2 b_1$$

and we define  $\frac{a}{b} := [(a, b)]$ .

It is a field under addition and multiplication of fractions

$$\frac{a_1}{b_1} + \frac{a_2}{b_2} = \frac{a_1 b_2 + a_2 b_1}{b_1 b_2} \quad \text{and} \quad \frac{a_1}{b_1} \frac{a_2}{b_2} = \frac{a_1 a_2}{b_1 b_2}$$

We have an injective ring homomorphism

$$\begin{aligned} R &\hookrightarrow \text{Frac}(R) \\ a &\mapsto \frac{a}{1} \end{aligned}$$

### Example 1

$$\mathbb{Q} = \text{Frac}(\mathbb{Z}).$$

### Remark:

$\mathbb{Z}$  is a domain.

Its ideals are  $n\mathbb{Z}$  for  $n = 0, 1, 2, \dots$

Its prime ideals are  $(0)$  and  $p\mathbb{Z}$  for  $p$  prime.

## Definition: Root

Let  $R$  be a commutative ring and  $f \in R[x]$ .

A root or zero of  $f$  is an element  $r \in R$  such that  $f(r) = 0$ .

$$\begin{aligned} R[x] &\xrightarrow{\text{ev}_a} R \\ f &\longmapsto 0 \end{aligned}$$

The kernel is  $(x - a)$ .

That is  $f(a) = 0$  if and only if  $f \in (x - a)$ , if and only if  $x - a \mid f$ , if and only if  $f(x) = (x - a)g(x)$  for some  $g \in R[x]$ .

## Proposition:

Let  $R$  be a domain. Then

1.  $R[x]$  is a domain.
2.  $\deg(fg) = \deg(f) + \deg(g)$ .
3.  $R[x]^\times = R^\times$  (i.e.  $f \in R[x]^\times \iff f(x) = b_0$  with  $b_0 \in R^\times$ ).

## Example 1

If  $R = F$  a field,  $F[x]^\times =$  the nonzero constant polynomials.

## Remark:

If  $R$  a domain and  $a \in R$  a root of  $f \in R[x]$ , then

$$f(x) = (x - a)^m g(x)$$

with  $g(a) \neq 0$ . The  $m$  is uniquely determined and called the multiplicity of the root. Roots of multiplicity 1 are called simple roots.

## Remark:

If  $R$  is a domain, a polynomial  $f \in R[x]$  of degree  $d$  has at most  $d$  roots. In fact, at most  $d$  roots counted with multiplicity.

## Definition: Formal Derivative

The formal derivative of a polynomial

$$f(x) = a_d x^d + a_{d-1} x^{d-1} + \cdots + a_2 x^2 + a_1 x + a_0 \in R[x]$$

is the polynomial  $Df = f' \in R[x]$  defined by

$$f'(x) = d a_d x^{d-1} + \cdots + 2 a_2 x + a_1$$

## Remark: Properties

$$\begin{aligned} (f + g)' &= f' + g' & R[x] \rightarrow R[x] \text{ is } R\text{-linear} \\ (af)' &= a f' & \text{for } a \in R \\ (fg)' &= f g' + f' g & \text{(Leibniz Formula)} \end{aligned}$$

## Proposition:

$a \in R$  is a multiple root of  $f \in R[x]$  if and only if  $f(a) = 0$  and  $f'(a) = 0$ .

## Proof

$$f(x) = (x-a)^m g(x), g(a) \neq 0.$$

Therefore, by Leibniz,  $f'(x) = m(x-a)^{m-1}g(x) + (x-a)^m g'(x)$ .

## Recall:

For a field  $F$ , the polynomial ring  $F[x]$  is a PID.

$\mathbb{Z}$  is also a PID.

## Proposition:

Let  $R$  be a PID.

Every nonzero prime ideal is maximal.

## Proof

Let  $0 \neq p$  be a nonzero prime ideal.

Suppose  $p \subseteq I$ . Then  $p = (p)$  and  $I = (a)$  for some  $a \in R$  and prime element  $p \in R$ .

Then  $(p) \subseteq (a)$  and  $p = ab$  for some  $b \in R$ . So  $p|a$  or  $p|b$ .

If  $p|a$ , then  $p = I$ . If, instead,  $b = pc$  for some  $c \in R$ , then

$$p = acp \implies 1 = ac \implies a \in R^\times \implies (a) = R$$

## Example 1

If  $f \in F[x]$  is an irreducible polynomial then  $F[x]/(f)$  is a field.

For example,  $R[x]/(x^2 + 1)$  is a field ( $\cong \mathbb{C}$ ).

Also,  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$  is a field.

## Example 2

On the other hand,

$$(\mathbb{Z}/n)^\times = \{a \in \mathbb{Z}/n : \gcd(a, n) = 1\}$$

$$|(\mathbb{Z}/n)^\times| = |\{0 \leq k \leq n-1 : \gcd(k, n) = 1\}| = \phi(n)$$

Euler's Totient Function.

## Remark

Later in the course, we will prove the Fundamental Theorem of Algebra which states that every nonconstant complex polynomial  $f \in \mathbb{C}[x]$  has a root.

This implies that if  $f \in \mathbb{C}[x]$  is a monic polynomial with complex coefficients then  $f(x) = (x - \alpha_1)(x - \alpha_2) \cdots (x - \alpha_n)$  with  $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ .

Write it as

$$f(x) = x^d + a_1 x^{d-1} + \cdots + a_n$$

with coefficients  $a_1, \dots, a_n \in \mathbb{C}$ . Then

$$a_1 = -s_1(\alpha_1, \dots, \alpha_n) = -(\alpha_1 + \dots + \alpha_n)$$

$$a_k = (-1)^k s_k(\alpha_1, \dots, \alpha_n)$$

$$a_n = (-1)^n \alpha_1 \cdots \alpha_n$$

### Example 1

$$f(x) = x^2 + bx + c = (x - \alpha_1)(x - \alpha_2)$$

where  $\alpha_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2}$  and  $\alpha_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2}$ .

So  $\alpha_1 + \alpha_2 = -b$  and  $\alpha_1 \alpha_2 = c$ .

### Bottom Line

The coefficients of a monic polynomial are very simple expressions of the roots of the polynomial.

### Motivating Question

Can we go the other around?

Can we find simple expressions of the roots of a polynomial in terms of the coefficients.

**April 8, 2024**

## Chapter 1: Field Theory

### Definition: Field Homomorphism

If  $F$  and  $K$  are fields, a field homomorphism  $F \xrightarrow{\phi} K$  is just a ring homomorphism.

#### Remark

The kernel of a field homomorphism  $\phi : F \rightarrow K$  is an ideal of  $F$ .

Hence, it is either  $(0)$  or  $F$ . Since  $\phi(1_F) = 1_K \neq 0$ ,  $\ker(\phi) = (0)$ .

Thus every field homomorphism is automatically injective and embeds  $F$  as a subfield of  $K$ .

#### Notation

If  $F \subseteq K$  is a subfield, we say that  $K$  is an extension of  $F$  or that  $K/F$  is a field extension.

#### Remark

The ring of integers  $\mathbb{Z}$  is the initial object in the category of rings.

That is, given any ring  $R$ , there is a unique ring homomorphism  $\mathbb{Z} \rightarrow R$  given by  $n \mapsto n1_R = \begin{cases} \overbrace{1_R + \dots + 1_R}^n & \text{if } n \geq 0 \\ -\underbrace{(1_R + \dots + 1_R)}_n & \text{if } n < 0 \end{cases}$ .

The kernel of an ideal of  $\mathbb{Z}$ . We have three possibilities

1.  $\ker = \mathbb{Z} \implies 1_R = 0 \text{ in } R \implies R = 0$ .

$$2. \ker = (0) \implies \mathbb{Z} \hookrightarrow R.$$

$$3. \ker = n\mathbb{Z} \text{ for some } n \geq 2 \implies \mathbb{Z}/n\mathbb{Z} \hookrightarrow R.$$

## Proposition

Let  $F$  be a field and consider the unique ring homomorphism  $\mathbb{Z} \xrightarrow{\phi} F$ . Then the kernel of  $\phi$  is either  $(0)$  or  $p\mathbb{Z}$  for some prime number  $p$ .

### Proof

Note that  $\mathbb{Z}/n\mathbb{Z} \hookrightarrow F$ , but all subrings of fields are domains and  $\mathbb{Z}/n\mathbb{Z}$  is a domain if and only if  $n\mathbb{Z}$  is a prime ideal of  $\mathbb{Z}$ .

### Corollary

Let  $F$  be a field. It contains precisely one of the following as a subfield

1.  $\mathbb{Q}$  or
2.  $\mathbb{F}_p$  for  $p$  prime.

### Proof

The proposition implies either  $\mathbb{Z} \hookrightarrow F$  or  $\mathbb{Z}/p\mathbb{Z} \hookrightarrow F$  for  $p$  prime.

If  $\mathbb{Z} \hookrightarrow F$  then this extends to an embedding  $\mathbb{Q} \hookrightarrow F$  by the universal property of the field of fractions.

On the other hand,  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$  by definition.

Claim: we can't have more than one such field as a subfield of  $F$ .

Observe that if  $F$  has a subfield isomorphic to  $\mathbb{F}_p$ , then  $p \cdot 1 = 0$  in  $F$ .

On the other hand, if  $\mathbb{Q} \subseteq F$  then  $p \cdot 1 \neq 0$  for all  $p$ .

Finally, if  $p \neq q$  primes and  $\mathbb{F}_p \subseteq F$  and  $\mathbb{F}_q \subseteq F$ , then  $p \cdot 1 = 0$  and  $q \cdot 1 = 0$  in  $F$ .

By Bezout's, this means that  $a, b \in \mathbb{Z} : ap + bq = 1$ . So

$$1 = 1 \cdot 1 = (ap + bq) \cdot 1 = (ap)1 + (bq)1 = a(p \cdot 1) + b(q \cdot 1) = 0 + 0 = 0$$

which cannot be true.

## Definition: Field Characteristic

We define the characteristic of a field  $F$  by  $\text{char}(F) = \begin{cases} 0 & \text{if } \mathbb{Q} \subseteq F \\ p & \text{if } \mathbb{F}_p \subseteq F \end{cases}$ .

### Remark

Note that the kernel of  $\mathbb{Z} \rightarrow F$  is  $\text{char}(F)\mathbb{Z} \subseteq \mathbb{Z}$ .

The characteristic of  $F$  is the smallest positive integer  $n$  such that  $n \cdot 1 = 0$  in  $F$  or 0 if  $n \cdot 1 \neq 0$  in  $F$  for all  $n \geq 1$ .

### Remark

If  $K/F$  is a field extension, then  $K$  and  $F$  have the same characteristic.

$n \cdot 1 = 0$  in  $F$  if and only if  $n \cdot 1 = 0$  in  $K$ . Observe that the composition  $\mathbb{Z} \rightarrow F \hookrightarrow K$  requires matching kernels.

## Aside

In math, one sometimes passes between characteristic zero and characteristic  $p$  through the integers.

$$\begin{array}{ccc} \mathbb{Z} & \hookrightarrow & \mathbb{Q} \\ \downarrow & & \\ \mathbb{F}_p & & \end{array}$$

## Examples

$\mathbb{Q}, \mathbb{R}, \mathbb{C}$  have characteristic 0.

$\mathbb{F}_p := \mathbb{Z}/p\mathbb{Z}$  has characteristic  $p$ .

$\mathbb{C}(t) := \text{Frac}(\mathbb{C}[t])$ .

$\mathbb{F}_p(t) := \text{Frac}(\mathbb{F}_p[t])$  is an infinite field of characteristic  $p$ .

## Remark

If  $R$  is a domain, then  $R[t]$  is a domain and

$$R(t) := \text{Frac}(R[t]) = \left\{ \frac{f}{g} : f, g \in R[t], g \neq 0 \right\}$$

the field of rational functions.

More generally,  $R[t_1, \dots, t_n]$  is a domain and

$$R(t_1, \dots, t_n) := \text{Frac}(R[t_1, \dots, t_n])$$

is the field of rational functions in  $n$  variables.

## Definition: Degree of a Field Extension

Let  $K/F$  be a field extension.

We can regard  $K$  as a vector space over  $F$  (restriction of scalars  $F \hookrightarrow K$ ).

The degree of the field extension  $K/F$  is  $\dim$  of the  $F$ -vector space  $K$ .

## Notation

$$[K : F] := \dim_F(K).$$

## Example

$\mathbb{C}/\mathbb{R}$  is a degree 2 extension. An  $\mathbb{R}$ -basis for  $\mathbb{C}$  is  $\{1, i\}$ .

## Remark

$K/F$  has degree 1 if and only if  $K = F$ .

## Terminology

A degree 2 extension  $K/F$  is a quadratic extension.

A degree 3 extension  $K/F$  is a cubic extension.

Etc.

## Definition: Finite Extension

A field extension  $K/F$  is said to be a finite extension if  $[K : F]$  is finite.

### Example

$F(t)/F$ , noting  $F \subseteq F[t] \subseteq F(t)$ , is an infinite extension. Write  $[F(t) : F] = \infty$ .

## Proposition

Let  $L/K/F$  be field extensions.

Then  $[L : F] = [L : K][K : F]$ .

### Proof (Sketch)

Idea: if  $\{a_i\}_{i \in I}$  is a basis for  $L/K$  and  $\{b_j\}_{j \in J}$  is a basis for  $K/F$ , then  $\{a_i b_j\}_{(i,j) \in I \times J}$  is a basis for  $L$  over  $F$ .  
Note that  $|I \times J| = |I||J|$ .

## Definition: Algebraic and Transcendental Elements

Let  $K/F$  be a field extension.

An element  $a \in K$  is said to be algebraic over  $F$  if it is a root of a nonzero polynomial with coefficients in  $F$ .

Otherwise, we say that  $a$  is transcendental over  $F$ .

### Example

Consider  $\mathbb{C}/\mathbb{Q}$ .

$\sqrt{2} \in \mathbb{C}$  is algebraic over  $\mathbb{Q}$  since  $t^2 - 2 \in \mathbb{Q}[t]$ .

$i \in \mathbb{C}$  is algebraic over  $\mathbb{Q}$  since  $t^2 + 1 \in \mathbb{Q}[t]$ .

$\omega_n = e^{2\pi i/n} \in \mathbb{C}$  is algebraic over  $\mathbb{Q}$  since  $t^n - 1 \in \mathbb{Q}[t]$ .

### Remark

Whether or not an element is algebraic or transcendental depends a lot on the ground field  $F$ .

e.g. every element  $a \in K$  is algebraic over  $K$ , since it is a root of  $t - a \in K[t]$ .

### Remark

Often the terms “algebraic number” and “transcendental number” mean a complex number which is algebraic or transcendental over  $\mathbb{Q}$ .

### Theorem: (Hermite 1873)

$e$  is a transcendental number.

### Theorem: (Lindemann 1882)

$\pi$  is a transcendental number.

### Exercise (Cantor)

There are only countably many algebraic numbers.

## Remark

Whether  $a \in K$  is algebraic or transcendental over  $F$  is described by the evaluation homomorphism

$$\begin{aligned} F[t] &\xrightarrow{\text{ev}_a} K \\ f &\longmapsto f(a) \end{aligned}$$

That is,  $a$  is transcendental if and only if  $\ker(\text{ev}_a) = (0)$ .

Then  $a$  is algebraic if and only if  $\ker(\text{ev}_a) \neq (0)$ .

$F[t]$  is PID, so if  $a \in K$  is algebraic over  $F$  then  $\ker(\text{ev}_a)$  is a nonzero principal ideal of  $F[t]$ .

A generator of this principal ideal is only determined up to association (that is up to multiplication by a nonzero constant polynomial).

We can pin it down by requiring the generator to be monic.

## Definition: Minimal Polynomial

The unique monic polynomial  $f$  of lowest degree with coefficients in  $F$  such that  $f(a) = 0$  is called the minimal polynomial of  $a$  over  $F$ .

## Notation

$$m_a(t) \in F[t].$$

## Remark

It generates  $\ker(\text{ev}_a)$ . For any  $f \in F[t]$ ,  $f(a) = 0$  if and only if  $m_a | f$ .

## Note

$F[t]/(m_a(t))$  is isomorphic to a subring of  $K$ .

So  $F[t]/(m_a(t))$  is a domain and  $m_a \in F[t]$  is an irreducible polynomial.

## Exercise

The minimal polynomial of  $a \in K$  over  $F$  is the unique, monic, irreducible polynomial  $f \in F[t]$  such that  $f(a) = 0$ .

## Example

Take  $\sqrt{2} \in \mathbb{C}$ , the root of  $t^2 - 2 \in \mathbb{Q}[t]$ . This is the minimal polynomial of  $\sqrt{2}$  over  $\mathbb{Q}$  since it is irreducible.

$i \in \mathbb{C}$  is a root of  $t^2 + 1 \in \mathbb{Q}[t]$  which is also irreducible and hence the minimal polynomial of  $i$  over  $\mathbb{Q}$ .

$a = \frac{1+i}{2} \in \mathbb{C}$  ( $\sqrt{i}$ ) is a root of  $t^4 + 1 \in \mathbb{Q}[t]$ , irreducible and therefore minimal of  $a$  over  $\mathbb{Q}$ .

Consider  $F = \mathbb{Q}[i] = \{\alpha + i\beta : \alpha, \beta \in \mathbb{Q}\}$ . Observe that  $t^4 + 1 = (t^2 - i)(t^2 + i) \in F[t]$ . We can show that the minimal polynomial of  $a$  over  $F$  is  $t^2 - i$ .

## Definition: Generated Subring

Let  $K/F$  be a field extension and let  $S \subseteq K$  be a subset.

The subring generated by  $S$  over  $F$  is defined to be  $F[S] :=$  the intersection of all subrings of  $K$  which contain  $F$  and  $S$ .

That is, the  $F$ -subalgebra generated by  $S$ .

## Exercise

$$F[S] = \{a \in K : a = f(s_1, \dots, s_n) \text{ for some } n \geq 0, f \in F[x_1, \dots, x_n], s_1, \dots, s_n \in S\}.$$



## Notation

$$S = \{a\} \rightsquigarrow F[a].$$

$$S = \{a_1, \dots, a_n\} \rightsquigarrow F[a_1, \dots, a_n].$$

Note that  $F[a] = \text{im}(F[t] \xrightarrow{\text{ev}_a} K)$  and  $F[a_1, \dots, a_n] = \text{im}(F[t_1, \dots, t_n] \xrightarrow{\text{ev}_a} K)$ .

## Definition: Generated Subfield

Let  $K/F$  be a field extension and let  $S \subseteq K$  be a subset.

Then the subfield generated by  $S$  over  $F$  is defined to be  $F(S) :=$  the intersection of all subfields of  $K$  which contain  $F$  and  $S$ .

Observe that  $F[S] \subset F(S)$ .

## Exercise

$$F(S) = \left\{ a \in K : a = \frac{\alpha}{\beta} \text{ for } \alpha, \beta \in F[S] \right\} = \text{Frac}(F[S]).$$

## Notation

$$S = \{a\} \rightsquigarrow F(a).$$

$$S = \{a_1, \dots, a_n\} \rightsquigarrow F(a_1, \dots, a_n).$$

## Definition: Finitely Generated Field Extension

A field extension  $K/F$  is finitely generated if  $K = F(S)$  for some  $S \subset K$  finite.

That is, finitely generated as a field over  $F$  not as an algebra over  $F$  or a vector space over  $F$ .

## Example

$F(t)/F$  is a finitely generated field extension but is not finitely generated as an  $F$ -algebra (exercise) nor as an  $F$ -vector space.

## Example

In  $F(t)/F$ , the indeterminant  $t \in F(t)$  is transcendental over  $F$ .

The evaluation homomorphism  $F[t] \hookrightarrow F(t)$ .

## April 10, 2024

Last time:  $K/F$ ,  $S \subset K$ ,  $F[S] \subset F(S)$ .

Example:  $S, T \subset K \rightsquigarrow F(S)(T) = F(S \cup T)$ .

$$F(a_1, \dots, a_n) = F(a_1, \dots, a_{n-1})(a_n).$$

## Remark

Let  $K/F$  be a field extension.

If  $a \in K$  is transcendental over  $F$ , then

$$F[t] \xrightarrow[\sim]{\text{ev}_a} F[a]$$

is an isomorphism. Hence,

$$F(a) \simeq \text{Frac}(F[t]) = F(t)$$

the field of rational functions.

Thus, the field extensions  $F(a)$  for  $a \in K$  transcendental over  $F$  are all isomorphic.

### Example

$\mathbb{Q}(\pi) \simeq \mathbb{Q}(e)$  are isomorphic fields.

### Bottom Line

Transcendental elements behave like indeterminates  $t$ .

The prototypical example is  $F(t)/F$ .

### Proposition

Let  $K/F$  be a field extension.

If  $a \in K$  is algebraic over  $F$ , then  $F[a] \simeq F[t]/(m_a(t))$  where  $m_a(t) \in F[t]$  is the minimal polynomial of  $a$  over  $F$ .

Moreover,  $F[a]$  is a field. Hence  $F[a] = F(a)$ .

Also,  $[F(a) : F] = \deg(m_a(t))$ .

An explicit  $F$ -basis of  $F(a)$  is  $\{1, a, a^2, \dots, a^{d-1}\}$  where  $d := \deg(m_a(t))$ .

### Proof

$$\begin{array}{ccc} F[t] & \xrightarrow{\text{ev}_a} & K \\ \downarrow & & \uparrow \\ F[t]/m_a(t) & \xrightarrow{\sim} & F[a] \end{array}$$

Now  $m_a(t)$  is irreducible, so  $(m_a(t))$  is a nonzero prime ideal.

$F[t]$  is PID, therefore every nonzero prime ideal is maximal.

Hence  $F[t]/(m_a(t))$  is a field.

Also,  $\dim_F(F[t]/(f(t))) = \deg(f)$ .

A basis  $\{\bar{1}, \bar{t}, \bar{t}^2, \dots, \bar{t}^{d-1}\}$ .

Suppose  $a_0\bar{t} + a_1\bar{t}^2 + \dots + a_{d-1}\bar{t}^{d-1} = 0 = a_0 + a_1\bar{t} + \dots + a_{d-1}\bar{t}^{d-1}$ .

That is,  $g(t) = a_0 + a_1t + \dots + a_{d-1}t^{d-1} \in (f(t))$ .

So  $f$  divides  $g$  but  $\deg(f) = d > d-1$ . Then  $g = 0$  in the new polynomial.

That is  $a_0 = a_1 = \dots = a_{d-1} = 0$ .

What is the span?  $g(t) = b_0 + b_1t + \dots + b_nt^n = b_0 + b_1t + \dots + b_{d-1}t^{d-1} + t^d(\dots)$ .

In  $F[t]/(f(t))$ ,  $f(t) = 0$ ,  $f(t) = t^d = a_{d-1} + \dots + a_0$ ,  $t^d = (a_{d-1}t^{d-1} + \dots + a_0)$  where  $g(t)$  is some polynomial of degree less than  $d$ .

Finally, note that  $F[t]/(m_a(t)) \xrightarrow{\sim} F[a]$  is given by  $\overline{f(t)} \mapsto f(a)$ .

Then  $\bar{1} \mapsto 1$ ,  $\bar{t} \mapsto a$ ,  $\dots$ ,  $\bar{t}^{d-1} \mapsto a^{d-1}$ .

### Remark

This proposition explains the choice of the term “degree” for  $[K : F]$ . If  $K = F(a)$  is generated by a single, algebraic element  $a \in K$ , then the  $[F(a) : F]$  is the degree of the minimal polynomial of  $a$  over  $F$ .

### Example 1

$\mathbb{Q}(i) = \mathbb{Q}[i] = \{a + bi : a, b \in \mathbb{Q}\}$  (since the minimal polynomial  $t^2 + 1$  of  $i$  over  $\mathbb{Q}$  has degree 2).  
 $\mathbb{Q}(\sqrt{2}) = \mathbb{Q}[\sqrt{2}] = \{a + b\sqrt{2} : a, b \in \mathbb{Q}\}$ .

### Example 2

$\xi_p := e^{2\pi i/p} \in \mathbb{C}$  for a prime  $p$  is a root of unity for

$$x^p - 1 = (x - 1) \overbrace{(x^{p-1} + x^{p-2} + \cdots + x + 1)}^{\Phi_p(x)}$$

Since  $\xi_p \neq 1$ ,  $\xi_p$  is a root of  $\Phi_p(x)$ .

Eisenstein's Criterion applied to  $\Phi_p(x+1)$  that  $\Phi_p(x)$  is irreducible over  $\mathbb{Z}$  and hence over  $\mathbb{Q}$ .

Hence  $\Phi_p(x)$  is the minimal polynomial of  $\xi_p$  over  $\mathbb{Q}$ .

Thus,  $\mathbb{Q}(\xi_p) = \mathbb{Q}[\xi_p] = \{a_0 + a_1\xi_p + a_2\xi_p^2 + \cdots + a_{p-2}\xi_p^{p-2} : a_i \in \mathbb{Q}\}$ .

### Example

Let  $p = 3$ .  $\mathbb{Q}(\xi_3) = \{a_0 + a_1\xi_3 : a_0, a_1 \in \mathbb{Q}\}$ .

So  $\mathbb{Q}(\xi_3) = \mathbb{Q}(\sqrt{3}i) = \mathbb{Q}(\sqrt{-3}) = \{a + b\sqrt{-3} : a, b \in \mathbb{Q}\}$ .

### Example 3

$\mathbb{R}(i) = \mathbb{R}[i] = \{a + bi : a, b \in \mathbb{R}\} = \mathbb{C}$ .

$\mathbb{R}[i] \simeq \mathbb{R}[t]/(t^2 + 1)$ .

## To Study:

Eisenstein's Criterion

Gauss' Lemma

## Remark

Given a field extension  $K/F$ , an element  $a \in K$  is algebraic over  $F$  if and only if  $[F(a) : F] < \infty$ .

The above proposition gives the  $\implies$  direction.

On the other hand, if  $a$  is transcendental then  $F(a) \simeq F(t)$  and  $[F(t) : F] = \infty$ .

## Proposition

Let  $K/F$  be a finite extension of degree  $n$ .

Every element of  $K$  is algebraic over  $F$  and has degree dividing  $n$ .

## Proof

$$[K : F] = [K : F(a)][F(a) : F]$$

## Corollary


Every irreducible polynomial  $f \in \mathbb{R}[x]$  has degree 1 or 2.

## Proof

(Assuming the FTA)

Let  $f \in \mathbb{R}[x]$  be irreducible. Then  $K : \mathbb{R}[x]/(f)$  is a field.

By the Fundamental Theorem of Algebra,  $f$  has a root  $a \in \mathbb{C}$ .



So  $\mathbb{R} \hookrightarrow \mathbb{R}[x]/(f) \hookrightarrow \mathbb{C}$ , and

$$2 = [\mathbb{C} : \mathbb{R}] = [\mathbb{C} : \mathbb{R}[x]/(f)][\mathbb{R}[x]/(f) : \mathbb{R}]$$

Therefore  $[\mathbb{R}[x]/(f) : \mathbb{R}] = \deg(f)$  is either 1 or 2.

## Definition: Algebraic Extension

A field extension  $K/F$  is said to be an algebraic extension if every element of  $K$  is algebraic over  $F$ .

## Example

We showed above that every finite extension is an algebraic extension.

## Theorem:

Let  $L/K/F$  be field extensions.

Then  $L/F$  is algebraic if and only if  $L/K$  is algebraic and  $K/F$  is algebraic.

## Proof

( $\implies$ ) trivial.

( $\impliedby$ ) Let  $a \in L$ . Then  $f(a) = 0$  for some nonzero polynomial  $f(t) \in K[t]$ . Write

$$f(t) = b_0 + b_1 t + \cdots + b_d t^d$$

with  $b_0, \dots, b_d \in K$ .

Each of these  $b_i$  is algebraic over  $F$ . Hence  $E := F(b_0, b_1, \dots, b_d)$  is a finite extension of  $F$ .

$$F(b_0, \dots, b_d)/F(b_0, \dots, b_{d-1})/\dots/F(b_0)/F$$

Note that  $f(t) \in E[t]$  and  $f(a) = 0$ , so  $a \in L$  is algebraic over  $E$ .

Observe

$$[E(a) : F] = \underbrace{[E(a) : E]}_{\text{finite}} \underbrace{[E : F]}_{\text{finite}}$$

so  $E(a)/F$  is finite, hence an algebraic extension.

Therefore  $a \in L$  is algebraic over  $F$ .

## Corollary

Let  $K/F$  be a field extension.

The elements of  $K$  which are algebraic over  $F$  form a subfield of  $K$ .

## Proof

Let  $a, b \in K$  be algebraic over  $F$ .

Then  $F(a, b)/F$  factors as  $F(a, b)/F(a)/F$ .

So  $F(a, b)/F$  is a finite, hence algebraic, extension.

Therefore  $a + b, a - b, ab, a^{-1}$  (for  $a \neq 0$ ) are algebraic over  $F$ .

## Example

Apply to  $\mathbb{C}/\mathbb{Q}$  to see that the collection of all algebraic numbers forms a subfield of  $\mathbb{C}$ .

$\overline{\mathbb{Q}}$  is defined as the field of algebraic numbers.

Recall: Theorem (Cantor),  $\overline{\mathbb{Q}}$  is countable.

Exercise:  $\overline{\mathbb{Q}}/\mathbb{Q}$  is an infinite extension.

## Adjoining Elements

Let  $F$  be a field and  $f(x) \in F[x]$  be irreducible.

Then  $K := F[x]/(f)$  is a field extension of degree  $\deg(f)$ .

Note:  $K = F(a)$  when  $a := \bar{x}$ .

Note also that  $a$  is a root of  $f(x)$ .

$$f(a) = f(\bar{x}) = \overline{f(x)} = 0$$

in  $K$ .

## Example 1

We could define  $\mathbb{C} := \mathbb{R}[x]/(x^2 + 1)$  and define  $i := \bar{x}$ .

## Example 2

Let  $F$  be a field.

Suppose  $a \in F$  which does not have a square root in  $F$  (i.e.  $\nexists \delta \in F$  such that  $\delta^2 = a$ ).

Then  $x^2 - a \in F[x]$  is irreducible.

Then  $K := F[x]/(x^2 - a)$  is a degree 2 extension.

Setting  $\delta := \bar{x} \in K$ ,  $K = F(\delta)$  and  $\delta^2 = a \in F$ .

In fact, every quadratic extension arises in this way – adjoining a square root.

## Example 3

Let  $F$  be a field of characteristic  $\neq 2$ .

Let  $K/F$  be a quadratic extension (i.e.  $[K : F] = 2$ ).

Let  $\delta \in K$  be an element such that  $\delta \notin F$ , then  $K = F(\delta)$  ( $K = F(\delta)/F$ ).

So the minimal polynomial of  $\delta$  over  $F$  is a quadratic polynomial.

$$m_\delta(t) = t^2 + bt + c \in F[t]$$

Consider  $\Delta := b^2 - 4c \in F$ .

Claim:  $\Delta$  does not have a square root in  $F$ . Otherwise

$$m_\delta(t) = \left(t - \frac{-b + \sqrt{\Delta}}{2}\right) \left(t - \frac{-b - \sqrt{\Delta}}{2}\right)$$

would not be irreducible.

Note:  $2\delta + b = \pm\sqrt{\Delta}$ . So

$$K = F(\delta) = F(2\delta + b) = F(\sqrt{\Delta})$$

**April 15, 2024**

## Remarks on Ruler and Compass Construction

### Game

Start with some “known” points in the plane and then we construct new points using a ruler and compass.

The ruler you may draw lines between two known points.

With the compass you may draw a circle with known center and known radius (distance between two known points).

### Example Problems

Squaring the circle: given a circle, construct, using ruler and compass, a square with the same area.

Trisecting a given angle.

Can you construct a regular  $n$ -gon using ruler and compass?

Doubling the cube: given a cube, construct a cube with double the volume.

### Remark

Starting with two given points, call one zero and the other one. Then we can construct a coordinate system system which identifies the plane with  $\mathbb{R}^2 = \mathbb{C}$ .

One can readily check that the constructible points (the points which are constructible starting from zero and one) form a subfield of  $\mathbb{C}$ .

One can readily check that every point whose coordinates are rational is constructible.

This transforms the geometric problems into problems of field theory.

That is, the field of constructible numbers is some field extension of  $\mathbb{Q}(i)$ .

### Key Point

Let  $K$  be a subfield of  $\mathbb{C}$  which contains  $i$  and is closed under complex conjugation.

e.g.  $K = \mathbb{Q}(i)$ . Then  $z = x + iy \in K$  if and only if  $x, y \in K$ .

Consider the intersection of two lines. Then the point of intersection is in  $K$ .

The coordinates of the points of intersection are rational expressions of the coordinates of the points which determine the lines.

Consider instead the intersection of a circle  $(x - a)^2 + (y - b)^2 = r^2$  and a line  $px + qy = h$  ( $p \neq 0$ ).

Then  $x = \frac{h - qy}{p}$  and

$$\left(\frac{h - qy}{p} - a\right)^2 + (y - b)^2 = r^2$$

is a quadratic equation in  $y$ . So  $y$  may be expressed as a rational expression, including square roots, of the coordinates of the points which determine the line and the circle.

Therefore, the point of intersection is contained in a quadratic extension of  $K$ .  
Finally, the intersection of two circles is the same as the intersection of line and circle.

### Theorem:

A point  $z \in \mathbb{C}$  is constructible if and only if there exists a tower of field extension  $\mathbb{Q} = F_0 \subseteq \cdots \subseteq F_m$  such that  $z \in F_m$  and each  $F_i/F_{i-1}$  is a quadratic extension.  
That is,  $F_i = F_{i-1}(u_i)$  where  $u_i^2 \in F_{i-1}$ .

### Corollary

If  $z \in \mathbb{C}$  is constructible, then  $z$  is algebraic and  $[\mathbb{Q}(z) : \mathbb{Q}]$  is a power of 2.

### Proof

$$[F_m : \mathbb{Q}(z)][\mathbb{Q}(z) : \mathbb{Q}] = [F_m : F_0] = 2^m$$

### Corollary

It is impossible to square the circle.

### Proof

A square of side length  $\sqrt{\pi}$ ,  $\sqrt{\pi}$  and  $\pi$  would be constructible. But Lindemann demonstrated that  $\pi$  is transcendental.

### Remark

Note that some angles can be trisected (e.g.  $\pi/2$ ).

### Corollary

Not every angle may be trisected.  
In particular,  $2\pi/3$  cannot be trisected.

### Proof

If we could trisect  $2\pi/3$ , then we could construct  $e^{2\pi i/9}$ .  
Let  $\xi = e^{2\pi i/9}$ . If  $\xi$  were constructible, then  $\alpha := \xi + \xi^{-1}$  would be constructible.  
Observe that  $\xi^3 = e^{2\pi i/3}$ . Then  $(\xi^3)^2 + \xi^3 = -1$ .  
Given that

$$\alpha^3 = (\xi + \xi^{-1})^3 = \xi^3 + 3\xi + 3\xi^{-1} + \overbrace{\xi^{-3}}^{\xi^6} = 3(\xi + \xi^{-1}) - 1 = 3\alpha - 1$$

we have that  $\alpha^3 - 3\alpha + 1 = 0$ . That is,  $\alpha$  is a root of  $x^3 - 3x + 1$  which is irreducible (check directly that it is irreducible over  $\mathbb{Z}$  and then Gauss implies it is irreducible over  $\mathbb{Q}$ ).  
Then  $[\mathbb{Q}(\alpha) : \mathbb{Q}] = 3$ , and we are done.

### Proposition

We can construct a regular  $n$ -gon using ruler and compass if and only if  $e^{2\pi i/n}$  is constructible.

## Corollary

We cannot construct a regular 9-gon using ruler and compass.

## Definition: Algebraic Independence

Let  $K/F$  be a field extension.

A collection of distinct elements  $a_1, \dots, a_n \in K$  are said to be algebraically independent over  $F$  if  $\forall f \in F[x_1, \dots, x_n]$ ,  $f(a_1, \dots, a_n) = 0$  implies  $f = 0$ .

A subset  $S \subseteq K$  is said to be algebraically independent over  $F$  if every finite subset of  $S$  is algebraically independent over  $F$ .

## Definition: Purely Transcendental Extension

$K/F$  is a purely transcendental extension if  $K = F(S)$  for some algebraically independent  $S \subseteq K$ .

## Example

$F(t_1, \dots, t_n)/F$  is a purely transcendental extension.

## Exercise

$K/F$  is purely transcendental if and only if  $K \cong F(\{t_i\}_{i \in S}) = \text{Frac}(F[\{t_i\}_{i \in S}])$ .

## Definition: Transcendence Basis

Let  $K/F$  be a field extension.

A transcendence basis for  $K/F$  is an algebraically independent subset  $S \subseteq K$  such that  $K/F(S)$  is an algebraic extension.

## Theorem:

Let  $K/F$  be a field extension and  $S \subseteq K$ . The following are equivalent

1.  $S$  is a transcendence basis for  $K/F$ .
2.  $S$  is maximal among all algebraically independent subsets of  $K/F$ .
3.  $S$  is minimal among all subsets of  $K$  such that  $K/F(S)$  is algebraic.

## Lemma:

A subset  $S \subseteq K$  is algebraically independent over  $F$  if and only if every element  $u \in S$  is transcendental over  $F(S \setminus \{u\})$ .

## Exercise

Using this lemma, prove the previous theorem.

## Theorem:

Let  $K/F$  be a field extension.



1. Any algebraically independent subset  $S \subseteq K$  is contained in the transcendence basis.
2. Any subset  $S \subseteq K$  such that  $K/F(S)$  is algebraic contains a transcendence basis.

In particular, a transcendence basis exists.

## Remark

Every field extension is an algebraic extension of a purely transcendental extension.

## Corollary / Exercise

Every finitely generated field extension  $K/F$  is an algebraic extension of  $F(t_1, \dots, t_n)$  for some  $n$ .

## Theorem:

Any two transcendence bases of  $K/F$  have the same cardinality.

## Definition: Transcendence Degree

Let  $K/F$  be a field extension.

The cardinality of a transcendence basis is called the transcendence degree of  $K/F$ .

$$\text{trdeg}_F(K) \quad \text{or} \quad \text{trdeg}(K/F)$$

## Examples

$$\text{trdeg}_F(F(t_1, \dots, t_n)) = n.$$

$$\text{trdeg}_F(K) = 0 \text{ if and only if } K/F \text{ is algebraic.}$$

## Exercise

$$\text{Given } L/K/F, \text{trdeg}(L/F) = \text{trdeg}(L/K) + \text{trdeg}(K/F).$$

## Example

Let  $X$  be a compact Riemann surface.

Then we have a field  $M(x)$  of meromorphic functions,  $M(x)/\mathbb{C}$ .

e.g.  $X = \mathbb{P}_{\mathbb{C}}^1 = S^2$ . Then, since every meromorphic function is rational,  $M(x) \cong \mathbb{C}(t)$ .

In general,  $\text{trdeg}_{\mathbb{C}}(M(x)) = 1$ .

For every finite extension,  $K/\mathbb{C}(t)$ , there exists a compact Riemann surface  $X$  such that  $M(x) = K$ .

There is an equivalence of categories between the compact Riemann surfaces with nonconstant holomorphic maps and the opposite category of finite extensions of  $\mathbb{C}(t)$ .

**April 17, 2023**

## Definition: Splitting Field

Given a polynomial,  $f \in F[t]$  we want to construct a field extension  $K/F$  in which  $f$  splits into linear factors:

$$f(t) = c(t - a_1)(t - a_2) \cdots (t - a_n)$$

where  $c \in F$ ,  $a_1, \dots, a_n \in K$ .

A splitting field for  $f \in F[t]$  is a field extension  $K/F$  such that  $f$  splits completely into linear factors as above over  $K$  and  $K$  is generated by the roots of  $f$  in  $K$  such that  $K = F(a_1, \dots, a_n)$ .

## Theorem:

Every polynomial  $f \in F[t]$  admits a splitting field.

## Proof

Key: if  $f \in F[t]$  is irreducible, then  $F[t]/(f(t))$  is a field which is an extension of  $F$

$$F \hookrightarrow F[t] \rightarrow F[t]/(f(t))$$

and  $f(t)$  has a root in it, namely  $\bar{t} \in F[t]/(f(t))$  ( $f(\bar{t}) = \overline{f(t)} = 0$ ).

For a general polynomial  $f(t) \in F[t]$ ,

$$f = cf_1f_2 \cdots f_m$$

with  $c \in F$ ,  $f_i$  irreducible ( $F[x]$  a UFD).

Take  $K_1 := F[t]/(f_1)$ . Over  $K_1$ , one of the irreducible factors (namely  $f_1$ ) gets a root.

Then consider the factorization of  $f$  into irreducibles over  $K_1$  and continue. Then

$$F = K_0 \hookrightarrow K_1 \hookrightarrow K_2 \hookrightarrow \cdots$$

By injection on the sum of the degrees of the nonlinear irreducible factors, this implies that we eventually obtain a field  $K_N$  in which  $f$  splits completely.

## Remark

Note that if  $K = F(a_1, \dots, a_n)$  is a splitting field of  $f$  where  $a_1, \dots, a_n$  are the roots of  $f$  in  $K$ , then  $K$  is a finite, hence algebraic, extension of  $F$ .

## Remark

Be careful. Even if  $f \in F[t]$  is irreducible, it is not enough in general just to go to  $K := F[t]/(f(t))$  in order to obtain a complete splitting.

## Example

Let  $F = \mathbb{Q}$ ,  $f(X) = X^4 - 2 \in \mathbb{Q}[X]$ .

This is irreducible by Eisenstein.

Take  $K := \mathbb{Q}[X]/(X^4 - 2)$ . Then  $x := \bar{X}$  is a root of the polynomial.

In  $K$ , we have that  $x^4 = 2$ ; in  $K[X]$ ,  $X^4 - 2 = x^4 - x^4 = (X^2 - x^2)(X^2 + x^2) = (X - x)(X + x)(X^2 + x^2)$ .

The last polynomial,  $X^2 + x^2$  is irreducible in  $K[X]$ .

We still need to do one more quadratic extension (adjoining  $i = \sqrt{-1}$ ) for  $X^2 + x^2$  to split as  $(X - ix)(X + ix)$ .

A splitting field of  $X^4 - 2$  over  $\mathbb{Q}$  is a degree 8 extension  $\mathbb{Q}(\sqrt[4]{2}, i)$ .

$$\begin{array}{c}
 \mathbb{Q}(\sqrt[4]{2}, i) \\
 \mid_2 \\
 \mathbb{Q}(\sqrt[4]{2}) \\
 \mid_4 \\
 \mathbb{Q}
 \end{array}$$

### Corollary

Let  $f_1, f_2, \dots, f_s \in F[t]$  be a finite collection of polynomials.

Then there exists a field extension  $K/F$  in which each polynomial  $f_1, \dots, f_s$  splits completely.

### Proof

Apply the theorem to  $f := f_1 f_2 \cdots f_s$ .

### Definition: Algebraically Closed Field

A field  $F$  is algebraically closed if it admits no non-trivial algebraic extension.

That is,  $K/F$  algebraic  $\implies K = F$ .

### Proposition:

A field  $F$  is algebraically closed if and only if every polynomial in  $F[t]$  splits completely into linear factors.

### Proof

( $\implies$ ) Let  $f \in F[t]$ .

By the theorem above, there exists a splitting field  $K$  of  $f$ . So  $K/F$  is algebraic, and therefore  $K = F$ .

So  $f$  splits completely over  $F$ .

( $\impliedby$ ) Suppose  $K/F$  is algebraic.

Let  $a \in K$ . The minimal polynomial of  $a$  over  $F$  is irreducible, hence linear. Therefore  $a \in F$ .

So  $K = F$ .

### Remark / Exercise

Every polynomial in  $F[t]$  splits completely if and only if every non-constant polynomial in  $F[t]$  has a root in  $F$ .

### Theorem:

Let  $F$  be a field.

There exists a field extension  $E/F$  with  $E$  algebraically closed.

### Aside: Zorn's Lemma

Take  $S := \{\text{proper ideals } I \subsetneq R\}$  ordered by inclusion,  $S \neq \emptyset$  since  $R \neq \emptyset$ .

If  $\{I_i\}$  a chain, then  $\bigcup I_i$  is an ideal.

Further, it is a proper ideal since  $1 \in \bigcup I_i \implies 1 \in I_i \implies I_i = R$  a contradiction.

Zorn's lemma implies that  $S$  contains a maximal ideal.

## Proof

Zorn's lemma implies that every non-zero commutative ring has a maximal ideal.

Therefore, every non-zero commutative ring admits a ring homomorphism  $R \rightarrow K$  to a field.

Let  $A := F[X_f]_{f \in F[t] \setminus F}$ , the (huge) polynomial algebra in infinitely many indeterminates, one for each non-constant polynomial  $f \in F[t] \setminus F$ .

Consider the ideal  $I := (f(X_f))_{f \in F[t] \setminus F} \subseteq A$ .

Here  $f(X_f)$  is the single-variable polynomial  $f$  in the variable  $X_f$ .

Key:  $I \neq A$ . Let  $Q = \sum_{i=1}^n Q_i f_i(X_{f_i}) \in I$ , for some  $Q_i \in A$  and  $f_i \in F[t] \setminus F$ .

There exists a field  $K/F$  in which  $f_1, \dots, f_n$  split completely.

In particular,  $\exists \alpha_1, \dots, \alpha_n \in K$  such that  $f_i(\alpha_i) = 0$ ,  $\forall i = 1, \dots, n$ .

Evaluate  $Q(X_{f_1}, \dots, X_{f_n}, X_{q_1}, \dots, X_{q_k})$  by setting  $X_{f_i} \mapsto \alpha_i$  and  $X_{q_j} \mapsto 0$  (for example).

Then  $Q \mapsto 0$ . Therefore,  $Q \neq 1$  because  $1 \mapsto 1$  in  $K$ .

So  $1 \notin I$  (i.e.  $I \neq A$ ), hence  $R := A/I$  is a non-zero ring and there exists a ring homomorphism  $R \rightarrow L$  for some field  $L$ .

This field  $L$  is an extension of  $F$ .

$$\begin{array}{ccccc}
 F & \hookrightarrow & & & L \\
 & \searrow & & \nearrow \phi & \uparrow \\
 & & F[\{X_f\}] = A & \longrightarrow & A/I = R
 \end{array}$$

Let  $f \in F[t] \setminus F$ .

Consider  $\alpha_f := \phi(X_f) \in L$ .

Well

$$f(\alpha_f) = f(\phi(X_f)) = \phi(\underbrace{f(X_f)}_{\in I}) = 0$$

That is, every non-constant polynomial in  $F[t]$  has a root in  $L$ .

Repeat this process inductively,

$$F = L_0 \hookrightarrow L_1 \hookrightarrow \dots \hookrightarrow L_n \hookrightarrow \dots$$

with the property that every non-constant polynomial with coefficients in  $L_n$  has a root in  $L_{n+1}$ .

Take  $L := \text{colim}_{n \rightarrow \infty} L_n = \bigcup_{n \geq 1} L_n$ . This  $L$  is algebraically closed.

$$L[T] = \bigcup_{n \geq 1} L_n[T]$$

Every non-constant polynomial over  $L$  is realized over  $L_n$ , hence has a root in  $L_{n+1}$  and therefore has a root in  $L$ .

## Definition: Algebraic Closure

Let  $F$  be a field.

An algebraic closure of  $F$  is an algebraic extension  $E/F$  with  $E$  algebraically closed.

## Corollary

Every field admits an algebraic closure.

## Proof

By the theorem,  $F \hookrightarrow L$  with  $L$  algebraically closed.

Consider  $E := \{a \in L : a \text{ is algebraic over } F\}$ . We previously proved that  $E$  is a subfield of  $L$ .

By construction,  $E/F$  is algebraic.

We claim  $E$  is algebraically closed.

Let  $f \in E[t]$  be a polynomial. Well

$$f(t) = c(t - \alpha_1) \cdots (t - \alpha_n)$$

in  $L[t]$  with  $c \in E$  and  $\alpha_1, \dots, \alpha_n \in L$ .

Each  $\alpha_i$  is algebraic over  $E$  (since  $f(\alpha_i) = 0$ ).

So  $E(\alpha_i)/E$  is algebraic and  $E/F$  is an algebraic extension.

Therefore  $E(\alpha_i)/F$  is algebraic. Therefore  $\alpha_i$  is algebraic over  $F$  so  $\alpha_i \in E$ .

So  $f$  splits on  $E$ .

## Example

The Fundamental Theorem of Algebra asserts that  $\mathbb{C}$  is algebraically closed.

## Example

$\overline{\mathbb{Q}} = \{z \in \mathbb{C} : z \text{ is algebraic over } \mathbb{Q}\}$  is an algebraic closure of  $\mathbb{Q}$ .

## Example

$\mathbb{C}$  is an algebraic closure of  $\mathbb{R}$ .