

# Set theory

Johannes Hahn

January 8, 2026

## 1 Well-ordering

### 1.1 Definition:

Let  $A$  be a well-ordered set. An initial segment is a subset  $S \subseteq A$  such that

$$\forall \alpha \in S, \alpha' \in A : \alpha' \leq \alpha \implies \alpha' \in S$$

### 1.2 Lemma:

Let  $A$  be any well-ordered set.

- a.)  $\emptyset$  and  $A$  itself are initial segments. Sets of the form  $A_{<\alpha}$  and  $A_{\leq\alpha}$  are initial segments.
- b.) All *proper*, initial segments of  $A$  are of the form  $A_{<\alpha}$  for a unique  $\alpha \in A$ , namely the smallest element of  $A \setminus S$ .
- c.)  $\alpha' \leq \alpha \iff A_{<\alpha'} \subseteq A_{<\alpha}$ . In particular: The set of initial segments

$$A^+ = \{ A_{<\alpha} \mid \alpha \in A \} \cup \{ A \}$$

is itself well-ordered w.r.t. to inclusion and  $A$  is its greatest element.

### 1.3 Lemma (Transfinite induction along well-ordered sets):

Let  $A$  be a well-ordered set and  $\phi$  a formula with a free variable.

- a.) If  $\phi$  satisfies

$$(\forall \alpha' < \alpha : \phi(\alpha')) \implies \phi(\alpha)$$

for all  $\alpha \in A$ , then  $\forall \alpha \in A : \phi(\alpha)$  holds.

- b.) If  $\phi$  satisfies

- i.)  $\phi(0)$  holds

- ii.) If  $\phi(\alpha')$  holds and  $\alpha$  is the successor of  $\alpha'$ , then  $\phi(\alpha)$  holds too.
  - iii.) If  $\alpha$  is a limit and  $\phi(\alpha')$  holds for all  $\alpha' < \alpha$ , then  $\phi(\alpha)$  holds too.
- then  $\forall \alpha \in A : \phi(\alpha)$  holds.

*Proof.* Obviously, it is sufficient to prove a., since b. is just a restatement.  
Let  $F := \{ \alpha \in A \mid \neg \phi(\alpha) \}$ . Assume  $F \neq \emptyset$  and let  $\alpha$  be its smallest element. Then  $\phi(\alpha')$  holds for all  $\alpha' < \alpha$ . But then by assumption on  $\phi$ ,  $\phi(\alpha)$  also holds contradicting to the minimality of  $\alpha$ .  $\square$

**1.4 Lemma:** a.) For any two well-ordered sets  $A, B$  there is at most one isomorphism  $A \rightarrow B$ .

- b.) The only initial segment of  $A$  isomorphic to  $A$  is  $A$  itself. In particular: All initial segments of  $A$  are pairwise non-isomorphic.

*Proof.* a. Transfinite induction along  $A$ : Let  $f, g : A \rightarrow B$  be two isomorphism,  $\alpha \in A$  and assume  $f(\beta) = g(\beta)$  for all  $\beta < \alpha$ . Set  $A' := A_{<\alpha}$ .  
Then by construction  $\alpha$  is the smallest element of  $A \setminus A'$  and therefore the isomorphisms  $f$  and  $g$  must map it to the smallest element of  $B \setminus f(A')$  and  $B \setminus g(A')$  respectively. But by assumption,  $f(A') = g(A')$  so that  $f(\alpha) = g(\alpha)$ .  
Therefore,  $f(\alpha) = g(\alpha)$  for all  $\alpha \in A$ .

b. Assume  $f : A \rightarrow A_0$  is an isomorphism onto an initial segment  $A_0 \subsetneq A$ . If there was an  $\alpha \in A \setminus A_0$ , then  $f(\alpha) < \alpha$ . But then by induction  $f^{n+1}(\alpha) < f^n(\alpha)$  for all  $n \in \mathbb{N}$  so that we get an infinite descending sequence in  $A$  which does not exist because  $A$  is well-ordered. Therefore  $A_0 = A$ .  $\square$

**1.5:** This means that we can strengthen the previous observation:  $\alpha' \leq \alpha$  iff  $A_{<\alpha'}$  is merely isomorphic to an initial segment of  $A_{<\alpha}$ .

**1.6 Lemma** (Transfinite recursion along well-ordered sets):

Let  $A$  be a well-ordered set and  $B$  an arbitrary set.

- a.) Let

$$g : \text{part.Maps}(A \rightarrow B) \rightarrow B$$

Then there is a unique function  $f : A \rightarrow B$  such that

$$\forall \alpha : f(\alpha) = g(f|_{A_{<\alpha}})$$

- b.) Let  $h : P(B) \rightarrow B$ . Then there is a unique function  $f : A \rightarrow B$  such that

$$\forall \alpha : f(\alpha) = h(f(A_{<\alpha}))$$

**1.7:** In a recursion over  $A = \mathbb{N}$ , we might define  $f(n)$  by any computation  $g$  that may depend on some or all of  $f(0), f(1), \dots, f(n-1)$ , i.e. any finite-length sequence of values in  $B$ .

The appropriate generalisation to sequences with well-ordered index sets would be something like  $\bigcup_{S \in A^+ \setminus \{A\}} \text{Maps}(S \rightarrow B)$ , i.e. certain partial maps  $A \rightarrow B$ .

Note that the set of partial maps  $A \rightarrow B$  is may be identified with the set

$$\{ G \in P(A \times B) \mid \forall x \in A \forall y, y' \in B : (x, y) \in G \wedge (x, y') \in G \implies y = y' \}$$

*Proof.* It is sufficient to prove a., because b. is the special case where  $g$  is the function  $g(f) := h(\text{im}(f))$ .

The idea is to apply transfinite induction to the statement “ $f$  is uniquely defined at  $\alpha$ ”. Consider

$$D := \left\{ S \in A^+ \mid \exists ! f_S : S \rightarrow B : \forall \alpha \in S : f(\alpha) = g(f_{\upharpoonright A_{<\alpha}}) \right\}$$

We prove  $D = A^+$  by transfinite induction along  $A^+$ :

Step 1.  $\emptyset \in D$  because the empty map has that property.

Step 2: If  $S = A_{<\alpha} \in D$ , then  $\tilde{S} = A_{\leq \alpha} \in D$  as well, because

$$f_{\tilde{S}}(\beta) = \begin{cases} g(f_S) & \beta = \alpha \\ f_S(\beta) & \beta < \alpha \end{cases}$$

is the unique extension of  $f_S$  to  $\tilde{S}$  with the required property.

Step 3:  $D$  is closed under unions.

If  $S$  is in  $D$ , then any smaller initial segment  $S' \subseteq S$  is also in  $D$ , because the restriction of  $f_S$  also has the property. In particular  $f_{S'} = (f_S)_{\upharpoonright S'}$ , i.e. the functions are all compatible with each another.

That means that for any  $D' \subseteq D$ ,  $S := \bigcup_{S' \in D'} S'$  is an initial segment and  $f_S := \bigcup_{S' \in D'} f_{S'}$  is the unique function with the required property.

Now note that the successor in  $A^+$  of a proper initial segment  $A_{<\alpha}$  is  $A_{\leq \alpha}$ . Therefore, steps 1, 2, and 3 correspond to the 0-case, the successor-case, and the limit-case respectively and transfinite inductions shows that all of  $A^+$  is contained in  $D$ . In particular  $A \in D$ .  $\square$

**1.8 Theorem** (“The class of well ordered sets is well-ordered”):

Given two well-ordered sets  $A, B$  exactly one of the following happens:

- a.)  $A$  is isomorphic to a proper initial segment of  $B$
- b.)  $A \cong B$
- c.)  $B$  is isomorphic to a proper initial segment of  $A$

*Proof.* By the previous lemma, the three cases are mutually exclusive. Thus, we assume that we're neither in the second nor the third case and will prove that we're in the first.

We build a strictly monoton map  $f : A \hookrightarrow B$  such that the image of  $f$  is a proper initial segment of  $B$  by transfinite recursion along  $A$ .

Note that the empty map is already a strictly monoton map  $\emptyset \rightarrow B$  whose image  $\emptyset$  is an initial segment so that we're done if  $A$  is empty and  $B$  is non-empty (which it must be if  $A = \emptyset$  because we're not in case 2 by assumption).

Assume that  $f$  is already defined for all  $\alpha' < \alpha$  and that  $\text{im}(f) = f(A_{<\alpha})$  a proper initial segment of  $B$ , i.e.  $\text{im}(f) = B_{<\beta}$  for some  $\beta \in B$ .

Define an extension of  $f$  by  $\hat{f}(\alpha) := \beta$ . That is strictly monoton by construction and

$$\text{im}(\hat{f}) = \{ \beta \} \cup \text{im}(f) = \{ \beta \} \cup B_{<\beta} = B_{\leq \beta}$$

is an initial segment of  $B$ . It is proper because otherwise  $\hat{f}$  would be an isomorphism between  $A_{\leq \alpha}$  and all of  $B$  so that we'd be in cases 2 (if  $\alpha = \max A$ ) or 3 (otherwise) contrary to our assumption.  $\square$