

# AIIMS CRE OT Technician PYQ

Memory-Based Previous Year Questions & Explanations (Part 4: Q61-Q80)

## Q61. What is the next step after preoxygenation?

- A) Reversal
- B) Muscle relaxant
- C) IV induction
- D) Intubation

**Answer: C) IV induction**

**Explanation:** Preoxygenation prepares the patient for the induction phase. The immediate next step is the administration of an IV induction agent, which is then followed by a muscle relaxant and eventually intubation.

## Q62. Which equipment is kept ready before induction?

- A) Ambu bag
- B) Laryngoscope
- C) Suction
- D) All of these

**Answer: D) All of these**

**Explanation:** Before inducing a patient, all airway equipment must be ready. An Ambu bag is needed for manual positive ventilation, suction is required for clearing secretions (creating negative pressure), and a laryngoscope is used to visualize the glottis.

## Q63. Perforated eardrum surgery is known as:

- A) Myringotomy
- B) Tympanoplasty
- C) Stapedectomy
- D) Tonsillectomy

**Answer: B) Tympanoplasty**

**Explanation:** Tympanoplasty is the surgical grafting or repair of a perforated eardrum (tympanic membrane). Myringotomy is merely an incision to drain accumulated fluid.

**Q64. The commonest treatment of branchial cyst is:**

- A) Cystectomy
- B) Aspiration
- C) Excisional biopsy
- D) None of these

**Answer: C) Excisional biopsy**

**Explanation:** The standard surgical procedure for treating a branchial cyst is complete surgical removal via an excisional biopsy, preventing recurrence compared to simple aspiration.

**Q65. As per ASA guidelines, which is NOT correct regarding pre-operative fasting?**

- A) Light meal up to 6 hours
- B) Clear fluids up to 2 hours
- C) Pulp juice up to 2 hours
- D) Both A and B

**Answer: C) Pulp juice up to 2 hours**

**Explanation:** Clear fluids can be taken up to 2 hours before surgery, but fruit juices containing pulp delay gastric emptying and are not allowed, as they compromise the NPO status.

**Q66. The most common triggers of anaphylaxis under anesthesia are:**

- A) Muscle relaxants
- B) Antibiotics
- C) Latex
- D) All of these

**Answer: D) All of these**

**Explanation:** Muscle relaxants (neuromuscular blockers), intraoperative antibiotics, and latex are the three most frequent causative agents of allergic (anaphylactic) shocks in the operating theater.

**Q67. Total body fluid in a child accounts for how much percent of body weight?**

- A) 60%
- B) 70%
- C) 75%
- D) 65%

**Answer: D) 65%**

**Explanation:** Total body water changes with age: neonates are around 75-80%, children are typically 50-65%, and average adults are around 55-60%.

**Q68. Maximum concentration of potassium is present in:**

- A) Normal saline
- B) Ringer lactate
- C) D5
- D) Darrow's solution

**Answer: D) Darrow's solution**

**Explanation:** Darrow's solution is a specialized crystalloid fluid that contains a high concentration of potassium, explicitly used to manage severe hypokalemia.

**Q69. Which IV induction agent can cause hypertension?**

- A) Thiopentone
- B) Ketamine
- C) Propofol
- D) Etomidate

**Answer: B) Ketamine**

**Explanation:** Ketamine stimulates the sympathetic nervous system, increasing heart rate, blood pressure, and cardiac output, unlike Propofol and Thiopentone which tend to decrease blood pressure.

**Q70. Which LMA is used as an intubation guide for an ET tube?**

- A) ProSeal LMA
- B) Ambu LMA
- C) Classic LMA
- D) ILMA (Intubating Laryngeal Mask Airway)

**Answer: D) ILMA**

**Explanation:** The ILMA (Intubating Laryngeal Mask Airway), also known as Fastrach, is uniquely designed to act as a conduit to facilitate blind endotracheal intubation.

**Q71. Which is NOT a component of the APGAR score?**

- A) Muscle tone
- B) Color of the body
- C) Heart rate
- D) ABG (Arterial Blood Gas)

**Answer: D) ABG**

**Explanation:** The APGAR score evaluates Appearance (color), Pulse (heart rate), Grimace (reflex), Activity (muscle tone), and Respiration. ABG is an invasive lab test, not part of this scoring system.

**Q72. Which drug does NOT cause cardiac depression?**

- A) Propofol
- B) Thiopentone
- C) Ketamine
- D) Halothane

**Answer: C) Ketamine**

**Explanation:** Ketamine maintains or increases cardiac output and heart rate, thus avoiding the cardiac depression commonly seen with agents like propofol (which causes vasodilation).

**Q73. Which of the following anesthetic agents causes post-operative delirium and hallucinations?**

- A) Etomidate
- B) Sevoflurane
- C) Propofol
- D) Ketamine

**Answer: D) Ketamine**

**Explanation:** Ketamine is known for its psychological side effects during recovery, including emergence delirium, vivid dreams, and auditory/visual hallucinations.

**Q74. NaOH in soda lime acts as:**

- A) Directly absorption of CO<sub>2</sub>
- B) A catalyst
- C) Prevention formation of water
- D) None of these

**Answer: B) A catalyst**

**Explanation:** Sodium hydroxide (NaOH) functions as a catalyst to speed up the chemical reaction in which soda lime absorbs carbon dioxide from the anesthesia circuit.

**Q75. The color indicator in soda lime that changes from white to violet is:**

- A) Phenolphthalein
- B) Methyl orange
- C) Ethyl violet
- D) Litmus

**Answer: C) Ethyl violet**

**Explanation:** Ethyl violet is a pH indicator used in soda lime. As the granules exhaust their absorbing capacity, the pH changes, causing the white granules to turn violet.

**Q76. Depth of anesthesia is best measured by:**

- A) BIS (Bispectral Index System)
- B) MAC (Minimum Alveolar Concentration)
- C) TOF (Train of Four)
- D) PTP (Post Tetanic Potentiation)

**Answer: A) BIS**

**Explanation:** The Bispectral Index System (BIS) processes EEG signals to provide a direct, numerical measurement of the patient's brain activity and depth of anesthesia. TOF measures muscle paralysis, not anesthesia depth.

**Q77. The color of a Nitrous Oxide cylinder is:**

- A) Black with white shoulder
- B) French Blue
- C) Grey
- D) White

**Answer: B) French Blue**

**Explanation:** Standard medical gas color coding designates French Blue for Nitrous Oxide (N<sub>2</sub>O) cylinders.

**Q78. Maximum airway irritation is caused by:**

- A) Desflurane
- B) Enflurane
- C) Sevoflurane
- D) Halothane

**Answer: A) Desflurane**

**Explanation:** Desflurane has a highly pungent odor and causes severe airway irritation, making it unsuitable for inhalational induction (unlike Sevoflurane, which is smooth and non-irritating).

**Q79. Airway resistance to the flow of gases through an endotracheal tube is primarily due to:**

- A) Curvature
- B) Length
- C) Material
- D) Diameter

---

**Answer: D) Diameter**

**Explanation:** Based on the principles of fluid dynamics (Poiseuille's law), resistance is inversely proportional to the radius to the fourth power. Thus, the diameter is the most critical factor.

**Q80. A patient on Halothane develops muscle rigidity and hyperthermia. Which agent also contributed to this condition?**

- A) Succinylcholine
- B) D-tubocurarine
- C) Rocuronium
- D) Vecuronium

---

**Answer: A) Succinylcholine**

**Explanation:** Malignant Hyperthermia is a severe reaction triggered by a combination of volatile anesthetics (like Halothane) and depolarizing muscle relaxants (like Succinylcholine).