

# Computational Linear Algebra

## EK103

Giacomo Cappelletto

21/1/2

# Contents

<b>Chapter 1</b>	<b>Basics</b>	<b>Page 2</b>
1.1	Vectors, Norms and Products Interpretation of vectors in $\mathbb{R}^{2,3}$ — 6	2
1.2	Rotation Matrices Examples of 2D Matrix Transformations — 8	7
1.3	Mapping Method 1 – Vector Method — 9 • Method 2 – Vertex Method — 9	9
1.4	A System of Linear Equations	11
1.5	How to Determine Consistency, Uniqueness, and the Number of Solutions for a Linear System FRR — 13	11
1.6	Back Row Reduction	13
<b>Chapter 2</b>	<b>Linear Combinations</b>	<b>Page 16</b>
2.1	Matrix–Vector Product as Linear Combination	16
2.2	Linear Combinations and the Span	17
2.3	Geometric Plots in the Notes Key Conclusions — 18 • Dropping a vector to retain the same span — 18	17
2.4	Evaluating Linear Dependence Procedurally	19
2.5	Identity Matrix ERP's and Identity Matrices — 23	20
<b>Chapter 3</b>	<b>Vector Spaces</b>	<b>Page 24</b>
3.1	Column Space	25
3.2	Linear Combinations and Column Space Basic Definitions and Concepts — 25 • Dimension and Basis of the Column Space — 25 • Parametric Form of the Column Space — 26 • Geometry of the Column Space — 26 • Example — 26	25
3.3	Nullspace of a Matrix	27
3.4	Row Space of a Matrix	29
3.5	Dimension and Basis of Row Space	29
<b>Chapter 4</b>	<b>Properties of Matrices</b>	<b>Page 31</b>
4.1	Determinant of a Square Matrix Cofactor Algorithm for Computing Determinant — 31 • Properties of Determinants — 32	31
4.2	EigenVectors and EigenValues	32
4.3	Properties of Eigenvectors	36

# Chapter 1

## Basics

### 1.1 Vectors, Norms and Products

**Note:-**

Let us consider two vectors in  $\mathbb{R}^3$ :

$$u = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad \text{and} \quad v = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}.$$

We wish to compute their magnitudes (norms and norm-squared), the angle between them, and the plane that they span. These methods are directly applicable to computational tools such as MATLAB.

**Definition 1.1.1: Norm of a Vector**

For a vector  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ , its norm is

$$\|x\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}.$$

In many programming languages (including MATLAB), this is computed via `norm(x)`, while the square of the norm is  $\|x\|^2 = x \cdot x = x_1^2 + \dots + x_n^2$ .

Norm squared is the result of the dot product of a vector with itself. For example, the norm squared of  $x$  is

$$\|x\|^2 = x \cdot x = x_1^2 + \dots + x_n^2 = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \cdot \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix}.$$

**Example 1.1.1** (Norms and Norm-Squared of  $u$  and  $v$ )

$$\|u\| = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}, \quad \|v\| = \sqrt{1^2 + (-1)^2 + 1^2} = \sqrt{3}.$$

Thus, both vectors have the same magnitude  $\sqrt{3}$ . Their squared norms are

$$\|u\|^2 = 3, \quad \|v\|^2 = 3.$$

In MATLAB notation, one could write:

- `norm(u)` or `norm(u,2)` for the norm of  $u$ .
- `dot(u,u)` or `norm(u)^2` for  $\|u\|^2$ .

### Definition 1.1.2: Angle Between Two Vectors

The angle  $\theta$  between two nonzero vectors  $u$  and  $v$  in  $\mathbb{R}^n$  is given by

$$\theta = \arccos\left(\frac{u \cdot v}{\|u\|\|v\|}\right).$$

#### Example 1.1.2 (Angle Between $u$ and $v$ )

First, compute the dot product:

$$u \cdot v = (1)(1) + (1)(-1) + (1)(1) = 1 - 1 + 1 = 1.$$

Hence,

$$\theta = \arccos\left(\frac{u \cdot v}{\|u\|\|v\|}\right) = \arccos\left(\frac{1}{\sqrt{3}\sqrt{3}}\right) = \arccos\left(\frac{1}{3}\right).$$

In MATLAB, one could write:

$$\text{theta} = \text{acos}(\text{dot}(u,v)/(\text{norm}(u)*\text{norm}(v)));$$

### Definition 1.1.3: Plane Spanned by Two Vectors

The plane containing vectors  $u$  and  $v$  and passing through the origin is given by

$$\{ \alpha u + \beta v \mid \alpha, \beta \in \mathbb{R} \}.$$

An equivalent description is all points  $x \in \mathbb{R}^3$  such that  $x \cdot (u \times v) = 0$ .

#### Example 1.1.3 (Plane Containing $u$ and $v$ )

- *Span form:*

$$\text{Plane} = \left\{ \alpha \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \beta \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \mid \alpha, \beta \in \mathbb{R} \right\}.$$

- *Normal form:* The cross product

$$u \times v = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 1 \\ 1 & -1 & 1 \end{vmatrix} = (2, 0, -2).$$

Hence, the plane also can be described by the set of points  $x = (x_1, x_2, x_3)$  for which

$$(2, 0, -2) \cdot (x_1, x_2, x_3) = 0 \implies 2x_1 - 2x_3 = 0 \implies x_1 = x_3.$$

In many computational environments, one simply keeps the span form or uses a symbolic package to compute the cross product and normal equation.

### Definition 1.1.4: Cross Product

Construct a system of linear equations where the dot product of the vector is orthogonal to both the vectors in the matrix.

$$\begin{bmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

**Example 1.1.4** (Finding the Plane Spanned by Two Vectors)

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = t \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

Then, we must formulate an equation for any vector perpendicular to the normal of the plane, i.e. the cross product of the two original vectors.

$$\begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} = 0$$

Hence,

$$y_1 - y_3 = 0$$

**Definition 1.1.5: Dot product**

We can take a vector  $\vec{v}$  in  $\mathbb{R}^n$  and a vector  $\vec{w}$  in  $\mathbb{R}^n$ . Then, the dot product of  $\vec{v}$  and  $\vec{w}$  is defined as

$$\vec{v} \cdot \vec{w} = \vec{v}^T \vec{w} = \vec{w}^T \vec{v}$$

Therefore

$$\|\vec{v}\|^2 = \vec{v}^T \vec{v}$$

**Definition 1.1.6: Scalar Multiplication**

Scalar multiplication is the operation of multiplying a vector by a scalar. The result is a new vector with the same direction as the original vector, but with a magnitude that is the product of the original magnitude and the scalar.

$$t \cdot \vec{v} = t \cdot \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} t a_1 \\ t a_2 \\ \vdots \\ t a_n \end{bmatrix}$$

Where

$$\|t \cdot \vec{v}\| = \|\vec{v}\| \cdot |t|$$

**Definition 1.1.7: Vector Addition**

Vector addition is the operation of adding two vectors together. The result is a new vector that is the sum of the two original vectors.

$$\vec{v} + \vec{w} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} = \begin{bmatrix} a_1 + b_1 \\ a_2 + b_2 \\ \vdots \\ a_n + b_n \end{bmatrix}$$

### Definition 1.1.8: Matrix to Vector Multiplication

Matrix to vector multiplication is the operation of multiplying a matrix by a vector. The result is a new vector that is the result of the matrix-vector multiplication. Matrices are represented as  $n \times m$ , where  $n$  is the number of rows and  $m$  is the number of columns. The number of columns of the matrix must be equal to the number of rows of the vector.

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n \end{bmatrix}$$

### Definition 1.1.9: Matrix to Matrix Multiplication

Matrix to matrix multiplication is the non-commutative operation of multiplying two matrices together. The result is a new matrix that is the product of the two original matrices.

For two matrices  $A$  and  $B$  to be multiplied, the number of columns of  $A$  must be equal to the number of rows of  $B$ . If  $A$  is an  $m \times n$  matrix and  $B$  is an  $n \times p$  matrix, then their product  $C = AB$  is an  $m \times p$  matrix. The element  $c_{ij}$  of the resulting matrix  $C$  is computed as:

$$c_{ij} = \sum_{k=1}^n a_{ik}b_{kj}$$

where  $a_{ik}$  is the element from the  $i$ -th row and  $k$ -th column of matrix  $A$ , and  $b_{kj}$  is the element from the  $k$ -th row and  $j$ -th column of matrix  $B$ .

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1p} \\ b_{21} & b_{22} & \cdots & b_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{np} \end{bmatrix} = \begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1p} \\ c_{21} & c_{22} & \cdots & c_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ c_{m1} & c_{m2} & \cdots & c_{mp} \end{bmatrix}$$

### Example 1.1.5 (Matrix to Matrix Multiplication)

Consider the matrices

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 7 & 8 & 9 \\ 10 & 11 & 12 \end{bmatrix}.$$

Their product  $C = AB$  is computed as follows:

$$C = \begin{bmatrix} 1 \cdot 7 + 2 \cdot 10 & 1 \cdot 8 + 2 \cdot 11 & 1 \cdot 9 + 2 \cdot 12 \\ 3 \cdot 7 + 4 \cdot 10 & 3 \cdot 8 + 4 \cdot 11 & 3 \cdot 9 + 4 \cdot 12 \\ 5 \cdot 7 + 6 \cdot 10 & 5 \cdot 8 + 6 \cdot 11 & 5 \cdot 9 + 6 \cdot 12 \end{bmatrix} = \begin{bmatrix} 27 & 30 & 33 \\ 61 & 68 & 75 \\ 95 & 106 & 117 \end{bmatrix}.$$



Figure 1.1: Matrix-Matrix Multiplication Diagram

**Definition 1.1.10: Matrix Multiplication of a Matrix with Itself**

When a matrix  $A$  is multiplied by its transpose  $A^T$ , the resulting matrix is a symmetric matrix. The element  $c_{ij}$  of the resulting matrix  $C = AA^T$  is computed as:

$$c_{ij} = \sum_{k=1}^n a_{ik}a_{jk}$$

where  $a_{ik}$  is the element from the  $i$ -th row and  $k$ -th column of matrix  $A$ , and  $a_{jk}$  is the element from the  $j$ -th row and  $k$ -th column of matrix  $A$ .

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1m} \\ c_{21} & c_{22} & \cdots & c_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ c_{m1} & c_{m2} & \cdots & c_{mm} \end{bmatrix}$$

The zeros in this matrix represent orthogonality between the corresponding rows of the original matrix  $A$ . Specifically, if  $c_{ij} = 0$ , it means that the  $i$ -th row and the  $j$ -th row of matrix  $A$  are orthogonal to each other.

**Example 1.1.6 (Orthogonality in Matrix Multiplication)**

Consider the matrix

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Its transpose is

$$A^T = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

The product  $C = AA^T$  is computed as follows:

$$C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Notice that the resulting matrix is the identity matrix, which is symmetric and has zeros in all off-diagonal elements, indicating that the rows (and columns) of the original matrix  $A$  are orthogonal to each other.

**1.1.1 Interpretation of vectors in  $\mathbb{R}^{2,3}$** **Definition 1.1.11: Position Vector**

A position vector is a vector that describes the position of an object in space with reference to an origin.

**Definition 1.1.12: Translational Vector**

A translational vector is a vector that describes the displacement of an object in space with reference to an origin.

## 1.2 Rotation Matrices

### Note:-

We are considering vectors in  $\mathbb{R}^2$ , denoted by

$$\vec{v} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}.$$

We also have a  $2 \times 2$  matrix (linear operator or transformation) given by

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}.$$

### Note:-

Applying  $A$  to an input vector  $\vec{v}_i$  produces the output vector  $\vec{v}_o$ :

$$A \vec{v}_i = \vec{v}_o.$$

Explicitly,

$$\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}.$$

### Question 1: Transforming a region

Suppose we restrict the input vectors  $\vec{v}_i$  to the square region in the  $(x_1, x_2)$ -plane with coordinates

$$0 \leq x_1 \leq 1, \quad 0 \leq x_2 \leq 1.$$

- How does  $A$  map this square region in the input space to a region in the  $(y_1, y_2)$ -plane?
- Geometrically, what does that image region look like?

**Solution:** We can answer this question *mechanically* by observing what happens to the corners of the square, or by spanning the space with two basis vectors:

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

Each point in the square can be written as

$$x_1 \vec{e}_1 + x_2 \vec{e}_2 \quad \text{with } 0 \leq x_1, x_2 \leq 1.$$

Applying  $A$  to these basis vectors, we get

$$A \vec{e}_1 = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad A \vec{e}_2 = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

Hence, the image of the unit square (spanned by  $\vec{e}_1$  and  $\vec{e}_2$ ) is the parallelogram spanned by

$$A \vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad A \vec{e}_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

In other words, every point  $(x_1, x_2)$  in the original square is mapped to

$$x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix},$$

with  $0 \leq x_1, x_2 \leq 1$ . Geometrically, this results in a parallelogram in the  $(y_1, y_2)$ -plane whose vertices are  $(0, 0)$ ,  $(1, 0)$ ,  $(1 + 2, 1)$ , and  $(2, 1)$ .



**Note:-**

Thus, restricting  $\vec{v}_i$  to a square region in the input space restricts  $\vec{v}_o$  to a parallelogram in the output space.

**Algorithm 1:** Mapping a unit square under the linear transformation  $A$ 


---

**Input:** Input vector  $\vec{v}_i = [x_1 \ x_2]^T$  with  $0 \leq x_1, x_2 \leq 1$   
**Output:** Output vector  $\vec{v}_o = [y_1 \ y_2]^T$  in the parallelogram

```

/* Matrix A: */
1  $A \leftarrow \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix};$ 
/* Apply A to input: */
2  $y_1 \leftarrow x_1 + 2x_2;$ 
3  $y_2 \leftarrow x_2;$ 
4 return  $\vec{v}_o;$ 


```

---

**1.2.1 Examples of 2D Matrix Transformations**


**Example 1.2.1** ( $A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ )

**Transformation:** Vertical stretch scaling  $x_2$  by a factor of 2.




**Example 1.2.2** ( $A = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$ )

**Transformation:** Reflection about the vertical axis.




**Example 1.2.3** ( $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ )

**Transformation:** Reflection about the line  $x_1 = x_2$ .




**Example 1.2.4** ( $A = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ )

**Transformation:** Counterclockwise rotation by  $\theta$  about the origin.



**Example 1.2.5** ( $A = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ )



**Transformation:** Projection onto the vertical axis.

## 1.3 Mapping

$$[x_1, x_2] \mapsto \text{point } (x_1, x_2).$$

Under the linear transformation given by

$$[x_1, x_2] \mapsto \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix},$$

we see that points are “stretched to the right” in the  $x_1$ -direction, while the  $x_2$  coordinate remains unchanged.

### 1.3.1 Method 1 – Vector Method


The matrix under consideration is

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}.$$

The columns of this matrix are:

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

These are the images of the standard basis vectors  $\mathbf{e}_1$  and  $\mathbf{e}_2$ , respectively.



### 1.3.2 Method 2 – Vertex Method


Consider the unit square with vertices

$$(0, 0), \quad (1, 0), \quad (1, 1), \quad (0, 1).$$

We apply  $A$  to each vertex:

$$A \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad A \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}.$$

Hence the new vertices are  $(0, 0)$ ,  $(1, 0)$ ,  $(2, 1)$ , and  $(3, 1)$ . Plotting these points yields the transformed parallelogram (the image of the original unit square).



### Definition 1.3.1: Rules for Linear Transformations

1. **Straight lines remain straight.**
2. **Parallel lines remain parallel.**
3. **Distances along lines scale in a consistent, proportional way.**

#### Justification

**1. Straight lines stay straight.** A line in parametric form is:

$$r(t) = t \mathbf{v} + \mathbf{w}.$$

Applying  $A$  gives:

$$A(r(t)) = A(t \mathbf{v} + \mathbf{w}) = t A(\mathbf{v}) + A(\mathbf{w}),$$

which is again a parametric line.


**2. Parallel lines stay parallel.** If two lines are parallel, their direction vectors are scalar multiples of each other. After applying  $A$ , the resulting direction vectors are  $A(\mathbf{v})$  for each original direction  $\mathbf{v}$ . Since  $A$  is linear, any scalar multiples remain so, preserving parallelism.

**3. Distances scale proportionally.** For two points  $\mathbf{v}_1$  and  $\mathbf{v}_2$ , the difference is  $\mathbf{d} = \mathbf{v}_2 - \mathbf{v}_1$ . Under  $A$ :

$$A\mathbf{v}_2 - A\mathbf{v}_1 = A(\mathbf{v}_2 - \mathbf{v}_1) = A(\mathbf{d}).$$

Thus the new distance  $\|\mathbf{d}'\| = \|A(\mathbf{d})\|$  is a consistent transform of  $\|\mathbf{d}\|$ , depending on the nature of  $A$ .

**Example 1.3.1** ( A matrix  $A$  that acts on a parallelogram (spanned by two vectors  $\mathbf{x}_1, \mathbf{x}_2$ ) will produce another parallelogram in the output plane. )



*Here, lines remain lines, parallels remain parallel.*

## 1.4 A System of Linear Equations

We have a matrix  $A$  of size  $m \times n$ , multiplying an unknown vector  $\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$  (which belongs to  $\mathbb{R}^n$ ), producing a result

vector  $\begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{bmatrix}$  in  $\mathbb{R}^m$ :

$$A_{m \times n} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{bmatrix}.$$

This can be viewed as a *system of  $m$  linear equations in  $n$  unknowns*. We are often interested in two main questions:

1. **Does a solution exist?** That is, can we find  $x_1, x_2, \dots, x_n$  so that  $A\mathbf{x} = \mathbf{v}$ ? Geometrically, this asks if  $\mathbf{v}$  lies in the *column space* (or *image*) of  $A$ .
2. **If at least one solution exists, is it unique or are there infinitely many?** Uniqueness is typically tied to whether the columns of  $A$  are linearly independent (and whether  $m, n$  are related in a way that gives a single solution). If there are *fewer* pivots than unknowns, or if the system is underdetermined, infinitely many solutions can occur.

*Key intuition:*

- The question of *existence* boils down to whether  $\mathbf{v}$  is in the span of the columns of  $A$ .
- The question of *uniqueness* depends on whether those columns form a set of independent vectors and on the relationship between  $m$  and  $n$ .

## 1.5 How to Determine Consistency, Uniqueness, and the Number of Solutions for a Linear System

**Note:-**

We consider a linear system

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m \end{cases}$$

and write it in *augmented matrix* form:

$$[A \mid \mathbf{b}] = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{bmatrix}.$$

- We perform *elementary row operations* (EROs) on  $[A \mid \mathbf{b}]$  to attempt to solve the system. The EROs are:

1. **Scaling (multiplying a row by a nonzero constant).**

2. **Row replacement** ( $R_i = R_i + c R_j$  for  $i \neq j$ ).

3. **Interchanging two rows** ( $R_i \leftrightarrow R_j$ ).

These operations do not change the solution set of the system.

**Row Echelon Form (REF)** We say  $[A | \mathbf{b}]$  is in *row echelon form* if:

- All rows of all zeros (if any) are at the bottom of the matrix.
- Each *pivot* (leftmost nonzero entry in a nonzero row) is strictly to the right of the pivot in the row above.

A pivot column in  $A$  corresponds to a *leading variable* (or *basic variable*), and any other column (apart from the augmented column) is called a *free column* (its corresponding variable is a *free variable*).

**Consistency & Number of Solutions**

- If, in the augmented matrix, there is a pivot in the last column (meaning a row of the form

$$[0 \ 0 \ \cdots \ 0 | c], \quad c \neq 0,$$

) then the system is *inconsistent* (no solutions). This corresponds to an equation  $0 = c$  where  $c \neq 0$ , which is impossible.

- If no such contradiction is found, then at least one solution exists (the system is *consistent*).
- The number of pivot columns in  $A$  (i.e. the number of leading variables) tells us whether solutions are unique or infinite:
  1. If the number of pivots equals the number of unknowns  $n$ , then there is exactly one solution (assuming no inconsistency).
  2. If the number of pivots is less than  $n$ , then there are free variables, implying infinitely many solutions (again, assuming no inconsistency).

**Example 1.5.1** (Example: Augmented Matrix and REF)

$$\begin{bmatrix} 1 & 2 & 3 & 5 \\ 2 & 4 & 6 & 9 \\ 3 & 6 & 9 & 15 \end{bmatrix} \longrightarrow (\text{REF}).$$

One performs row operations to get an upper-triangular or echelon form. If the last row becomes something like

$$[0 \ 0 \ 0 | c], \quad c \neq 0,$$

then there is no solution. Otherwise, we identify pivot columns, read off the relationships among variables, and find a general solution (unique or infinite).

**Definition 1.5.1: Free Columns and Basic Columns**

Once an augmented matrix is in REF, we label each pivot column (in  $A$ ) as a *basic column*, and any other column (except the last augmented column) as a *free column*.

- If  $x_j$  corresponds to a free column, we may label  $x_j$  as a parameter (e.g.  $t_1, t_2, \dots$ ).
- The variables in pivot columns can then be written in terms of these parameters.

In this manner, we get a general solution describing the entire solution set to  $A\mathbf{x} = \mathbf{b}$ .

**Algorithm to Convert  $[A | \mathbf{b}]$  to  $\text{REF}(A)$**

1. *Select a candidate row:* Choose the topmost row *among those not yet having a pivot* in which a pivot might appear.

2. *Pivot search and possible row swap*: Among this candidate row and those below it, find a row having a leftmost nonzero entry in the desired pivot column. Interchange (swap) that row with the candidate row if needed, placing a nonzero entry where your pivot should be.
3. *Declare the pivot and eliminate below*: Scale the pivot row (if desired) so that the pivot becomes 1. Then use row replacement to produce zeros below that pivot in the same column.
4. Move to the next row down and next column to the right, and repeat until you have a row echelon form.

One can then further use row replacement operations to clear the entries *above* each pivot, yielding the *reduced* row echelon form (RREF). However, for most solution purposes, REF is already sufficient to read off whether solutions exist, how many, and so on.

### 1.5.1 FRR

---

**Algorithm 2:** Forward Row Reduction (Forward Elimination)

---

**Input:** Matrix  $A$  of size  $m \times n$  and vector  $\mathbf{b}$  of size  $m$

**Output:** Upper triangular matrix  $A$  and modified vector  $\mathbf{b}$

*/\* Forward elimination process \*/*

```

1 for  $k \leftarrow 1$  to  $\min(m, n)$  do
2   for  $i \leftarrow k + 1$  to  $m$  do
3     if  $A_{kk} \neq 0$  then
4        $f \leftarrow A_{ik}/A_{kk}$ ;
5       for  $j \leftarrow k$  to  $n$  do
6          $A_{ij} \leftarrow A_{ij} - f \cdot A_{kj}$ ;
7       end
8        $b_i \leftarrow b_i - f \cdot b_k$ ;
9     end
10  end
11 end
12 return  $A, \mathbf{b}$ ;
```

---

## 1.6 Back Row Reduction

**Note:-**

Back row reduction, also known as back substitution, is a method used to solve a system of linear equations that has been transformed into an upper triangular form through Gaussian elimination. This method involves solving the equations starting from the last row and moving upwards.

### Definition 1.6.1: Back Row Reduction

Consider a system of linear equations represented in matrix form as  $A\mathbf{x} = \mathbf{b}$ , where  $A$  is an upper triangular matrix:

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}.$$

The solution is obtained by solving the last equation first and then substituting the obtained values into the preceding equations.

### Example 1.6.1 (Example of Back Row Reduction)

Consider the following upper triangular system:

$$\begin{cases} 2x_1 + 3x_2 + 4x_3 = 5, \\ 0x_1 + 6x_2 + 7x_3 = 8, \\ 0x_1 + 0x_2 + 9x_3 = 10. \end{cases}$$

We start with the last equation:

$$9x_3 = 10 \Rightarrow x_3 = \frac{10}{9}.$$

Next, we substitute  $x_3$  into the second equation:

$$6x_2 + 7\left(\frac{10}{9}\right) = 8 \Rightarrow 6x_2 + \frac{70}{9} = 8 \Rightarrow 6x_2 = 8 - \frac{70}{9} \Rightarrow x_2 = \frac{2}{9}.$$

Finally, we substitute  $x_2$  and  $x_3$  into the first equation:

$$2x_1 + 3\left(\frac{2}{9}\right) + 4\left(\frac{10}{9}\right) = 5 \Rightarrow 2x_1 + \frac{6}{9} + \frac{40}{9} = 5 \Rightarrow 2x_1 = 5 - \frac{46}{9} \Rightarrow x_1 = \frac{1}{9}.$$

Thus, the solution is:

$$x_1 = \frac{1}{9}, \quad x_2 = \frac{2}{9}, \quad x_3 = \frac{10}{9}.$$

---

**Algorithm 3:** Back Row Reduction (Back Substitution)

---

**Input:** Upper triangular matrix  $A$  of size  $n \times n$  and vector  $\mathbf{b}$  of size  $n$

**Output:** Solution vector  $\mathbf{x}$  of size  $n$

```

/* Initialize solution vector */
1 for  $i \leftarrow n$  to 1 do
2    $x_i \leftarrow b_i$ ;
3   for  $j \leftarrow i + 1$  to  $n$  do
4      $x_i \leftarrow x_i - A_{ij} \cdot x_j$ ;
5   end
6    $x_i \leftarrow x_i / A_{ii}$ ;
7 end
8 return  $\mathbf{x}$ ;

```

---

**Example 1.6.2** (Full SLE solving)

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & -1 & 0 & -2 \\ 2 & 0 & m & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & -2 & -2 & -4 \\ 0 & -2 & m-4 & -4 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & -2 & -2 & -4 \\ 0 & 0 & m-2 & 0 \end{bmatrix}$$

given  $m=0$

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & -2 & -2 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$



## Chapter 2

# Linear Combinations

### 2.1 Matrix–Vector Product as Linear Combination

$$Ax = \begin{bmatrix} 1 & -2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ 2 \end{bmatrix} = b.$$

*Intuition:* “Scaling each column of  $A$  by its corresponding entry in  $x$ .”

In other words, to compute  $Ax$ , you take  $x_1$  times the first column of  $A$  plus  $x_2$  times the second column of  $A$ .

#### Example 2.1.1 ( A Specific Choice of $x$ )

If

$$A = \begin{bmatrix} 1 & -2 \\ 3 & 2 \end{bmatrix},$$

then the first column is

$$a_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix} \quad \text{and} \quad a_2 = \begin{bmatrix} -2 \\ 2 \end{bmatrix}.$$

A linear combination of these columns is

$$x_1 \begin{bmatrix} 1 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} -2 \\ 2 \end{bmatrix} = b.$$

As an example, if  $x_1 = -\frac{1}{3}$  and  $x_2 = 1$ ,

$$-\frac{1}{3} \begin{bmatrix} 1 \\ 3 \end{bmatrix} + \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{3} - 2 \\ -1 + 2 \end{bmatrix} = \begin{bmatrix} -\frac{7}{3} \\ 1 \end{bmatrix}.$$

In the notes, a similar combination yields  $\begin{bmatrix} -2 \\ -2 \end{bmatrix}$ . The key point is that any pair  $(x_1, x_2)$  gives a vector in  $\mathbb{R}^2$ .

### Definition 2.1.1: Span

**Span** of a set of vectors is the collection of all linear combinations of those vectors. Concretely, for

$$a_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \quad a_2 = \begin{bmatrix} -2 \\ 2 \end{bmatrix},$$

we write

$$\text{Span}\{a_1, a_2\} = \left\{ x_1 a_1 + x_2 a_2 \mid x_1, x_2 \in \mathbb{R} \right\}.$$


Any vector  $b$  not in this span means  $Ax = b$  has no solution.

## 2.2 Linear Combinations and the Span

Vectors “outside” the span of  $\{a_1, a_2\}$  are precisely those  $b$  for which the system  $Ax = b$  is not consistent. Equivalently, they are not expressible as a linear combination of  $a_1$  and  $a_2$ .


## 2.3 Geometric Plots in the Notes

1. *First Plot (showing the columns  $a_1$  and  $a_2$ ):*




Any linear combination  $x_1 a_1 + x_2 a_2$  lands in the parallelogram structure (and its extensions) formed by these two vectors.

2. *Second Plot (showing a vector not in the span):*




If  $v_1$  and  $v_2$  do not cover  $(1, 1)$  by any linear combination, then  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is not in their span, hence no solution exists for  $Ax = b$ .

3. *Plot illustrating parallel vectors:*



Here,  $v_1$  and  $v_2$  are multiples of each other, so their span is just the single dashed line. The vector  $b$  off that line cannot be written as any combination of  $v_1$  and  $v_2$ .

**Example 2.3.1** ( $\vec{b}$  not in span of  $\vec{v}_1, \vec{v}_2$ )



### 2.3.1 Key Conclusions

- $\text{Span}\{a_1, a_2\}$  is the set of all vectors  $b$  for which  $Ax = b$  has a solution.
- If  $b$  is not in that span, there is no solution.
- Geometrically, if  $a_1$  and  $a_2$  are not multiples of each other, their span is a 2D plane through the origin in  $\mathbb{R}^2$ . If they are multiples, the span is just a single line, and most vectors in  $\mathbb{R}^2$  lie outside that line (no solution).
- $\text{span}a_1$  is a line or point.  $\text{span}a_1, a_2$  is a plane, line or point. and so on

### 2.3.2 Dropping a vector to retain the same span

**Question 2:** Drop vector  $a_3$  to retain the same span as  $a_1, a_2$  and  $a_3$ .

$$a_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$a_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$a_3 = \begin{bmatrix} 3 \\ 3 \end{bmatrix}$$

**Solution:** yes,  $\text{span} a_1, a_2, a_3 = \text{span} a_1, a_2$  because  $a_3$  is a linear combination of  $a_1$  and  $a_2$  ( $a_1 + a_2 a_3$ )

## 2.4 Evaluating Linear Dependence Procedurally

**Example 2.4.1** ( Finding linear dependence of a set of vectors. )

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 5 \\ 2 \\ -1 \end{bmatrix}$$

Since  $v_3$  is a linear combination of  $v_1$  and  $v_2$ , we can drop  $v_3$  to retain the same span as  $v_1$  and  $v_2$ .

$$\text{span} v_1, v_2, v_3 = \text{span} v_1, v_2$$

As

$$v_3 = 2v_1 - v_2$$

**Note:-**

Working towards a procedure for evaluating linear dependence of a set of vectors.

$$-v_1 + 2v_2 - v_3 = 0$$

$$\begin{bmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

We know that one solution is

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ -1 \end{bmatrix}$$

Since this SLE has one particular solution as well as the trivial one, then it must have infinitely many solutions.

Then if  $A\vec{x} = 0$  has infinitely many solutions, then the set of vectors (columns) in  $A$  is linearly dependent.

Conversely, if  $A\vec{x} = 0$  has only one solution ( $\vec{x} = \vec{0}$ ), then the set of vectors (columns) in  $A$  is linearly independent.

### Definition 2.4.1: Linear Independence

We say a set of vectors  $\{v_1, v_2, \dots, v_n\}$  is *linearly independent* if the only solution to the SLE

$$\begin{bmatrix} \vec{v}_1 & \vec{v}_2 & \dots & \vec{v}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

is the trivial solution, i.e.  $\vec{x} = \vec{0}$ .

$$A' = \begin{bmatrix} 1 & 3 & 5 & 0 \\ 2 & 2 & 2 & 0 \\ 3 & 1 & -1 & 0 \end{bmatrix} \xrightarrow{R_2=R_2-2R_1, \quad R_3=R_3-3R_1} \begin{bmatrix} 1 & 3 & 5 & 0 \\ 0 & -4 & -8 & 0 \\ 0 & -8 & -16 & 0 \end{bmatrix} \xrightarrow{R_3=R_3-2R_2} \begin{bmatrix} 1 & 3 & 5 & 0 \\ 0 & -4 & -8 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \text{REF}(A')$$

1 Free variable  $\implies$  Linearly Independent

$$\text{REF}(A') \xrightarrow{R_2=R_2/(-4)} \begin{bmatrix} 1 & 3 & 5 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1=R_1-3R_2} \begin{bmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \text{RREF}(A') \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

Then

$$[\vec{v}_1 \quad \vec{v}_2 \quad \vec{v}_3] t_3 \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

And therefore the linear combination in its homogeneous form implies

$$t_3(\vec{v}_1 - 2\vec{v}_2 + \vec{v}_3) = \vec{0}$$

## 2.5 Identity Matrix

### Definition 2.5.1: Identity Matrix

special Square matrix where the diagonal elements are 1 and the off-diagonal elements are 0.

$$I_n = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Note that  $I_n \vec{x} = \vec{x}$  and  $I'_n = I_n$  and therefore is called a **symmetric matrix**

### Question 3: Matrix Multiplication that gives Identity matrix

$$A = \begin{bmatrix} 1 & 0.7 \\ 0 & 1 \end{bmatrix}$$

Find a matrix  $B$  such that

$$AB = I$$

**Solution:** Since  $A$  makes a horizontal shear of 0.7 then  $B$  should do the opposite

$$\therefore B = \begin{bmatrix} 1 & -0.7 \\ 0 & 1 \end{bmatrix} = A^{-1}$$

### Question 4: Matrix Multiplication that gives Identity matrix 2

$$A = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

Find a matrix  $B$  such that

$$AB = I$$

**Solution:** Since  $A$  makes a reflection about the  $x_2$  axis,  $B$  must do the same

$$\therefore B = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} = A^{-1}$$

**Question 5: Matrix Multiplication that gives Identity matrix**

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

Find a matrix  $B$  such that

$$AB = I$$

**Solution:**  $A$  flattens all vertical components. Geometrically speaking,  $A^{-1}$  doesn't exist, as shown here

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a & b \\ 0 & 0 \end{bmatrix}$$

Observe that  $\text{span}A$  is the line  $x_2 = 0$

**Definition 2.5.2: General Case for inverses of square matrices**

For a matrix  $A$  which is  $n \times n$ ,  $A^{-1}$  exists if and only if

$$\text{span}\{A\} = \mathbb{R}^n$$

**Definition 2.5.3: Finding  $A^{-1}$**

Get the Augmented matrix

$$A'' = [A | I]$$

where  $A, I$  are in the form  $n \times n$  and then perform RREF on this augmented matrix. Once done, you will have a matrix in the form

$$\text{RREF}(A'') = [I | A^{-1}]$$

**Example 2.5.1 (Finding  $A^{-1}$ )**

$$A = \begin{bmatrix} 1 & 0.7 \\ 0 & 1 \end{bmatrix}$$

$$A'' = \begin{bmatrix} 1 & 0.7 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

Then RREF

$$\xrightarrow{R_1 = R_1 - 0.7R_2} \begin{bmatrix} 1 & 0 & 1 & -0.7 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 1 & -0.7 \\ 0 & 1 \end{bmatrix}$$

**Example 2.5.2 (Finding  $A^{-1}$ )**

$$A = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 2 & 1 \\ 3 & 1 & 3 \end{bmatrix}$$

Check if  $\text{span}\{A\} = \mathbb{R}^3$

$$\begin{array}{l} R_2 = R_2 - 2R_1 \\ R_3 = R_3 - 3R_1 \end{array} \rightarrow \begin{bmatrix} 1 & 3 & 2 \\ 0 & -4 & -3 \\ 0 & -8 & -3 \end{bmatrix} \xrightarrow{R_3 = R_3 - 2R_2} \begin{bmatrix} \textcircled{1} & 3 & 2 \\ 0 & \textcircled{-4} & -3 \\ 0 & 0 & \textcircled{3} \end{bmatrix}$$

Here  $A$  has 3 pivots, so it is invertible. Next, make the augmented matrix  $A'' = [A \mid I]$  and perform RREF

$$\begin{bmatrix} 1 & 3 & 2 & 1 & 0 & 0 \\ 2 & 2 & 1 & 0 & 1 & 0 \\ 3 & 1 & 3 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{array}{l} R_2 = R_2 - 2R_1 \\ R_3 = R_3 - 3R_1 \end{array} \rightarrow \begin{bmatrix} 1 & 3 & 2 & 1 & 0 & 0 \\ 0 & -4 & -3 & -2 & 1 & 0 \\ 0 & -8 & -3 & -3 & 0 & 1 \end{bmatrix} \xrightarrow{R_3 = R_3 - 2R_2} \begin{bmatrix} 1 & 3 & 2 & 1 & 0 & 0 \\ 0 & -4 & -3 & -2 & 1 & 0 \\ 0 & 0 & 3 & 1 & -2 & 1 \end{bmatrix}$$

$$\xrightarrow{R_2 = -\frac{1}{4}R_2} \begin{bmatrix} 1 & 3 & 2 & 1 & 0 & 0 \\ 0 & 1 & \frac{3}{4} & \frac{1}{2} & -\frac{1}{4} & 0 \\ 0 & 0 & 3 & 1 & -2 & 1 \end{bmatrix} \xrightarrow{R_1 = R_1 - 3R_2} \begin{bmatrix} 1 & 0 & -\frac{1}{4} & -\frac{1}{2} & \frac{3}{4} & 0 \\ 0 & 1 & \frac{3}{4} & \frac{1}{2} & -\frac{1}{4} & 0 \\ 0 & 0 & 3 & 1 & -2 & 1 \end{bmatrix}$$

$$R_1 = R_1 + \frac{1}{4}R_3$$


$$\xrightarrow{R_3 = \frac{1}{3}R_3} \begin{bmatrix} 1 & 0 & -\frac{1}{4} & -\frac{1}{2} & \frac{3}{4} & 0 \\ 0 & 1 & \frac{3}{4} & \frac{1}{2} & -\frac{1}{4} & 0 \\ 0 & 0 & 1 & \frac{1}{3} & -\frac{2}{3} & \frac{1}{3} \end{bmatrix} \xrightarrow{R_2 = R_2 - \frac{3}{4}R_3} \begin{bmatrix} 1 & 0 & 0 & -\frac{5}{12} & \frac{7}{12} & \frac{1}{12} \\ 0 & 1 & 0 & \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ 0 & 0 & 1 & \frac{1}{3} & -\frac{2}{3} & \frac{1}{3} \end{bmatrix}.$$

Therefore


$$A^{-1} = \begin{bmatrix} -\frac{5}{12} & \frac{7}{12} & \frac{1}{12} \\ \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{1}{3} & -\frac{2}{3} & \frac{1}{3} \end{bmatrix}$$

### Example 2.5.3 (Review)

$$A(5,5), \quad B\left(\frac{5}{2}, \frac{5}{2}\right), \quad C\left(\frac{15}{2}, \frac{5}{2}\right), \quad D(5,0)$$



$$A'(0,0), \quad B'\left(-\frac{5}{2}, -\frac{5}{2}\right), \quad C'\left(\frac{5}{2}, -\frac{5}{2}\right), \quad D'(0,-5)$$



$$\vec{AB} = \vec{v}_2 = \begin{bmatrix} -\frac{5}{2} \\ -\frac{5}{2} \end{bmatrix}, \vec{AC} = \vec{v}_1 = \begin{bmatrix} \frac{5}{2} \\ -\frac{5}{2} \end{bmatrix}$$

$$\vec{v}_1^T \vec{v}_2 = -\frac{25}{4} + \frac{25}{4} = 0 = |\vec{v}_1| |\vec{v}_2| \cos(\theta)$$

$$\therefore \theta = 90^\circ$$

### 2.5.1 ERP's and Identity Matrices

Let  $A''$  be a matrix. Consider the augmented matrix:

$$[A \mid I] : m \times 2m$$

Perform a sequence of elementary row operations (ERPs) on  $A$  to turn it into the identity matrix  $I$ :

$$E_p \dots E_2 E_1 A'' = B[A \mid I] = [BA \mid BI]$$

where  $B$  is an  $m \times m$  matrix.

If we achieve  $BA = I$ , then:

$$B = A^{-1}$$

#### Question 6: Thinklet

Consider a set  $\mathcal{S}$  of vectors in  $\mathbb{R}^m$  who's tips form a straight line through the origin.  
If we linearly combine a set of vectors in  $\mathcal{S}$ , is the result guaranteed to be in  $\mathcal{S}$ ?

**Solution:** DUH



# Chapter 3

## Vector Spaces

### Definition 3.0.1: Vector Space

A vector space is a set of vectors in  $\mathbb{R}^m$  for which the following hold:

1. The set is closed under addition: that is, if  $\vec{v}_1, \vec{v}_2$  are both in the set, then  $\vec{v}_1 + \vec{v}_2$  is also in the set.
2. The set is closed under scalar multiplication: that is, if  $\vec{v}$  is in the set and  $c \in \mathbb{R}$ , then  $c\vec{v}$  is also in the set.

### Definition 3.0.2: Basis of a Vector Space

A basis for a vector space in  $\mathbb{R}^m$  is a set of independent vectors that span the vector space. The maximum number of vectors that could be in that basis is  $m$ .  
 $m$  is the **dimension** of the vector space.

### Example 3.0.1 (Examples of Vector Spaces)

$\mathcal{S}$	Basis	Dimension
$\{\vec{0}\}$	$\emptyset$	0
$\left\{ \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{bmatrix} t \right\}$	$\left\{ \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{bmatrix} \right\}$	1
$\left\{ \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{bmatrix} t_1 + \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} t_2 \right\}$	$\left\{ \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{bmatrix}, \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix} \right\}$	2
$\left\{ \sum_{i=1}^k \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{mi} \end{bmatrix} t_i \right\}$	$\left\{ \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}, \begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}, \dots, \begin{bmatrix} a_{1k} \\ a_{2k} \\ \vdots \\ a_{mk} \end{bmatrix} \right\}$	$k$

## 3.1 Column Space

### Definition 3.1.1: Column space of matrix $A$

The *column space* of an  $m \times n$  matrix  $A$ , denoted as  $\text{Col}(A)$ , is the subspace of  $\mathbb{R}^m$  spanned by the columns of  $A$ . Formally,

$$\text{Col}(A) = \text{span}(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$$

where  $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$  are the column vectors of  $A$ . This means that  $\text{Col}(A)$  consists of all possible linear combinations of the columns of  $A$ .

## 3.2 Linear Combinations and Column Space

### 3.2.1 Basic Definitions and Concepts

#### Definition 3.2.1: Matrix Notation

An  $m \times n$  matrix  $A$  is a rectangular array of numbers with  $m$  rows and  $n$  columns. We can represent  $A$  as a collection of column vectors:

$$A = [\vec{v}_1 \ \vec{v}_2 \ \dots \ \vec{v}_n]$$

where each  $\vec{v}_i$  is a vector in  $\mathbb{R}^m$ .

#### Definition 3.2.2: Column Space (Col $A$ )

The column space of a matrix  $A$ , denoted as  $\text{Col } A$ , is the set of all linear combinations of the columns of  $A$ . Formally, if  $A = [\vec{v}_1 \ \vec{v}_2 \ \dots \ \vec{v}_n]$ , then

$$\text{Col } A = \text{span}\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\} = \{t_1\vec{v}_1 + t_2\vec{v}_2 + \dots + t_n\vec{v}_n \mid t_1, t_2, \dots, t_n \in \mathbb{R}\}$$

### 3.2.2 Dimension and Basis of the Column Space

#### Question 7: What is the dimension of $\text{Col } A$ ?

The dimension of  $\text{Col } A$  is equal to the number of pivot columns in the reduced row echelon form (REF) of  $A$ .

**Solution:** The dimension corresponds to the number of linearly independent columns in matrix  $A$ .

#### Question 8: What is the basis for $\text{Col } A$ ?

The basis for  $\text{Col } A$  is formed by the column vectors of the **original** matrix  $A$  that correspond to the pivot columns in the REF of  $A$ . These are referred to as the "basic columns."

**Solution:**

1. Find  $\text{REF}(A)$ .
2. Identify the pivot columns in  $\text{REF}(A)$ .
3. Select the corresponding columns from the **original\*** matrix **A**. These columns form the basis.

### 3.2.3 Parametric Form of the Column Space


#### Question 9: What is the parametric form of Col A?

If the basis for Col A is  $\{\vec{w}_1, \vec{w}_2, \dots, \vec{w}_k\}$ , then the parametric form of any vector  $\vec{v}$  in Col A is given by:

$$\vec{v} = t_1\vec{w}_1 + t_2\vec{w}_2 + \dots + t_k\vec{w}_k$$

where  $t_1, t_2, \dots, t_k$  are scalar parameters (real numbers).

**Solution:** This expresses any vector in the column space as a linear combination of the basis vectors.



### 3.2.4 Geometry of the Column Space

#### Question 10: What is the geometry of Col A?

The geometric interpretation of Col A depends on its dimension:

- If  $\dim(\text{Col } A) = 0$ , Col A is a point (the origin,  $\vec{0}$ ).
- If  $\dim(\text{Col } A) = 1$ , Col A is a line through the origin.
- If  $\dim(\text{Col } A) = 2$ , Col A is a plane through the origin.
- If  $\dim(\text{Col } A) = 3$ , Col A is a volume (all of  $\mathbb{R}^3$ ) through the origin.
- In general, if A is an  $m \times n$  matrix, and  $\dim(\text{Col } A) = k$ , Col A is a k-dimensional subspace of  $\mathbb{R}^m$ .

**Solution:** The dimension indicates the "degrees of freedom" or the number of independent directions spanned by the column vectors.

### 3.2.5 Example

#### Example 3.2.1 (Example Calculation)

Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 2 & 4 & 6 \end{bmatrix}$$

Find the REF of A, the dimension of Col A, a basis for Col A, the parametric form of Col A, and the geometry of Col A.

**Solution:** First, find the reduced row echelon form (REF) of A:

$$\text{REF}(A) = \begin{bmatrix} 1 & 0 & -1 \\ 0 & -4 & -8 \\ 0 & 0 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

- **Dimension of Col A:** The REF has two pivot columns (the first and second columns). Therefore,  $\dim(\text{Col } A) = 2$ .

- **Basis for Col A:** The pivot columns in  $\text{REF}(A)$  are the first and second columns. We take the corresponding columns from the \*original\* matrix  $A$  to form the basis:

$$\text{Basis}(\text{Col } A) = \left\{ \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 4 \end{bmatrix} \right\}$$

- **Parametric Form of Col A:** Any vector  $\vec{v}$  in  $\text{Col } A$  can be written as a linear combination of the basis vectors:

$$\vec{v} = t_1 \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix} + t_2 \begin{bmatrix} 2 \\ 2 \\ 4 \end{bmatrix}$$

where  $t_1$  and  $t_2$  are any real numbers.

- **Geometry of Col A:** Since  $\dim(\text{Col } A) = 2$ ,  $\text{Col } A$  is a plane in  $\mathbb{R}^3$  that passes through the origin.

### 3.3 Nullspace of a Matrix

#### Definition 3.3.1: Nullspace

The nullspace of a matrix  $A$ , denoted by  $\text{Nul } A$ , is the set of all solutions to the homogeneous equation  $A\vec{x} = \vec{0}$ .

#### Note:-

The solution space for  $A\vec{x} = \vec{0}$  is a vector space. The nullspace of  $A$  resides in  $\mathbb{R}^n$ , where  $n$  is the number of columns of  $A$ .

#### Question 11: Dimension of Nullspace

What is the dimension of  $\text{Nul } A$ ?

**Solution:** The dimension of  $\text{Nul } A$  is equal to the number of free columns in the reduced row echelon form (RREF) of  $A$ . We write this as  $\dim(\text{Nul } A) = \# \text{ of free columns in RREF}(A)$ .

#### Question 12: Basis for Nullspace

What is the basis for  $\text{Nul } A$ ?

**Solution:** To find the basis for  $\text{Nul } A$ , we first find the RREF of  $A$ . If the solution space of  $A\vec{x} = \vec{0}$  is given by

$$\vec{x} = t_1 \vec{w}_1 + t_2 \vec{w}_2 + \cdots + t_k \vec{w}_k$$

where  $k$  is the number of free columns in  $\text{RREF}(A)$ , and  $t_1, t_2, \dots, t_k$  are arbitrary scalars, then the basis of  $\text{Nul } A$  is the set  $\{\vec{w}_1, \vec{w}_2, \dots, \vec{w}_k\}$ .

#### Question 13: Parametric Form of Nullspace

What is the parametric form of  $\text{Nul } A$ ?

**Solution:** The parametric form of  $\text{Nul } A$  is given by the linear combination of the basis vectors:

$$\vec{v} = t_1 \vec{w}_1 + t_2 \vec{w}_2 + \cdots + t_k \vec{w}_k$$

where  $\vec{w}_1, \vec{w}_2, \dots, \vec{w}_k$  are the basis vector for the nullspace.

**Example 3.3.1** (Example)

Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 2 & 4 & 6 \end{bmatrix}$$

Find the nullspace, its dimension, basis, and the geometric interpretation.

First, find the RREF of  $A$ :

$$\text{REF}(A) = \begin{bmatrix} 1 & 2 & 3 \\ 0 & -4 & -8 \\ 0 & 0 & 0 \end{bmatrix}$$

Further reducing to RREF:

$$\text{RREF}(A) = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

- The first column corresponds to a basic variable.
- The second column corresponds to a basic variable.
- The third column corresponds to a free variable.

The dimension of the nullspace is the number of free variables:

$$\text{Dim}(\text{N}(A)) = 1$$

To find the solution space for  $A\vec{x} = \vec{0}$ , we use the RREF:

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ . The RREF gives us the following equations:

$$\begin{aligned} x_1 - x_3 &= 0 \\ x_2 + 2x_3 &= 0 \end{aligned}$$


$x_3$  is free. Let  $x_3 = t_3$ . Then  $x_1 = t_3$  and  $x_2 = -2t_3$ . So,

$$\vec{x} = \begin{bmatrix} t_3 \\ -2t_3 \\ t_3 \end{bmatrix} = t_3 \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$$

Therefore, the basis for  $\text{Nul } A$  is:

$$\text{Basis}(\text{Nul } A) = \left\{ \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix} \right\}$$

The geometry of  $\text{Nul } A$  is a line in  $\mathbb{R}^3$  passing through the origin and in the direction of the vector  $\begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix}$ .



### 3.4 Row Space of a Matrix

#### Definition 3.4.1: Row Space

The row space of a matrix  $A$ , denoted as  $\text{Row } A$ , is defined as the column space of the transpose of  $A$ , i.e.,  $\text{Row } A = \text{Col } A^T$ .


Let's consider a matrix  $A$  and its transpose  $A^T$ :

$$A = \begin{bmatrix} - & r_1 & - \\ - & r_2 & - \\ & \vdots & \\ - & r_m & - \end{bmatrix} \quad A^T = \begin{bmatrix} | & | & & | \\ r_1^T & r_2^T & \dots & r_m^T \\ | & | & & | \end{bmatrix}$$

Where  $r_1, r_2, \dots, r_m$  represent the rows of matrix  $A$ . The rows of  $A$  become the columns of  $A^T$ .

#### Note:-

The row vectors  $r_1, r_2, \dots, r_m$  live in  $\mathbb{R}^n$ .



### 3.5 Dimension and Basis of Row Space

#### Question 14: What is the dimension of Row $A$ ?

What is the dimension of the row space of a matrix  $A$ ?

**Solution:-** The dimension of  $\text{Row } A$  is equal to the number of pivots in the row echelon form (REF) of  $A$ . It is also equal to the number of pivots in the REF of  $A^T$ .

$$\text{Dim}(\text{Row } A) = \# \text{ pivots in REF}(A) = \# \text{ pivots in REF}(A^T)$$

#### Question 15: What is the Basis of Row $A$

How to find the basis for the Row Space of  $A$

**Solution:**

1. Identify all the rows of  $\text{REF}(A)$  that have pivots.
2. The basis vectors for Row  $A$  are:
  - Vectors corresponding to rows (with pivots in  $\text{REF}(A)$ ) of  $A$ .

OR

- Vectors corresponding to rows (with pivots in  $\text{REF}(A)$ ) of  $\text{REF}(A)$ .

OR

- Vectors corresponding to rows (with pivots in  $\text{REF}(A)$ ) of  $\text{RREF}(A)$ .

**Example 3.5.1** (Example: Finding the Basis and Dimension)

Let

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 1 & 2 & 3 \end{bmatrix}$$

Find the dimension and a basis for Row  $A$ .

**Solution:** First, find the REF of  $A$ :

$$\text{REF}(A) = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

The REF of  $A$  has one pivot (in the first row).

Dimension of Row  $A$ :

$$\text{Dim}(\text{Row } A) = 1$$

Basis for Row  $A$ :

- Using rows of  $A$  with pivots in  $\text{REF}(A)$ :  $[1 \ 2 \ 3]$
- Using rows of  $\text{REF}(A)$  with pivots:  $[1 \ 2 \ 3]$

---

**Algorithm 4:** Finding Dimension and Basis of Row Space

---

**Input:** Matrix  $A$

**Output:** Dimension of Row  $A$ , Basis of Row  $A$

*/\* Compute the Row Echelon Form (REF) of A \*/*

1  $\text{REF\_A} \leftarrow \text{REF}(A)$ ;

*/\* Count the number of pivots in REF(A) \*/*

2  $\text{num\_pivots} \leftarrow$  Number of pivots in  $\text{REF\_A}$ ;

*/\* The dimension of Row A is the number of pivots \*/*

3  $\text{dim\_row\_A} \leftarrow \text{num\_pivots}$ ;

*/\* Initialize an empty set to store basis vectors \*/*

4  $\text{basis\_row\_A} \leftarrow \emptyset$ ;

*/\* Identify rows with pivots in REF(A) \*/*

5 **foreach** row  $r$  in  $\text{REF\_A}$  **do**

6     **if** row  $r$  has a pivot **then**

*/\* Add corresponding row from original matrix A to basis \*/*

7         Let  $r'$  be the corresponding row in matrix  $A$ .  $\text{basis\_row\_A} \leftarrow \text{basis\_row\_A} \cup \{r'\}$ ;

8     **end**

9 **end**

10 **return**  $\text{dim\_row\_A}$ ,  $\text{basis\_row\_A}$ ;

---

# Chapter 4

## Properties of Matrices

### 4.1 Determinant of a Square Matrix

Consider a square Matrix

$$A : m \times m$$

If

$$A^{-1} \text{ exists} \Leftrightarrow \dim(\text{Col}(A)) = \dim(\text{Row}(A)) = m$$

$$\therefore \text{Rank}(A) = m \text{ and } \text{Nul}(A) = \{0\}$$

And most importantly, since the columns of A are linearly independent,

$$\det(\text{ref}(A)) \neq 0$$

To make  $\text{REF}(A)$  unique for  $A$ , we require that in constructing  $\text{REF}(A)$ , we follow three rules:

1. Use the “REPLACE” ERP freely.
2. When using the “SWAP” ERP, apply the “SCALE” ERP to one of the rows by  $-1$ .
3. Don’t use the “SCALE” ERP for any other purpose than the one in Rule 2.

#### 4.1.1 Cofactor Algorithm for Computing Determinant

Given a square matrix  $A = [a_{ij}]$  of size  $n \times n$ , the determinant can be computed by expanding along any row  $i$  (where  $1 \leq i \leq n$ ) as follows:

1. **Base Case:** If  $n = 1$ , then  $\det(A) = a_{11}$ .
2. **Recursive Case:** For  $n > 1$ :

$$\det(A) = \sum_{j=1}^n (-1)^{i+j} a_{ij} \det(M_{ij}),$$

where:

- $M_{ij}$  is the **minor matrix** obtained by deleting row  $i$  and column  $j$  from  $A$ ,
- $(-1)^{i+j} \det(M_{ij})$  is the **cofactor** of the element  $a_{ij}$ .

**Steps:**

- Choose a row  $i$  (commonly the first row,  $i = 1$ , but any row works).
- For each element  $a_{ij}$  in row  $i$ :
  - Compute the minor  $M_{ij}$ .



- Recursively calculate  $\det(M_{ij})$ .
- Multiply  $a_{ij}$ ,  $(-1)^{i+j}$ , and  $\det(M_{ij})$ .
- Sum the results for all columns  $j$  in row  $i$ .

Say  $B$  which has the Form

$$\begin{bmatrix} a_1 & a_2 & \dots \\ b_1 & b_2 & \dots \end{bmatrix}$$

And delimits an area in  $\mathbb{R}^2$ , then the absolute value of the determinant of a matrix  $A$  which applies a transformation to  $B$  gives the scaling of the area of  $B$

### 4.1.2 Properties of Determinants

- $\det(A \cdot B) = \det(A) \cdot \det(B)$
- $\det(A^{-1}) = \frac{1}{\det(A)}$
- $\det(A^T) = \det(A)$
- $\det(I) = 1$
- $\det(E_{\text{REPLACE}} \cdot A) = \det(A)$
- $\det(E_{\text{SCALE}=\alpha} \cdot A) = \alpha \cdot \det(A)$
- $\det(A) = 0 \Leftrightarrow \det(\text{REF}(A)) = 0$

## 4.2 EigenVectors and EigenValues

Given  $A \in \mathbb{R}^{n \times n}$

$$A\vec{x} = \vec{0} \leftarrow \vec{x} \in \text{Null}(A)$$

Can rewrite As

$$A\vec{x} = 0\vec{x}$$

Generalise

$$A\vec{x} = \lambda\vec{x}$$

where the solution for  $\vec{x}$  also defines a vector space, called an **EigenSpace**, where any vector in this space is called an **EigenVector**, and has a corresponding value of  $\lambda$  called an **EigenValue**.

#### Example 4.2.1 (Eigenvectors)

$$A = \begin{bmatrix} -2 & 0 \\ 0 & 3 \end{bmatrix}$$

is  $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$  an eigenvector for  $A$ ?

$$\begin{bmatrix} -2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \end{bmatrix} = \begin{bmatrix} -4 \\ 6 \end{bmatrix} = \lambda \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

No such Lambda, therefore not eigenvector, but

$$\begin{bmatrix} -2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \end{bmatrix} = \begin{bmatrix} -4 \\ 0 \end{bmatrix} = -2 \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

So  $\lambda = -2$  is the EigenValue for the EigenVector that is  $\begin{bmatrix} 2 \\ 0 \end{bmatrix}$

# Eigenvectors and Eigenvalues

## Definition 4.2.1: Eigenvector and Eigenvalue

Let  $A$  be an  $n \times n$  matrix. A non-zero vector  $\vec{x}$  is an eigenvector of  $A$  if there exists a scalar  $\lambda$  (called an eigenvalue) such that:

$$A\vec{x} = \lambda\vec{x}$$


### Note:-

The geometric interpretation of  $A\vec{x} = \lambda\vec{x}$  is that applying the transformation represented by  $A$  to the eigenvector  $\vec{x}$  results in a vector that is simply a scaled version of  $\vec{x}$ . The direction is either preserved (if  $\lambda > 0$ ), reversed (if  $\lambda < 0$ ), or the vector becomes the zero vector (if  $\lambda = 0$ ). Crucially,  $\vec{x}$  \*cannot\* be the zero vector, but  $\lambda$  can be.

### Example 4.2.2 (Rotation Matrix)

Consider a rotation matrix  $A$  in 2D that rotates vectors by 45 degrees counterclockwise:

$$A = \begin{bmatrix} \cos(45^\circ) & -\sin(45^\circ) \\ \sin(45^\circ) & \cos(45^\circ) \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} \end{bmatrix}$$




This matrix  $A$  has \*no\* real eigenvectors (and thus no real eigenvalues). Applying  $A$  to any non-zero vector  $\vec{x}$  will \*always\* rotate it; there's no non-zero vector that will simply be scaled by  $A$ . The resulting vector  $A\vec{x}$  will never have the same (or opposite) direction as the original  $\vec{x}$ .

### Example 4.2.3 (Horizontal Shear and Stretch)

Consider the matrix:

$$A = \begin{bmatrix} 2 & 0.5 \\ 0 & 1 \end{bmatrix}$$

This matrix represents a horizontal shear and stretch. Let's visualize its effect on a unit square:



- Vectors along the  $x_1$ -axis (of the form  $\alpha \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ) are stretched by a factor of 2, but their direction remains unchanged.
- Vectors along the  $x_2$ -axis (of the form  $\alpha \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ ) are sheared horizontally by 0.5 units, but their vertical

component is unchanged.

- Consider a general vector  $\begin{bmatrix} \alpha \\ \beta \end{bmatrix}$ . Applying the transformation yields  $A \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \begin{bmatrix} 2\alpha + 0.5\beta \\ \beta \end{bmatrix}$ .

Let's find the eigenvectors and eigenvalues for this matrix.

- For vectors along the  $x_1$  axis:

$$A \begin{bmatrix} \alpha \\ 0 \end{bmatrix} = \begin{bmatrix} 2 & 0.5 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \alpha \\ 0 \end{bmatrix} = \begin{bmatrix} 2\alpha \\ 0 \end{bmatrix} = 2 \begin{bmatrix} \alpha \\ 0 \end{bmatrix}$$

So,  $\begin{bmatrix} \alpha \\ 0 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 2$ .

- For vectors along the  $x_2$  axis, we might \*think\* they are eigenvectors, but let's check:

$$A \begin{bmatrix} 0 \\ \alpha \end{bmatrix} = \begin{bmatrix} 2 & 0.5 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ \alpha \end{bmatrix} = \begin{bmatrix} 0.5\alpha \\ \alpha \end{bmatrix} = \alpha \begin{bmatrix} 0.5 \\ 1 \end{bmatrix}$$

This is \*not\* a scalar multiple of the original vector  $\begin{bmatrix} 0 \\ \alpha \end{bmatrix}$ . So, vectors along the  $x_2$  axis are \*not\* eigenvectors.

- Now consider the case where the transformed vector is  $\alpha$  times the original:

$$\begin{bmatrix} 2 & 0.5 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -0.5\alpha \\ \alpha \end{bmatrix} = \begin{bmatrix} -1\alpha + 0.5\alpha \\ \alpha \end{bmatrix} = \begin{bmatrix} -0.5\alpha \\ \alpha \end{bmatrix} = 1 * \begin{bmatrix} -0.5\alpha \\ \alpha \end{bmatrix}$$

So  $\begin{bmatrix} -0.5\alpha \\ \alpha \end{bmatrix}$  is an eigenvector corresponding to eigenvalue 1.

## Finding Eigenvalues and Eigenvectors

### Theorem 4.2.1 Finding Eigenvalues

To find the eigenvalues of a matrix  $A$ , we solve the characteristic equation:

$$\det(A - \lambda I) = 0$$

where  $I$  is the identity matrix of the same size as  $A$ .

*Proof.* We start with the definition  $A\vec{x} = \lambda\vec{x}$ . Rearranging, we get:

$$A\vec{x} - \lambda\vec{x} = \vec{0}$$

$$A\vec{x} - \lambda I\vec{x} = \vec{0}$$

$$(A - \lambda I)\vec{x} = \vec{0}$$

Let  $B = A - \lambda I$ . We have  $B\vec{x} = \vec{0}$ . We are looking for non-zero vectors  $\vec{x}$  that satisfy this equation.

- If  $B$  is invertible (i.e.,  $\det(B) \neq 0$ ), then the only solution is  $\vec{x} = \vec{0}$ . But eigenvectors are \*non-zero\* by definition. Therefore, we must have  $\det(B) = 0$ .
- If  $\det(B) = 0$ , then  $B$  is \*not\* invertible, and the system  $B\vec{x} = \vec{0}$  has infinitely many solutions (i.e., a non-trivial null space). These non-zero solutions are the eigenvectors.

Therefore, to find the eigenvalues  $\lambda$ , we must solve  $\det(A - \lambda I) = 0$ . ☺

**Example 4.2.4** (Finding Eigenvalues (Continued))

Let's go back to our example:

$$A = \begin{bmatrix} 2 & 0.5 \\ 0 & 1 \end{bmatrix}$$

We form  $B = A - \lambda I$ :

$$B = \begin{bmatrix} 2 & 0.5 \\ 0 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 - \lambda & 0.5 \\ 0 & 1 - \lambda \end{bmatrix}$$

Now we find the determinant of  $B$ :

$$\det(B) = (2 - \lambda)(1 - \lambda) - (0.5)(0) = (2 - \lambda)(1 - \lambda)$$

Setting  $\det(B) = 0$ , we get:

$$(2 - \lambda)(1 - \lambda) = 0$$

This gives us the eigenvalues  $\lambda_1 = 2$  and  $\lambda_2 = 1$ .

**Finding Eigenvectors****Algorithm 5:** Finding Eigenvalues and Eigenvectors

**Input:** A matrix  $A$

**Output:** Eigenvalues and corresponding eigenvectors of  $A$

- 1 Calculate the characteristic polynomial:  $\det(A - \lambda I)$ ;
- 2 Solve the characteristic equation  $\det(A - \lambda I) = 0$  to find the eigenvalues  $\lambda_i$ ;
- 3 **for** each eigenvalue  $\lambda_i$  **do**
- 4     Form the matrix  $B_i = A - \lambda_i I$ ;
- 5     Solve the homogeneous system  $B_i \vec{x} = \vec{0}$  to find the eigenvectors corresponding to  $\lambda_i$ ;
- 6     The solution set will typically involve free variables. Express the eigenvectors in terms of these free variables.;
- 7 **end**
- 8 **return** The eigenvalues  $\lambda_i$  and their corresponding eigenvectors.

**Example 4.2.5** (Finding Eigenvectors (Continued))

We found the eigenvalues  $\lambda_1 = 2$  and  $\lambda_2 = 1$  for the matrix  $A = \begin{bmatrix} 2 & 0.5 \\ 0 & 1 \end{bmatrix}$ . Now let's find the eigenvectors.

- For  $\lambda_1 = 2$ :

$$B_1 = A - 2I = \begin{bmatrix} 2 - 2 & 0.5 \\ 0 & 1 - 2 \end{bmatrix} = \begin{bmatrix} 0 & 0.5 \\ 0 & -1 \end{bmatrix}$$

We solve  $B_1 \vec{x} = \vec{0}$ :

$$\begin{bmatrix} 0 & 0.5 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Row reducing (although not strictly necessary here, we do it for demonstration):

$$\begin{bmatrix} 0 & 0.5 \\ 0 & -1 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 \\ 0 & -1 \end{bmatrix} \rightarrow \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

This gives us  $x_2 = 0$ , and  $x_1$  is free. Let  $x_1 = t_1$ . Then the eigenvector is:

$$\vec{x} = \begin{bmatrix} t_1 \\ 0 \end{bmatrix} = t_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

- For  $\lambda_2 = 1$ :

$$B_2 = A - I = \begin{bmatrix} 2-1 & 0.5 \\ 0 & 1-1 \end{bmatrix} = \begin{bmatrix} 1 & 0.5 \\ 0 & 0 \end{bmatrix}$$

We solve  $B_2 \vec{x} = \vec{0}$ :

$$\begin{bmatrix} 1 & 0.5 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

This system is already in row-echelon form. We have  $x_1 + 0.5x_2 = 0$ , so  $x_1 = -0.5x_2$ .  $x_2$  is free. Let  $x_2 = t_2$ . Then  $x_1 = -0.5t_2$ , and the eigenvector is:

$$\vec{x} = \begin{bmatrix} -0.5t_2 \\ t_2 \end{bmatrix} = t_2 \begin{bmatrix} -0.5 \\ 1 \end{bmatrix}$$

### 4.3 Properties of Eigenvectors

#### Definition 4.3.1: Dimension of eigenspace

When an eigenvalue repeats  $k$  times,

$$1 \leq \dim(\text{eigenspace for that eigenvalue}) \leq k$$

**Example 4.3.1** (Example 1 of this property)

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A - \lambda I = \begin{bmatrix} 1-\lambda & 0 & 0 \\ 0 & 1-\lambda & 0 \\ 0 & 0 & 1-\lambda \end{bmatrix}$$

$$\det(A - \lambda I) = (1 - \lambda)^3$$

$$\lambda_1 = 1, \lambda_2 = 1, \lambda_3 = 1, k = 3$$

$$\lambda_1 \text{ repeats } 3 (= k) \text{ times}$$

And the eigenspace for  $\lambda_1 = 1$  is

$$A - \lambda_1 I = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

therefore the eigenspace is

$$\left\{ t_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + t_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + t_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} : t_1, t_2, t_3 \in \mathbb{R} \right\}$$

**Example 4.3.2** (Example 2 of this property)

$$A = \begin{bmatrix} 1 & 0.5 \\ 0 & 1 \end{bmatrix}$$

$$\det(A - \lambda I) = (1 - \lambda)^3$$

$$(1 - \lambda)^2 - (0.5)(0) = 0$$

$$\lambda_1 = 1, \lambda_2 = 1, k = 2$$

Therefore the eigenspace is

$$A - \lambda_1 I = \begin{bmatrix} 0 & 0.5 \\ 0 & 0 \end{bmatrix} \xrightarrow{R_1=2R_1} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Therefore the eigenspace for  $\lambda = 1$  is

$$\left\{ t_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} : t_1 \in \mathbb{R} \right\}$$

$$\therefore \dim(\text{eigenspace for } \lambda = 1) = 1 \leq k = 2$$

#### Definition 4.3.2: Dominant Eigenvector

An eigenvector whose eigenvalue (among all eigenvalues for some  $A : m \times m$ ) has the largest magnitude ( $|\lambda|$ ) gives the direction along which an initial blob of points is stretched the most.

#### Example 4.3.3 (Dominant Eigenvector Example)

A=

$$\begin{bmatrix} 2 & 4 \\ 3 & 3 \end{bmatrix}$$

$$\det(A - \lambda I) = (2 - \lambda)(3 - \lambda) - 4 * 3 = 0$$

$$\therefore \lambda_1 = 6, \lambda_2 = -1$$

Eigenspace for largest magnitude of eigenvalue  $\lambda_1 = 6$

$$A = \begin{bmatrix} -4 & 4 \\ 3 & -3 \end{bmatrix} \xrightarrow{R_2=R_2+\frac{3}{4}R_1} \begin{bmatrix} -4 & 4 \\ 0 & 0 \end{bmatrix} \xrightarrow{R_1=-\frac{1}{4}R_1} \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix}$$

$$\therefore \vec{v}_1 = t_2 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$


and the eigenspace for  $\lambda_1 = 6$  is

$$\left\{ t \begin{bmatrix} 1 \\ 1 \end{bmatrix} : t \in \mathbb{R} \right\}$$

Now apply the transformation to a set of points (unit square)

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 4 & 3 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 4 & 3 \end{bmatrix}$$

Visually represented this looks like this



**Example 4.3.4** (Dominant Eigenvector Example 2)

Consider

$$A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}.$$

The eigenvalues come from

$$\det(A - \lambda I) = (3 - \lambda)(2 - \lambda) = 0,$$

so  $\lambda_1 = 3$  and  $\lambda_2 = 2$ . The dominant eigenvalue is 3. To find its eigenvector, solve

$$(A - 3I)\vec{v} = \begin{bmatrix} 0 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \vec{0}.$$

The first row gives  $v_2 = 0$ ; choosing  $v_1 = 1$  yields

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

This indicates that the largest stretch occurs in the horizontal direction.

**Transformation on a set of points:**

Consider the original set of points

$$\mathcal{P} = \{(x, y) \mid -1 \leq x \leq 1, -0.5 \leq y \leq 0.5\}.$$

Under the transformation  $A$ , each point  $(x, y)$  is mapped to

$$(x', y') = (3x + y, 2y).$$

The four corners transform as follows:

- $(-1, -0.5)$  maps to  $(3(-1) + (-0.5), 2(-0.5)) = (-3.5, -1)$ .
- $(1, -0.5)$  maps to  $(3(1) + (-0.5), 2(-0.5)) = (2.5, -1)$ .


- $(1, 0.5)$  maps to  $(3(1) + (0.5), 2(0.5)) = (3.5, 1)$ .
- $(-1, 0.5)$  maps to  $(3(-1) + (0.5), 2(0.5)) = (-2.5, 1)$ .

Thus, the image of  $\mathcal{P}$  is the quadrilateral

$$\mathcal{P}' = \{(x', y')\}$$

with vertices at  $(-3.5, -1)$ ,  $(2.5, -1)$ ,  $(3.5, 1)$ , and  $(-2.5, 1)$ .

**Diagram:**



#### Definition 4.3.3: Eigenvalues of Triangular Matrices

The elements on the main diagonal of a triangular matrix are the eigenvalues of that matrix.

**Example 4.3.5** (EigenValues of Triangular matrices example)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}$$

$$\det(A - \lambda I) = (1 - \lambda)(4 - \lambda)(6 - \lambda)$$

$$\lambda_1 = 1, \lambda_2 = 4, \lambda_3 = 6$$

$$\text{Eigenvalues of } A = \{1, 4, 6\}$$

#### Definition 4.3.4: Number of Distinct EigenValues

The number of distinct eigenvalues of a matrix  $A : m \times m$  is  $\leq m$



**Definition 4.3.5: Eigenvalues and Determinant**

For a matrix  $A : m \times m$

$$\det(A) = \lambda_1 \cdot \lambda_2 \cdot \dots \cdot \lambda_m$$

Proof:

$$\det(A - \lambda I) = \pm \lambda^m + a_{m-1} \lambda^{m-1} + \dots + a_1 \lambda + a_0$$

where the  $\pm$  follows

$$\begin{cases} +1, & m \% 2 = 0 \\ -1, & m \% 2 = 1 \end{cases}$$

$$\therefore |A - \lambda I| = \begin{cases} \lambda^m - a_{m-1} \lambda^{m-1} + \dots - a_1 \lambda + a_0, & m \% 2 = 0 \\ -\lambda^m + a_{m-1} \lambda^{m-1} - \dots + a_1 \lambda - a_0, & m \% 2 = 1 \end{cases}$$

Set  $\lambda = 0$

$$|A| = \begin{cases} \lambda_1, \lambda_2, \dots, \lambda_m, & m \% 2 = 0 \\ \lambda_1, \lambda_2, \dots, \lambda_m, & m \% 2 = 1 \end{cases}$$

**Definition 4.3.6: Non-invertible Square matrix**

For a non-invertible (singular) square matrix  $A : m \times m$ , the determinant is 0, therefore at least one eigenvalue is 0.

**Definition 4.3.7: Spanning Eigenvectors**

The eigenvectors of  $A$  span all of  $\mathbb{R}^m$  if and only if  $A$  has  $m$  distinct eigenvalues. If an eigenvalue repeats  $k$  times, the dimension of the corresponding space is  $k$ .