Slow eccentrics compared to fast eccentrics in resistance training

Bibliography:

Certainty assessment							Summary of findings				
Participants (studies) Follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)	Anticipated absolute effects	
							With fast eccentrics	With slow eccentrics		Risk with fast eccentrics	Risk difference with slow eccentrics
Maximal strength (follow-up: range 4 weeks to 12 weeks; assessed with: 1RM)											
171 (8 RCTs)	not serious	serious[a]	not serious	very serious[b]	all plausible residual confounding would reduce the demonstrated effect	⨁⨁◯◯ Low			-		**0** (0 to 0)
Muscle hypertrophy (follow-up: range 6 weeks to 12 weeks; assessed with: US/MRI)											
100 (5 RCTs)	not serious	serious[c]	not serious	very serious[b]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)
Countermovement Jump Height (follow-up: range 4 weeks to 6 weeks; assessed with: Vertec/Force Plate/MyoJump)											
74 (3 RCTs)	not serious	not serious	not serious	very serious[b]	strong association	⨁⨁⨁◯ Moderate			-		**0** (0 to 0)
STR - Trained Subgroup (follow-up: range 4 weeks to 12)											
0 (RCTs)	not serious	not serious	not serious	serious[d]	none	⨁⨁⨁◯ Moderate			-		**0** (0 to 0)
STR - Untrained Subgroup (follow-up: range 6 weeks to 12 weeks)											
0 (RCTs)	not serious	serious[c]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)
STR - Multi-joint Subgroup (follow-up: range 4 weeks to 12 weeks)											

Slow eccentrics compared to fast eccentrics in resistance training

Bibliography:

Certainty assessment							Summary of findings				
0 (RCTs)	not serious	serious[c]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

STR - Single-joint Subgroup (follow-up: range 7 weeks to 12 weeks)

0 (RCTs)	not serious	serious[c]	not serious	very serious[b]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

STR - Sets-to-Failure Subgroup (follow-up: range 6 weeks to 12 weeks)

0 (RCTs)	not serious	very serious[f]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

STR - Volume-Load Subgroup (follow-up: range 4 weeks to 12 weeks)

0 (RCTs)	not serious	not serious	not serious	serious[d]	none	⨁⨁⨁◯ Moderate			-		**0** (0 to 0)

STR - Within-Subjects Subgroup (follow-up: range 8 weeks to 8 weeks)

0 (RCTs)	not serious	not serious	not serious	extremely serious[b]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

STR - Between-Subjects Subgroup (follow-up: range 4 weeks to 12 weeks)

0 (RCTs)	not serious	serious[c]	not serious	extremely serious[b]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

HYP - Trained Subgroup (follow-up: range 8 weeks to 12 weeks)

0 (RCTs)	serious[g]	serious[c]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

HYP - Untrained Subgroup (follow-up: range 6 weeks to 8 weeks)

0 (RCTs)	not serious	serious[c]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

HYP - Arm Subgroup (follow-up: range 7 to 12)

Slow eccentrics compared to fast eccentrics in resistance training

Bibliography:

Certainty assessment							Summary of findings				
0 (RCTs)	serious[g]	serious[c]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

HYP - Thigh Subgroup (follow-up: range 6 weeks to 8 weeks)

0 (RCTs)	not serious	serious[c]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

HYP - Within-Subjects Subgroup (follow-up: range 8 weeks to 8 weeks)

0 (RCTs)	not serious	not serious	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

HYP - Between-Subjects Subgroup (follow-up: range 6 weeks to 12 weeks)

0 (RCTs)	not serious	very serious[f]	not serious	extremely serious[e]	none	⨁◯◯◯ Very low			-		**0** (0 to 0)

CI: confidence interval; **SMD:** standardised mean difference

Explanations

a. Serious inconsistency (moderate heterogeneity I2 > 50% and point estimates vary considerably), that may be explained by training status.
b. Very serious imprecision (95% crosses both equivalence bounds)
c. Serious unexplained inconsistency (point estimates vary considerably).
d. Serious imprecision (95% CI crosses one equivalence bound).
e. Extremely serious imprecision (95% CI crosses both equivalence bounds with large effects in both directions).
f. Very serious unexplained inconsistency (large point estimates in each direction).
g. Serious risk of bias due to allocation concealment and lack of blinding.