Outcome (*n*, *k*)	**Hedge's *g* (95% CI) [95% PI]**	***p*-value**
Strength		
Pooled (*n = 166, k = 9*) *Follow-up: 4-12 weeks*	0.25 (-0.27, 0.78) [-1.13, 1.63]	0.30
Trained (*n = 77, k = 4*) *Follow-up: 4-12 weeks*	0.33 (-0.05, 0.72) [-0.05, 0.72]	0.06
Untrained (*n = 89, k = 5*) *Follow-up: 6-12 weeks*	0.15 (-1.08, 1.39) [-2.66, 2.96]	0.75
Multi-joint (*n = 111, k = 5*) *Follow-up: 4-12 weeks*	0.08 (-0.97, 1.14) [-2.23, 2.40]	0.83
Single-joint (*n = 74, k = 4*) *Follow-up: 7-12 weeks*	0.40 (-0.47, 1.27) [-1.10, 1.90]	0.23
Sets-to-failure (*n = 82, k = 5*) *Follow-up: 7-12 weeks*	0.26 (-1.03, 1.55) [-2.64, 3.16]	0.60
Volume-load (*n = 84, k = 4*) *Follow-up: 4-12 weeks*	0.25 (-0.04, 0.53) [-0.04, 0.53]	0.07
Between-participants (*n = 143, k = 7*) *Follow-up: 4-12 weeks*	0.31 (-0.47, 1.09) [-1.60, 2.23]	0.37
Within-participants (*n = 23, k = 2*) *Follow-up: 8 weeks*	0.11 (-1.17, 1.39) [-1.17, 1.39]	0.47
Hypertrophy		
Pooled (*n = 95, k = 6*) *Follow-up: 6-12 weeks*	0.05 (-0.31, 0.41) [-0.73, 0.83]	0.71
Trained (*n = 25, k = 2*) *Follow-up: 8-12 weeks*	0.46 (8.20, 9.13) [-13.54, 14.47]	0.62

Untrained (*n = 70, k = 4*) *Follow-up: 6-8 weeks*	0.02 (-0.50, 0.55) [-1.08, 1.12]	0.90
Arm (*n = 32, k = 2*) *Follow-up: 7-12 weeks*	0.29 (-1.39, 1.97) [-2.53, 3.11]	0.51
Thigh (*n = 63, k = 4*) *Follow-up: 6-8 weeks*	-0.01 (-0.85, 0.84) [-1.61, 1.59]	0.98
Between-participants (*n = 72, k = 4*) *Follow-up: 6-12 weeks*	0.15 (-0.86, 1.16) [-1.85, 2.15]	0.65
Within-participants (*n = 23, k = 2*) *Follow-up: 8 weeks*	0.00 (-1.66, 1.67) [-1.84, 1.85]	0.98
CMJ		
Pooled (*n = 74, k = 3*) Follow-up: 4-6 weeks	-0.73 (-1.64, 0.17) [-1.64, 0.17]	0.07

SF2 – Analysis results using the traditional 95% NHST interpretive lens. This table presents the 95% confidence and prediction intervals, along with p-values, following conventional NHST relative to the null.