Slow eccentrics compared to fast eccentrics in resistance training

Bibliography:

Certainty assessment							Summary of findings				
Participants (studies) Follow-up	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Overall certainty of evidence	Study event rates (%)		Relative effect (95% CI)	Anticipated absolute effects	
							With fast eccentrics	With slow eccentrics		Risk with fast eccentrics	Risk difference with slow eccentrics
Maximal strength (follow-up: range 4 weeks to 12 weeks; assessed with: 1RM)											
0 (9 RCTs)	serious[a]	serious[b]	not serious	serious[c]	all plausible residual confounding would reduce the demonstrated effect	⨁⨁◯◯ Low[a,b,c]			-	-	SMD **0.18 SD higher** (0.38 lower to 74 higher)
Muscle hypertrophy (follow-up: range 6 weeks to 12 weeks; assessed with: US/MRI)											
0 (6 RCTs)	not serious	serious[d]	not serious	serious[c]	none	⨁⨁◯◯ Low[c,d]			-	-	SMD **0.03 SD higher** (0.43 lower to 0.48 higher)
Countermovement Jump Height (follow-up: range 4 weeks to 6 weeks; assessed with: Vertec/Force Plate/MyoJump)											
74 (3 RCTs)	not serious	not serious	not serious	serious[c]	strong association	⨁⨁⨁⨁ High[c]	32	42	-	-	SMD **0.73 SD lower** (1.64 lower to 0.17 lower)
STR - Trained Subgroup (follow-up: range 4 weeks to 12)											
0 (RCTs)	not serious	not serious	not serious	serious[e]	none	⨁⨁⨁◯ Moderate[e]			-		**0** (0 to 0)
STR - Untrained Subgroup (follow-up: range 6 weeks to 12 weeks)											
0 (RCTs)	not serious	serious[d]	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[d,f]			-		**0** (0 to 0)
STR - Multi-joint Subgroup (follow-up: range 4 weeks to 12 weeks)											

Slow eccentrics compared to fast eccentrics in resistance training

Bibliography:

Certainty assessment							Summary of findings				
0 (RCTs)	not serious	serious[d]	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[d,f]			-		**0** (0 to 0)

STR - Single-joint Subgroup (follow-up: range 7 weeks to 12 weeks)

0 (RCTs)	not serious	serious[d]	not serious	serious[c]	none	⨁⨁◯◯ Low[c,d]			-		**0** (0 to 0)

STR - Sets-to-Failure Subgroup (follow-up: range 6 weeks to 12 weeks)

0 (RCTs)	serious[a]	very serious[g]	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[a,f,g]			-		**0** (0 to 0)

STR - Volume-Load Subgroup (follow-up: range 4 weeks to 12 weeks)

0 (RCTs)	not serious	not serious	not serious	serious[e]	none	⨁⨁⨁◯ Moderate[e]			-		**0** (0 to 0)

STR - Within-Subjects Subgroup (follow-up: range 8 weeks to 8 weeks)

0 (RCTs)	not serious	not serious	not serious	extremely serious[c]	none	⨁◯◯◯ Very low[c]			-		**0** (0 to 0)

STR - Between-Subjects Subgroup (follow-up: range 4 weeks to 12 weeks)

0 (RCTs)	serious[a]	serious[d]	not serious	very serious[c]	none	⨁◯◯◯ Very low[a,c,d]			-		**0** (0 to 0)

HYP - Trained Subgroup (follow-up: range 8 weeks to 12 weeks)

0 (RCTs)	serious[a]	serious[d]	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[a,d,f]			-		**0** (0 to 0)

HYP - Untrained Subgroup (follow-up: range 6 weeks to 8 weeks)

0 (RCTs)	serious[a]	serious[d]	not serious	very serious[f]	none	⨁◯◯◯ Very low[a,d,f]			-		**0** (0 to 0)

Slow eccentrics compared to fast eccentrics in resistance training

Bibliography:

Certainty assessment							Summary of findings				
HYP - Arm Subgroup (follow-up: range 7 to 12)											
0 (RCTs)	serious[a]	serious[d]	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[a,d,f]			-		**0** (0 to 0)
HYP - Thigh Subgroup (follow-up: range 6 weeks to 8 weeks)											
0 (RCTs)	serious[a]	serious[d]	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[a,d,f]			-		**0** (0 to 0)
HYP - Within-Subjects Subgroup (follow-up: range 8 weeks to 8 weeks)											
0 (RCTs)	not serious	not serious	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[f]			-		**0** (0 to 0)
HYP - Between-Subjects Subgroup (follow-up: range 6 weeks to 12 weeks)											
0 (RCTs)	serious[a]	very serious[g]	not serious	extremely serious[f]	none	⨁◯◯◯ Very low[a,f,g]			-		**0** (0 to 0)

CI: confidence interval; **SMD:** standardised mean difference

Explanations

a. Serious risk of bias due to allocation concealment and lack of blinding.
b. Serious inconsistency (moderate heterogeneity I2 > 50% and point estimates vary considerably), that may be explained by training status.
c. Very serious imprecision (95% crosses both equivalence bounds)
d. Serious unexplained inconsistency (point estimates vary considerably).
e. Serious imprecision (95% CI crosses one equivalence bound).
f. Extremely serious imprecision (95% CI crosses both equivalence bounds with large effects in both directions).
g. Very serious unexplained inconsistency (large point estimates in each direction).