

## Seasonal Function OLS

Average temperature  $t$ -th day:

$$T_t = \frac{T_{\max_t} + T_{\min_t}}{2}$$

following an Ornstein–Uhlenbeck (OU) process:

$$dT_t = a(T_m - T_t) dt + \sigma_t dW_t$$

Seasonality with a time-dependent mean that accounts for this seasonality implies -

$$\frac{dT_t^m}{dt} = B + \omega \cdot C \cdot \cos(\omega t + \phi)$$

SDE -

$$dT_t = \left\{ \frac{dT_t^m}{dt} + a(T_t^m - T_t) \right\} dt + \sigma_t dW_t$$

Distribution of differences  $\Delta T_t = T_t - T_{t-1}$  has mean and variance :

$$\begin{aligned} \overline{\Delta T_t} &= \frac{1}{n} \sum_{t=1}^n \Delta T_t \\ \frac{1}{n} \sum_{t=1}^n (\Delta T_t - \overline{\Delta T_t})^2. \end{aligned}$$

OLS to be fitted -

$$T_t^m = a_1 + a_2 t + a_3 \sin(\omega t) + a_4 \cos(\omega t)$$

where:

- $T_t^m$ : Modeled temperature at time  $t$
- $\omega$ : Angular frequency, typically set to  $\frac{2\pi}{365}$  for daily data with an annual cycle

such that -

$$T_t^m = A + Bt + C \sin(\omega t + \phi)$$

- $A = a_1$ ,  $B = a_2$ ,  $C = \sqrt{a_3^2 + a_4^2}$
- $\phi = \arctan\left(\frac{a_4}{a_3}\right) - \pi$