

# The American Monsoon System: variability and teleconnections

Jorge Luis García Franco

Wadham College  
University of Oxford

*A thesis submitted for the degree of  
Doctor of Philosophy*

Michaelmas 2020

## Abstract

Your abstract text goes here. Check your departmental regulations, but generally this should be less than 300 words. See the beginning of Chapter 4 for more.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Pellentesque sit amet nibh volutpat, scelerisque nibh a, vehicula neque. Integer placerat nulla massa, et vestibulum velit dignissim id. Ut eget nisi elementum, consectetur nibh in, condimentum velit. Quisque sodales dui ut tempus mattis. Duis malesuada arcu at ligula egestas egestas. Phasellus interdum odio at sapien fringilla scelerisque. Mauris sagittis eleifend sapien, sit amet laoreet felis mollis quis. Pellentesque dui ante, finibus eget blandit sit amet, tincidunt eu neque. Vivamus rutrum dapibus ligula, ut imperdiet lectus tincidunt ac. Pellentesque ac lorem sed diam egestas lobortis.

Suspendisse leo purus, efficitur mattis urna a, maximus molestie nisl. Aenean porta semper tortor a vestibulum. Suspendisse viverra facilisis lorem, non pretium erat lacinia a. Vestibulum tempus, quam vitae placerat porta, magna risus euismod purus, in viverra lorem dui at metus. Sed ac sollicitudin nunc. In maximus ipsum nunc, placerat maximus tortor gravida varius. Suspendisse pretium, lorem at porttitor rhoncus, nulla urna condimentum tortor, sed suscipit nisi metus ac risus.

Aenean sit amet enim quis lorem tristique commodo vitae ut lorem. Duis vel tincidunt lacus. Sed massa velit, lacinia sed posuere vitae, malesuada vel ante. Praesent a rhoncus leo. Etiam sed rutrum enim. Pellentesque lobortis elementum augue, at suscipit justo malesuada at. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Praesent rhoncus convallis ex. Etiam commodo nunc ex, non consequat diam consectetur ut. Pellentesque vitae est nec enim interdum dapibus. Donec dapibus purus ipsum, eget tincidunt ex gravida eget. Donec luctus nisi eu fringilla mollis. Donec eget lobortis diam.

Suspendisse finibus placerat dolor. Etiam ornare elementum ex ut vehicula. Donec accumsan mattis erat. Quisque cursus fringilla diam, eget placerat neque bibendum eu. Ut faucibus dui vitae dolor porta, at elementum ipsum semper. Sed ultrices dui non arcu pellentesque placerat. Etiam posuere malesuada turpis, nec malesuada tellus malesuada.

# The American Monsoon System: variability and teleconnections



Jorge Luis García Franco  
Wadham College  
University of Oxford

A thesis submitted for the degree of  
*Doctor of Philosophy*

Michaelmas 2020

# Acknowledgements

## Personal

This is where you thank your advisor, colleagues, and family and friends.

  Lorem ipsum dolor sit amet, consectetur adipiscing elit. Vestibulum feugiat et est at accumsan. Praesent sed elit mattis, congue mi sed, porta ipsum. In non ullamcorper lacus. Quisque volutpat tempus ligula ac ultricies. Nam sed erat feugiat, elementum dolor sed, elementum neque. Aliquam eu iaculis est, a sollicitudin augue. Cras id lorem vel purus posuere tempor. Proin tincidunt, sapien non dictum aliquam, ex odio ornare mauris, ultrices viverra nisi magna in lacus. Fusce aliquet molestie massa, ut fringilla purus rutrum consectetur. Nam non nunc tincidunt, rutrum dui sit amet, ornare nunc. Donec cursus tortor vel odio molestie dignissim. Vivamus id mi erat. Duis porttitor diam tempor rutrum porttitor. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Sed condimentum venenatis consectetur. Lorem ipsum dolor sit amet, consectetur adipiscing elit.

  Aenean sit amet lectus nec tellus viverra ultrices vitae commodo nunc. Mauris at maximus arcu. Aliquam varius congue orci et ultrices. In non ipsum vel est scelerisque efficitur in at augue. Nullam rhoncus orci velit. Duis ultricies accumsan feugiat. Etiam consectetur ornare velit et eleifend.

  Suspendisse sed enim lacinia, pharetra neque ac, ultricies urna. Phasellus sit amet cursus purus. Quisque non odio libero. Etiam iaculis odio a ex volutpat, eget pulvinar augue mollis. Mauris nibh lorem, mollis quis semper quis, consequat nec metus. Etiam dolor mi, cursus a ipsum aliquam, eleifend venenatis ipsum. Maecenas tempus, nibh eget scelerisque feugiat, leo nibh lobortis diam, id laoreet purus dolor eu mauris. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Nulla eget tortor eu arcu sagittis euismod fermentum id neque. In sit amet justo ligula. Donec rutrum ex a aliquet egestas.

## Institutional

If you want to separate out your thanks for funding and institutional support, I don't think there's any rule against it. Of course, you could also just remove the subsections and do one big traditional acknowledgement section.

  Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut luctus tempor ex at pretium. Sed varius, mauris at dapibus lobortis, elit purus tempor neque, facilisis sollicitudin felis nunc a urna. Morbi mattis ante non augue blandit pulvinar. Quisque nec euismod mauris. Nulla et tellus eu nibh auctor malesuada quis imperdiet quam. Sed eget tincidunt velit. Cras molestie sem ipsum, at faucibus quam mattis vel. Quisque vel placerat orci, id tempor urna. Vivamus mollis, neque in

aliquam consequat, dui sem volutpat lorem, sit amet tempor ipsum felis eget ante. Integer lacinia nulla vitae felis vulputate, at tincidunt ligula maximus. Aenean venenatis dolor ante, euismod ultrices nibh mollis ac. Ut malesuada aliquam urna, ac interdum magna malesuada posuere.

# Abstract

Your abstract text goes here. Check your departmental regulations, but generally this should be less than 300 words. See the beginning of Chapter 4 for more.

  Lorem ipsum dolor sit amet, consectetur adipiscing elit. Pellentesque sit amet nibh volutpat, scelerisque nibh a, vehicula neque. Integer placerat nulla massa, et vestibulum velit dignissim id. Ut eget nisi elementum, consectetur nibh in, condimentum velit. Quisque sodales dui ut tempus mattis. Duis malesuada arcu at ligula egestas egestas. Phasellus interdum odio at sapien fringilla scelerisque. Mauris sagittis eleifend sapien, sit amet laoreet felis mollis quis. Pellentesque dui ante, finibus eget blandit sit amet, tincidunt eu neque. Vivamus rutrum dapibus ligula, ut imperdiet lectus tincidunt ac. Pellentesque ac lorem sed diam egestas lobortis.

  Suspendisse leo purus, efficitur mattis urna a, maximus molestie nisl. Aenean porta semper tortor a vestibulum. Suspendisse viverra facilisis lorem, non pretium erat lacinia a. Vestibulum tempus, quam vitae placerat porta, magna risus euismod purus, in viverra lorem dui at metus. Sed ac sollicitudin nunc. In maximus ipsum nunc, placerat maximus tortor gravida varius. Suspendisse pretium, lorem at porttitor rhoncus, nulla urna condimentum tortor, sed suscipit nisi metus ac risus.

  Aenean sit amet enim quis lorem tristique commodo vitae ut lorem. Duis vel tincidunt lacus. Sed massa velit, lacinia sed posuere vitae, malesuada vel ante. Praesent a rhoncus leo. Etiam sed rutrum enim. Pellentesque lobortis elementum augue, at suscipit justo malesuada at. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Praesent rhoncus convallis ex. Etiam commodo nunc ex, non consequat diam consectetur ut. Pellentesque vitae est nec enim interdum dapibus. Donec dapibus purus ipsum, eget tincidunt ex gravida eget. Donec luctus nisi eu fringilla mollis. Donec eget lobortis diam.

  Suspendisse finibus placerat dolor. Etiam ornare elementum ex ut vehicula. Donec accumsan mattis erat. Quisque cursus fringilla diam, eget placerat neque bibendum eu. Ut faucibus dui vitae dolor porta, at elementum ipsum semper. Sed ultrices dui non arcu pellentesque placerat. Etiam posuere malesuada turpis, nec malesuada tellus malesuada.

# Contents

<b>List of Figures</b>	<b>vi</b>
<b>List of Abbreviations</b>	<b>viii</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Motivation . . . . .	1
1.2 Contribution . . . . .	1
<b>2 Background</b>	<b>3</b>
2.1 The tropical circulation and the global monsoon . . . . .	3
2.2 The American Monsoon System . . . . .	5
2.3 Review of the Midsummer drought . . . . .	6
2.4 El Niño Southern Oscillation: impacts to the American monsoon system . . . . .	9
2.5 Stratosphere-Troposphere Coupling in the Tropics . . . . .	11
<b>3 Data and methods</b>	<b>12</b>
3.1 Observations and reanalysis data . . . . .	12
3.2 Model data . . . . .	14
<b>4 On the dynamical and thermodynamical mechanisms of the MSD in the Met Office CMIP6 models</b>	<b>17</b>
4.1 Climatological features . . . . .	17
4.1.1 On the mechanisms of the MSD in the UK Met Office models . . . . .	20
4.1.2 Summary and discussion . . . . .	27
<b>Appendices</b>	
<b>A Review of Cardiac Physiology and Electrophysiology</b>	<b>33</b>
A.1 Anatomy . . . . .	33
A.2 Mechanical Cycle . . . . .	36
A.3 Electrical Cycle . . . . .	37
A.4 Cellular Electromechanical Coupling . . . . .	38
<b>Bibliography</b>	<b>40</b>

# List of Figures

2.1	a) Climatological mean annual rainfall rates in the tropics in the TRMM dataset (1999-2018). b) The mean rainfall rate difference between boreal summer (JJAS) and austral summer (DJFM). The red contours highlight the regions where the summer rainfall amount accounts for more than 55% of the total annual rainfall accumulation.	4
2.2	Four-chamber illustration of the human heart.	10
4.1	Pentad-mean precipitation in (a) southern Mexico and northern Central America and (b) Cuba. Shading shows uncertainty obtained by bootstrapping the interannual variability of the TRMM dataset.	18
4.2	(a, d, g) Climatological JJAS rainfall and the difference between (b, e, h) the midsummer drought and the first peak periods and (c, f, i) between the second peak and the midsummer drought periods for (a-c) TRMM, (d-f) ERA5 and (g-i) CHIRPS.	19
4.3	(a, d, g, j) JJAS model bias compared to TRMM and the difference between (b, e, h, k) the midsummer drought and the first peak periods and (c, f, i, l) between the second peak and the midsummer drought periods for four different simulations.	20
4.4	Pentad-mean seasonal cycle of indices associated with the MSD in Central America and the Caribbean.	21
4.5	Out-going longwave radiation (OLR) [ $\text{W m}^{-2}$ ] (shaded) and $\omega$ 500-hPa [ $10^{-2} \text{ Pa s}^{-1}$ ] (line contours) differences between the MSD and MSDO and the MSDE and MSD.	23
4.6	As in Figure 4.5 but showing wind vectors at the 850 hPa level.	24
4.7	As in Figure 4.5 but the anomalies are shown for SSTs [K] (contours) and incoming shortwave radiation [ $\text{W m}^{-2}$ ] at the surface (line-contours). Incoming shortwave is defined such as negative differences imply less incoming shortwave and positive anomalies represent more incoming shortwave at the surface.	25
4.8	As in Figure 4.5 but showing in shading, moisture flux divergence $\nabla \cdot \vec{u}q$ at the 850 hPa level with units of $10^{-7} \text{ s}^{-1} \text{ kg / kg}$ and zonal wind anomalies (line contours) in $\text{m s}^{-1}$ .	26

4.9 Scatter plot of the (a, b) area-averaged precipitation over land (Box in Figure ??) during the different stages of the MSD. (c) scatter of the East Pacific SSTs against the precipitation over land during the first peak period. (d-l) show the scatter differences in several variables between the different stages of onset of the MSD (MSDO), the drier MSD and the end of the MSDE. The differences are shown for area-averaged quantities in the East Pacific (EP), the Caribbean Sea (CSEA) and overland (MSD) as above. The units for  $\Delta U$  are [ $m s^{-1}$ ],  $\Delta MFC$  [ $10^{-11} s^{-1}$ ],  $\Delta SW$  and  $\Delta LHF$  [ $W m^{-2}$ ] and  $\Delta pr$   $mm d^{-1}$ . The Pearson correlation coefficient for the 38 yr of reanalysis or observations ( $r_{obs}$ ) is shown for each panel. 28

## List of Abbreviations

- 1-D, 2-D** . . . One- or two-dimensional, referring in this thesis to spatial dimensions in an image.
- Otter** . . . . . One of the finest of water mammals.
- Hedgehog** . . . Quite a nice prickly friend.

*Neque porro quisquam est qui dolorem ipsum quia dolor sit amet, consectetur, adipisci velit...*

*There is no one who loves pain itself, who seeks after it and wants to have it, simply because it is pain...*

— Cicero's *de Finibus Bonorum et Malorum*

# 1

## Introduction

### Contents

---

<b>1.1 Motivation</b>	1
<b>1.2 Contribution</b>	1

---

### 1.1 Motivation

The American Monsoon System (AMS) provides the majority of rainfall for the large regions in Latin America and southwestern United States. Climate variability and teleconnections to this monsoon system can impact the population through changes in extreme precipitation, the timings of the monsoon or the overall rainfall during the rainy or the dry seasons causing floods or droughts.

General circulation models (GCMs) have been used to provide climate projections of future climate in the AMS. However, GCMs may also be used to understand physical mechanisms associated with climate variability and teleconnections.

This thesis focuses on the American Monsoon System and the outstanding questions regarding the climate variability and teleconnections affecting this monsoon.

### 1.2 Contribution

Chapter 3 evaluates two state-of-the-art CMIP6 models for their representation of the monsoon system. In general, the models show a good representation of the seasonal cycle as they are

able to simulated detail aspects such as the Midsummer drought. ENSO teleconnections in these models appear to be non-linear, as are the observations. Chapter 4 provides a method that is able to better characterise the MSD timings and strengths, as a way of analysing the mechanisms of the MSD in observations and models, analysis that is done in chapter 5. The Quasi-biennial Oscillation is proposed to be responsible for the different ENSO teleconnections shown in chapter 3 and are thus further explored using modelling experiments in chapter 6.

*Alles Gescheite ist schon gedacht worden.  
Man muss nur versuchen, es noch einmal zu denken.  
All intelligent thoughts have already been thought;  
what is necessary is only to try to think them again.*

— Johann Wolfgang von Goethe ?

# 2

## Background

### Contents

---


2.1	The tropical circulation and the global monsoon . . . . .	3
2.2	The American Monsoon System . . . . .	5
2.3	Review of the Midsummer drought . . . . .	6
2.4	El Niño Southern Oscillation: impacts to the American monsoon system . . . . .	9
2.5	Stratosphere-Troposphere Coupling in the Tropics . . . . .	11

---

This chapter summarises the main aspects of the tropical circulation and of the global monsoon. Then, the relevant background of the characteristics of American monsoon system and outstanding issues is given.

### 2.1 The tropical circulation and the global monsoon

Tropical climate is characterized by the strong incoming solar insolation year-round which makes the tropics the warmest region of the planet. The differential heating between the tropics and higher latitudes drives a meridional transport of energy by the atmosphere. The tropical oceans that receive such a strong solar insolation exhibit strong evaporative fluxes. This in combination with the differential heating of land over ocean modulate the tropical circulation. This tropical circulation can be described to a first order through the zonal and meridional circulations known as the Hadley and Walker cells.



**Figure 2.1:** a) Climatological mean annual rainfall rates in the tropics in the TRMM dataset (1999–2018). b) The mean rainfall rate difference between boreal summer (JJAS) and austral summer (DJFM). The red contours highlight the regions where the summer rainfall amount accounts for more than 55% of the total annual rainfall accumulation.

The Hadley cell is the meridional overturning circulation that arises from the differential heating between the tropics and the midlatitudes. The Hadley cell is characterized by ascending motions in the tropics and descending motions in the subtropics, and acts to transport heat poleward from the equator (Lorenz, 1967). The boreal summer Hadley cell, for instance, is primarily a result of ascent in the Indian Ocean and the west Pacific regions with a minor contribution from ascending motions in Central and North America (Hoskins et al., 2020).

The Walker circulation is the zonal overturning circulation found in the equatorial Pacific Ocean characterized by ascending motion over the West Pacific and descending motions over the East Pacific(Bjerknes, 1969; Gill, 1980). The dynamic and thermodynamic effects of the location and strength of convection in the Walker circulation have strong impacts across all the tropics and also the extratropics (Cai et al., 2019).

The Inter-tropical Convergence Zone (ITCZ) is a band of precipitation that migrates meridionally with the seasons (Schneider et al., 2014). The ITCZ is arguably one of the most relevant features of tropical climate because of the strong influence on the low- and upper-level

circulation associated with ITCZ and the fact that the vast majority of rainfall in the tropics is due to convection in the ITCZ. The ITCZ is characterized by a strong convergent flow in the low levels and a strong divergent flow at upper levels. The meridional migration of the ITCZ, as well as the mean latitude of the ITCZ, results from the energy and momentum balances so that the ITCZ is predominantly north of the equator because of the inter-hemispheric temperature contrast (Donohoe et al., 2013; Bischoff and Schneider, 2016).

The traditional view of a monsoon is that of a large-scale land-sea breeze associated with the different heatings of the land and the ocean that force a seasonal reversal of the winds. However, this framework has recently been challenged by the notion that monsoons are simply a manifestation of the ITCZ migration (Gadgil, 2018). This has created the concept of the global monsoon as a phenomenon that exists all across the continental tropics (Zhou et al., 2016; Gadgil, 2018).

The global monsoon refers to the those regions of the planet where more than 70% of the total annual rainfall falls during the summer season (Zhou et al., 2016; Wang et al., 2017). Figure 2.1 shows the global monsoon as depicted by the TRMM dataset. By this definition, the majority of the regions over land between 5 and 10 degrees away from the equator are part of the global monsoon.

A regional monsoon, such as the Indian Monsoon, is then a subset of the global monsoon with unique regional characteristics that shape this monsoon different to other regional monsoons in terms of the seasonality, the strength and the dynamics.

The American Monsoon System is then the regional monsoon that is located in the subtropics of North and South America.

## 2.2 The American Monsoon System

The AMS is typically subdivided into the North and South American monsoon systems (Vera et al., 2006). The North American Monsoon is the main source of rainfall in south-western North America, extending north from central-west Mexico into the southwestern United States (Adams and Comrie, 1997; Stensrud et al., 1997; Vera et al., 2006). The seasonal cycle of rainfall in the North American Monsoon is characterised by a wet July-August-September season and significantly drier conditions during the rest of the year (Adams and Comrie, 1997). A key aspect of this monsoon is the moisture advected by the low-level flow from the

Gulf of California and the East Pacific Ocean and to a lesser extent the moisture mixed in the mid-troposphere from the Caribbean Sea and Gulf of Mexico (e.g Stensrud et al., 1997; Pascale and Bordoni, 2016; Ordoñez et al., 2019).

The South American Monsoon is a primary source of precipitation for South America, especially in the Amazon region (Gan et al., 2004; Vera et al., 2006; Jones and Carvalho, 2013). During austral summer (DJF), monsoon rainfall accounts for over 60% of the total annual precipitation in the Amazon (Gan et al., 2004; Marengo et al., 2012), whereas austral winter rainfall accounts for less than 5% of the total annual rainfall (Vera et al., 2006). In the central Amazon, convective precipitation is observed from early October but the main rainy season extends from December to April (Machado et al., 2004; Adams et al., 2013), whereas convection in southeastern Brazil and Paraguay starts in November and peaks in January and February (Marengo et al., 2001; Nieto-Ferreira and Rickenbach, 2011).

## 2.3 Review of the Midsummer drought

The seasonality of precipitation in northwestern Central America and southern Mexico fits the new definition of a monsoon characterized by the sharp contrast between the wet and dry seasons. However, this region shows a unique climatological precipitation feature. After monsoon onset, rainfall decreases considerably around the midsummer; this decrease is followed by a secondary increase in precipitation in the late summer (Mosino and García, 1966).

These variations of precipitation are well known by local farmers who refer to the drier midsummer period as ‘El Veranillo’ in Central America and ‘canícula’ in southern Mexico because the drier period coincides with the Canis Major constellation appearing in the sky (Dilley, 1996). This feature of the seasonal cycle is most commonly referred to in the literature as Midsummer drought (MSD). Farmers in Central America who are subject to climatic stress due to droughts, have already perceived and adapted to changes in the characteristics of the rainy season, such as the timing and strength of the midsummer drought (Hellin et al., 2017; de Sousa et al., 2018; Harvey et al., 2018), but it is unclear whether these perceived changes are a real trend in the observations (Anderson et al., 2019).

Observations have characterised this feature in the climatological precipitation of several regions of Mexico, El Salvador, Belize, Guatemala, Cuba (e.g. Mosino and García, 1966; Magaña et al., 1999; Durán-Quesada et al., 2017; Perdigón-Morales et al., 2018). However, notable

differences in the seasonal cycle of precipitation have been found between the mainland and the Caribbean. The so-called first peak of precipitation occurs in May in Cuba and in June in northern Central America whereas the second peaks are observed in October and September for the mainland and the Caribbean, respectively.

In spite of extensive research to understand the physical mechanisms associated with the MSD (e.g. Magaña et al., 1999; Giannini et al., 2000; Gamble et al., 2008; Ryu and Hayhoe, 2014; Herrera et al., 2015; Maldonado et al., 2017; Straffon et al., 2019), debate remains over which is the leading-order mechanism that causes rainfall to decrease at midsummer and increase again at the end of the summer. Dynamical or the thermodynamical mechanisms have been put forth and different roles have been proposed for the Atlantic and the East Pacific Oceans (e.g. Magaña and Caetano, 2005; Gamble et al., 2008; Herrera et al., 2015).

Fundamental questions remain unclear such as whether the MSD is caused by two precipitation enhancing mechanisms (Karnauskas et al., 2013) or a mechanism that inhibits rainfall at midsummer (Durán-Quesada et al., 2017). Furthermore, the association between the MSD in Central America and in the Caribbean is still disputed (Gamble et al., 2008), as most studies suggest that the two regimes are unrelated and therefore two different explanations are required to account for the two MSDs in these regions.

Any complete theory or conceptual model must account for the following characteristics of the seasonal cycle. First, the theory must explain the timing and strength of the first peak of rainfall. Second, the timing and strength of the MSD, i.e., what causes rainfall to decrease at midsummer. Finally, the theory must explain the timing and mechanism driving the second increase in precipitation after the midsummer.

Magaña et al. (1999) and Magaña and Caetano (2005) proposed a mechanism driven by radiative-convective feedbacks between the East Pacific SSTs and deep tropical convective clouds. The height and strength of convection, the incoming shortwave and the SSTs are strongly coupled in their framework. The peak in SSTs during May (Figure 4.4a) triggers evaporation and deep convection in the East Pacific ITCZ and Central America (Figure 4.1). The high convective clouds produce a radiative cooling effect at the surface due to decreased incoming shortwave radiation (Figure 4.4d). This cooling decreases SSTs and deep convective activity and thus accounts for the modest decrease in rainfall during the midsummer. The second peak in September is driven by the feedback effect caused by decreased frequency of tall convective clouds during July and August, as convective activity decreased during the MSD,

which reduces the cooling effect of the clouds and increases incoming shortwave, SSTs and surface fluxes, eventually increasing precipitation (Magaña et al., 1999).

Other studies suggest the seasonal evolution of North Atlantic Subtropical High (NASH) and the associated geostrophic flow are the primary cause of the bi-modal regime, particularly for the MSD in the Caribbean (e.g. Mapes et al., 2005; Gamble et al., 2008; Curtis and Gamble, 2008). The NASH is a subtropical anticyclone in the Atlantic Ocean that shifts southwest early in boreal summer. The expansion and intensification of the NASH in boreal summer, according to this theory, strengthens the low-level trade winds, controlling the seasonal cycle of a low-level jet known as the Caribbean Low-Level Jet (CLLJ).

The CLLJ is key for regional variability and climate of the Caribbean, northern central America and southern United States, because the strength, height and direction determines the regional moisture transport (Giannini et al., 2000; Martinez et al., 2019; García-Martínez and Bollasina, 2020). The expansion of the western flank of the NASH is argued to strengthen the CLLJ which cools the SSTs, through the effect of wind stress and mixed-layer mixing (Gamble et al., 2008; Martinez et al., 2019). The cooling of SSTs diminishes evaporation and therefore low-level moisture which leads to less precipitation, at least locally. Because the expansion of the NASH is closely aligned with the timing of the MSD in the Caribbean, the NASH effect on the CLLJ is argued to generate the bimodal regime of precipitation (Gamble et al., 2008; Martinez et al., 2019).

The moisture flow from the Caribbean Sea to the continent and the easternmost Pacific Ocean is another active topic of research (Hidalgo et al., 2015; Corrales-Suastegui et al., 2020). The effect of the SST gradient between the easternmost Pacific Ocean and the Caribbean Sea on the moisture flow across the continent is a relevant for variability on interannual time-scales (Martinez et al., 2020). The Pacific Ocean is projected to warm more than the Caribbean Sea in future decades, which will change the SST gradient, strengthen the CLLJ and shifting the regional precipitation patterns (Corrales-Suastegui et al., 2020).

Herrera et al. (2015) shows that during the drier months in Central America, stronger convective activity is found west of the Central American coast. This evidence suggests that the coupling of EP SSTs to the gap flow that originated from the CLLJ in the Caribbean Sea controls the location of ascending and descending motions, thereby explaining some features of the Central American MSD. Herrera et al. (2015) argued that the exit region of the CLLJ

is located to the east of the region of strongest MSD signal, which suggests that the moisture divergence effect over the central American MSD is minimal.

A different mechanism, proposed by Karnauskas et al. (2013), argues that the biannual crossing of the solar declination angle can control precipitation to the extent of explaining the bimodal characteristics of the seasonal cycle. In this mechanism, the MSD is driven by two precipitation enhancing periods that are separated by a relatively normal, and drier, period. This theory differs from those previously discussed which explained the MSD through mechanisms that inhibit convective activity in the midsummer whereas Karnauskas et al. (2013) argues that the solar declination angle that crosses twice through Central America, once during June and a second time during September, increases convective activity during each crossing.


The variations of incoming shortwave radiation associated with the declination angle modulate the SSTs, surface fluxes and therefore convective activity. In other words, the first crossing of the solar declination angle increases the incoming shortwave radiation which increases the SSTs, evaporation and precipitation, i.e., the first peak. The second crossing, similarly, explains the second peak as the second increase in incoming shortwave promotes more deep convection than during the MSD.

Other mechanisms have been proposed arguing that the MSD is a result of the double crossing of the Intertropical Convergence Zone (ITCZ), the result of vertical wind shear affecting convective instability or the Saharan dust controlling the microphysics of clouds (Angeles et al., 2010). For instance, Perdigón-Morales et al. (2019) also finds a link between the frequency and spatial distribution of the first peak rainfall rates and the Madden-Julian Oscillation.

## 2.4 El Niño Southern Oscillation: impacts to the American monsoon system

ENSO is the leading mode of interannual variability in the tropics which produces changes in the atmospheric circulation that impact most regions on the planet. ENSO has motivated extensive research using GCMs to understand the mechanisms related the origin of ENSO (Christensen et al., 2017), the feedbacks and processes that phase-lock the phenomena (Neelin et al., 1998), as well as how will ENSO characteristics change in the future (Cai et al., 2015a; Santoso et al., 2017).

ENSO is the oscillation from a mean state of the Walker circulation and the SST gradient. El Niño was initially the term to refer to the SST changes and the Southern Oscillation



**Figure 2.2:** Four-chamber illustration of the human heart. Clockwise from upper-left: right atrium, left atrium, left ventricle, right ventricle.

(SO) is associated with changes in the zonal gradient of MSLP. Combined, El Niño and the SO compose the ENSO phenomenon.

The most prominent characteristic of ENSO is the effect that zonal gradients of MSLP and SST have on the location and strength of deep convection in the equatorial Pacific (Trenberth, 1997; Neelin et al., 1998). As ENSO shifts from the positive to the negative phase, deep convection shifts along the equatorial Pacific, generating precipitation anomalies on inter-annual timescales (Neelin et al., 1998; Wang and Picaut, 2004). In other words, ENSO poses a strong control on the location and strength of the Walker circulation which then produces effects in remote regions of the world where the Walker circulation is important, *e.g.*, in West Africa (Ropelewski and Halpert, 1986, 1987) or South America (Sulca et al., 2018).

The period of ENSO of 2-7 years (Neelin et al., 1998; Wang and Picaut, 2004) was poorly represented in CMIP3 and CMIP5 models (Guilyardi et al., 2009), particularly by models that had much stronger power spectrums than the observed.

## 2.5 Stratosphere-Troposphere Coupling in the Tropics

The QBO is the dominant mode of tropical stratospheric variability which modulates the characteristics of the tropopause in the tropics (Baldwin et al., 2001; Tegtmeier et al., 2020) such as height and cold point temperature. Observations have shown an influence of the QBO on deep convective features such as monsoons (Giorgetta et al., 1999), tropical cyclones (Chan, 1995) and the Walker circulation (Collimore et al., 2003). More recently a link between the QBO and the Madden-Julian Oscillation (MJO) was discovered (Son et al., 2017) and motivated extensive research (see e.g. Lee and Klingaman, 2018; Wang et al., 2019; Martin et al., 2020).

The MJO in observations shows a stronger amplitude and more predictability during QBO E, but further inspection in cloud-permitting and forecast models have not provided conclusive answers to this puzzle (Martin et al., 2019, 2020). Questions still arise as to whether this tropical link is real or due to chance, for instance Wang et al. (2019) argued that the increased predictability of the MJO under the QBO E phase was included in the initial conditions, and thus not a result of a mechanistic effect of the QBO on the MJO. More generally, whether the QBO has a considerable effect on deep convection in general is debated as several plausible mechanisms exist in the literature (see e.g. Nie and Sobel, 2015) such as the effect of wind shear, the tropopause height, the cold-point temperature, static stability and/or feedbacks with very high cirrus and cumulonimbus clouds.

The use of climate models to understand these tropical teleconnections of the QBO has proven difficult due to biases in both the MJO and the QBO representations. State-of-the-art CMIP6 models struggle to reproduce several of the characteristics of the QBO (Richter et al., 2020). For instance, weaker temperature QBO signals in the lowermost tropical stratosphere in the models, e.g., compare the QBOE-W difference plot in Figure ???. The weaker temperature signal may be key for possible biases in the tropical QBO links discussed above, such as the the QBO-MJO link which is missing from CMIP6 models (Kim et al., 2020) and from seasonal prediction forecast models (Wang et al., 2019; Martin et al., 2020).

# 3

## Data and methods

### Contents

---

3.1 Observations and reanalysis data . . . . .	12
3.2 Model data . . . . .	14

---

### 3.1 Observations and reanalysis data

Table 3.1 summarises relevant information of the observations and reanalysis datasets used in this study. In short, surface and satellite observations were used where available, whereas other metrics were taken from reanalysis data from the European Centre for Medium-Range Weather Forecasts (ECMWF): ERA-5, downloaded from <https://climate.copernicus.eu/climate-reanalysis>. Four different precipitation datasets are used.

The Tropical Rainfall Measurement Mission (TRMM) dataset is a multi-satellite multi-sensor product that in some versions is calibrated with gauge analyses (Huffman et al., 2007). A set of microwave sensors onboard low earth orbit (LEO) satellites, such as the Microwave Imager (TMI) and the Advanced Microwave Scanning Radiometer-Earth Observing System (AMSR-E), provide the main source of information about hydrometeors for TRMM. However, even using the products of several satellites there is a sparse sampling of time-space precipitation in passive microwave techniques. Therefore, this data is complimented by infrared measurements onboard geosynchronous earth orbit satellites. Other sources of information include a radar

**Table 3.1:** Summary of the datasets used in this study. For each dataset, the acronym used hereafter, the period of coverage, the field used and the horizontal resolution are shown. Some datasets extend further back in time, but only the satellite-era period is used in most of the datasets. The variables used are: precipitation, surface-air temperature ( $2mT$ ), sea-level pressure (SLP), SSTs, the x and y components of the wind ( $u, v$ ), the lagrangian tendency of air pressure ( $\omega$ ), outgoing longwave radiation (OLR), geopotential height (GPH) and specific humidity ( $q$ ).

Dataset/ Version	Acronym	Variable	Period	Data type	Resolution	Reference
Global Precipitation Climatology Project v2.3	GPCP	Precipitation	(1979-2018)	Surface and satellite	2.5°x2.5°	(Adler et al., 2003)
Global Precipitation Climatology Centre	GPCC	Precipitation	(1940-2013)	Surface station	0.5°x0.5°	(Becker et al., 2011)
Climate Prediction Center	CMAP	Precipitation	(1979-2016)	Satellite calibrated with surface rain-gauge	2.5x2.5°	(Xie and Arkin, 1997)
Merged Analysis of Precipitation						
Climatic Research Unit TS v4.	CRU4	Surface temperature	(1979-2017)	Surface station	0.5°x0.5°	(Harris et al., 2014)
Climate Hazards Infrared Precipitation with Stations	CHIRPS	Precipitation	(1981-2018)	Surface rain-gauge and satellite	0.05°x0.05°	(Funk et al., 2015)
Tropical Rainfall Measurement Mission 3B42 V7	TRMM	Precipitation	(1999-2018)	Satellite calibrated with surface station	0.25°x0.25°	(Huffman et al., 2010)
Hadley Centre SST3	HadSST	SST	(1940-2018)	Buoy and satellite	2.5°x2.5°	(Kennedy et al., 2011)
European Centre for Medium-Range Forecasting ERA-5	ERA-5	$2mT$ , SLP, $u$ , $v$ , $\omega$ , OLR, $q$ , SST, GPH, precipitation	(1979-2018)	Reanalysis	0.75x0.75°	(C3S, 2017; Hersbach et al., 2020)

onboard TRMM and rain gauge analysis. Details of the research product can be found in Huffman et al. (2007) and Huffman et al. (2010).

The Climate Prediction Center Merged Analysis of Precipitation (CMAP) dataset is a global merged product of satellite and ground based observations but also constrained by a numerical model (Xie et al., 2007). This dataset was first produced at monthly-mean resolution (Xie and Arkin, 1997) but is now available as a collection of products at several temporal scales. The pentad-scale version of CMAP is used in this study.

The Climate Hazards Infrared Precipitation with Stations (CHIRPS) is relatively more recent merged product. This dataset uses high-resolution rain-gauge station data that is complimented by satellite cloud cold duration estimates on regions where station data is sparse. The products are calibrated with TRMM data (Funk et al., 2015), so they are cannot be considered an independent source of information from TRMM.

The TRMM dataset has a high horizontal and temporal resolution and was used in several CMIP assessments (Geil et al., 2013; Jones and Carvalho, 2013) as a reliable source of precipitation (Carvalho et al., 2012). Therefore, we use TRMM as our best estimate for the spatial and temporal characteristics of the AMS rainfall. However, the period covered by TRMM (1998-2018) is too short to analyse statistically robust teleconnections or variability, so we use GPCP, GPCC and CHIRPS for their longer period. Although a thorough validation and comparison of these datasets across the AMS domain is missing, several studies have analysed one or more of these datasets in regions of the AMS (e.g. Franchito et al., 2009; Dinku et al., 2010; Trejo et al., 2016).

## 3.2 Model data

The MOHC has submitted the output of two models for CMIP6: HadGEM3 GC3.1 (hereafter GC3) is the latest version of the Global Coupled (GC) Met Office Unified Model (UM) and UKESM1, the new U.K. Earth System Model. The most substantial change from the version used in CMIP5 (HadGEM2-AO) is the inclusion of the new GC configuration 3.1 (Walters et al., 2019) with the updated components: Global Atmosphere 7.0 (GA7.0), Global Land 7.0 (GL7.0), Global Ocean 6.0 (GO6.0), and Global Sea Ice 8.0 (GSI8.0). The GC3.1 configuration runs with 85 atmospheric levels, 4 soil levels and 75 ocean levels; for details see Williams et al. (2018) and Kuhlbrodt et al. (2018). The GC3 model was run for CMIP6 deck experiments

**Table 3.2:** Summary of the CMIP6 simulations in this study. For each simulation the acronym used hereafter, the experiment and the horizontal resolution are shown. The first 100 years of the piControl simulations are used and for historical experiments the period 1979-2014 is used.

Model	Experiment	Atmospheric resolution	Ocean resolution	Acronym	Reference
Hadley Centre Global Environment Model version 3 (HadGEM3)	Pre-industrial control	N96 1.875°x1.25°	1°	GC3 N96-pi	1 (Menary et al., 2018; Ridley et al., 2018)
HadGEM3	Pre-industrial control	N216 0.83°x0.56°	0.25°	GC3 N216-pi	1 (Menary et al., 2018; Ridley et al., 2019c)
HadGEM3	Historical	N96 1.875°x1.25°	1°	GC3-hist	4(r1-r4) (Andrews et al., 2020; Ridley et al., 2019b)
HadGEM3	Historical	N216 0.83°x0.56°	0.25°	N216-hist	1 (Ridley et al., 2019c)
HadGEM3	Atmospheric Model Inter-comparison (AMIP)	N96 1.875°x1.25°	1°	GC3-AMIP	5 (r1-r5) (Ridley et al., 2019a)
United Kingdom Earth System Model version 1 (UKESM1)	Pre-industrial control	1.875°x1.25°	1°	UKESM-pi	1 (Tang et al., 2019b)
UKESM1	Historical	1.875°x1.25°	1°	UKESM-hist	5 (r1-r5) (Tang et al., 2019a)

with two horizontal resolutions: a low resolution configuration, labelled as N96, with an atmospheric resolution of  $1.875^\circ \times 1.25^\circ$  and a  $1^\circ$  resolution in the ocean model and a medium resolution configuration, labelled N216, with atmospheric resolutions of  $0.83^\circ \times 0.56^\circ$  and a  $0.25^\circ$  oceanic resolution (Menary et al., 2018).

The UKESM1 was recently developed aiming to improve the UM climate model adding processes of the Earth System (Sellar et al., 2019). These additional components include ocean biogeochemistry with coupled chemical cycles, tropospheric-stratospheric interactive chemistry which aim to better characterise aerosol-cloud and aerosol-radiation interactions (Mulcahy et al., 2018; Sellar et al., 2019). The physical atmosphere-land-ocean-sea-ice core of the HadGEM3 GC3.1 underpins the UKESM1, so that the UKESM1 and the HadGEM3 have the same dynamical core but the UKESM1 has the additional components mentioned above.

This study uses three CMIP6 deck experiments. First, the pre-industrial control (piControl) simulations, which are run with constant forcing using the best estimate for pre-industrial (1850) forcing of aerosols and greenhouse gas levels. The historical experiments are 164-yr integrations for 1850-2014 that include historical forcings of aerosol, greenhouse gas, volcanic and solar signals since 1850 (Eyring et al., 2016; Andrews et al., 2019). For further details, Andrews et al. (2020) extensively describes the historical simulations of HadGEM3-GC3.1.

In contrast to the pre-industrial control experiments, the historical experiments use time-varying aerosol and greenhouse gas emissions and land-use change (Eyring et al., 2016). In Latin-America, land-use change for agricultural purposes has dramatically decreased tree cover in Central America and south-eastern Brazil since the 1950s (Lawrence et al., 2012), thereby affecting the surface energy balance. The regional emissions of carbonaceous aerosols, nitrogen oxides and volatile organic compound in Latin America are also considered in the historical experiments. These emissions are noteworthy, e.g., due to the impact of black carbon emissions by increased biomass burning in the Amazon and northern Central America (Chuvieco et al., 2008).

The historical experiments of HadGEM3 and UKESM1 are composed of 4 and 9 ensemble members, respectively, but the results will be presented as the ensemble mean for the 1979-2014 period. These experiments will be referred to as GC3-hist and UKESM1-hist hereafter. Finally, we use the five ensemble members of the AMIP experiment from GC3 N96 covering 1979-2014. Table 3.2 summarises the main features of the experiments used in this study.

*Alles Gescheite ist schon gedacht worden.  
Man muss nur versuchen, es noch einmal zu denken.  
All intelligent thoughts have already been thought;  
what is necessary is only to try to think them again.*

— Johann Wolfgang von Goethe ?

# 4

## On the dynamical and thermodynamical mechanisms of the MSD in the Met Office CMIP6 models

### Contents


---

<b>4.1 Climatological features</b> . . . . .	<b>17</b>
4.1.1 On the mechanisms of the MSD in the UK Met Office models . . . . .	20
4.1.2 Summary and discussion . . . . .	27

---


### 4.1 Climatological features

Figure 4.1 shows the pentad-mean seasonal cycle of precipitation in Central America and the Caribbean. The seasonal cycle in both regions follows that of a monsoon, i.e., a dry winter and a wet summer season. In the first region (Figures 4.1a, b), two precipitation maxima, in June and September, are separated by a decrease in precipitation during July and August, *i.e.* the MSD. In Central America, the difference between the first peak (June 15) to the driest pentad of the MSD (Aug 01) is of about  $2 \text{ mm day}^{-1}$ , according to TRMM. The two peak structure in the Caribbean (Figures 4.1c, d) is characterised by two peaks in May and October with a four-month drier period in between the two peaks (e.g. Giannini et al., 2000; Gamble et al., 2008; Angeles et al., 2010). In Cuba, the difference between the first peak (June 01) to the driest pentad of the MSD (Aug 01) is of about  $3 \text{ mm day}^{-1}$  in the TRMM dataset.



**Figure 4.1:** Pentad-mean precipitation in (a) southern Mexico and northern Central America and (b) Cuba. Shading shows uncertainty obtained by bootstrapping the interannual variability of the TRMM dataset.

Precipitation in these regions depends on several factors such as the seasonal migration of the East Pacific (EP) and Atlantic ITCZs. The SSTs in the Gulf of Mexico, the Caribbean Sea, the western tropical Atlantic and the Eastern Pacific are also very relevant for the seasonal cycle and interannual variations (Magaña et al., 1999; Amador, 2008; Straffon et al., 2019). Figures 4.4a, b show the seasonal cycle of SSTs in the EP and the Caribbean Sea. While the EP shows a maximum in SSTs in late May, during the early stages of the monsoon in Central




**Figure 4.2:** (a, d, g) Climatological JJAS rainfall and the difference between (b, e, h) the midsummer drought and the first peak periods and (c, f, i) between the second peak and the midsummer drought periods for (a-c) TRMM, (d-f) ERA5 and (g-i) CHIRPS.

America, the Caribbean SSTs peak in early fall, about five months later.

The Caribbean Low-level Jet (CLLJ) is a strong low-level easterly jet in the Caribbean Sea that peaks at the end of June (Figure 4.4e) at the 925 hPa level (Amador, 2008; Herrera et al., 2015; Maldonado et al., 2016). The CLLJ determines the moisture transport from the Caribbean Sea into the eastern Pacific across the Central American landmass as well as the northward moisture transport into the Gulf of Mexico and Florida (Muñoz et al., 2008; Hidalgo et al., 2015; Maldonado et al., 2016).

However, as shown in Figure 4.4a and as discussed for the radiative-convective feedback of Magaña et al. (1999), SSTs do not increase in the East Pacific in the late summer and the second increase in incoming shortwave is only modest in the reanalysis (Figure 4.4d).

However, SSTs in the easternmost Pacific do not increase after, during or at the end of the MSD (Figure 4.4a). In fact, the SSTs decrease with the second increase in deep convection and precipitation. The other hypothesis of this theory, referring to the incoming shortwave is also not consistent with observations, as the incoming shortwave only modestly increases during the midsummer (Figure 4.4d). There is perhaps a role for this modest increase in incoming shortwave, but the link to SSTs suggested by this theory does not agree with the reanalysis.




**Figure 4.3:** (a, d, g, j) JJAS model bias compared to TRMM and the difference between (b, e, h, k) the midsummer drought and the first peak periods and (c, f, i, l) between the second peak and the midsummer drought periods for four different simulations.

#### 4.1.1 On the mechanisms of the MSD in the UK Met Office models

Biases in the strength and position of the EP ITCZ in Global Coupled Models (GCMs) (Bellucci et al., 2010; Li and Xie, 2014; Schneider et al., 2014) are a major reason for biases in the model representation of rainfall in Central America (Rauscher et al., 2008).

Ryu and Hayhoe (2014) analyzed the performance of CMIP3 and CMIP5 models and found that the majority of CMIP5 models were unable to represent the total annual rainfall and the seasonal cycle of the MSD. Ryu and Hayhoe (2014) also finds that models that simulate a bimodal distribution of rainfall, HadGEM2-A for example, also show an accurate seasonal cycle of the NASH and the CLLJ. However, an exhaustive analysis as to whether these features are actually driving mechanisms for the MSD in GCMs as in observations is missing from the literature.

The CMIP6 Met Office models, HadGEM3 and UKESM1, are amongst the first models to simulate a bimodal regime in both Central America and Cuba (Figure ??a and 4.1). In Central America and southern Mexico, the models simulate a wetter-than-observed first peak



**Figure 4.4:** Pentad-mean seasonal cycle of indices associated with the MSD in Central America and the Caribbean.

of precipitation and a drier MSD period. The so-called second peak of precipitation found in late August is simulated in close agreement with TRMM, except in the AMIP experiment which has a far too strong second peak mean precipitation rate.


Figure 4.3 shows the distribution of rainfall in the different stages of boreal summer in different CMIP6 experiments and ERA-5. The main feature, the East Pacific ITCZ shows the maximum rainfall rates ( $>15 \text{ mm day}^{-1}$  in the models) and strong mid-level ascent ( $-0.1 \text{ Pa s}^{-1}$ ). Prior to the MSD, rainfall extends from the easternmost Pacific ITCZ into the North American continent. Therefore, the positive bias during the first peak over land is associated with the biased wetter EP ITCZ. However, during the MSD, rainfall decreases over land remaining only above  $10 \text{ mm day}^{-1}$  south west of the coastline in the models.

The wetter EP ITCZ is a common feature of GCMs, including the Met Office models, which results from multiple biases in the radiative and convective schemes (Oueslati and Bellon, 2013; Li and Xie, 2014). In UKESM1 and HadGEM3 several biases exist in the radiative balance in the easternmost Pacific Ocean. A positive bias in incoming shortwave in Central America of about 15% and a cold SST bias in both East Pacific and Caribbean Sea SSTs are observed in Figure 4.4. Increased incoming shortwave but cooler SSTs require increased surface fluxes to maintain energy balance. These higher latent heat fluxes (LHF<sub>s</sub>) in the models in both basins (Figs. 4.4g, h) are almost 40% larger than in ERA5 during the first peak of rainfall. The models also exhibit a larger seasonal cycle of the fluxes than the reanalysis. GC3 AMIP is the only simulation to also show a significantly positive bias in LHF<sub>s</sub> during the second peak of rainfall in the EP but also at the end of MSD in the Caribbean Sea.

In all the model experiments, the ITCZ prior to the MSD period is stronger than in ERA5 by more than 5 mm day<sup>-1</sup>, whereas after the MSD rainfall in the coupled models on the western coast of Central America agrees well the ERA5. This analysis suggests that the biases shown in Figure ?? are mostly coming from the period prior to the MSD. The models reasonably simulate the decrease in rainfall during the MSD (Figure 4.3) followed by the second increase or peak. Note that GC3 AMIP, forced by very similar SSTs as ERA-5, simulated a much larger mean precipitation in the ITCZ during MSDE in contrast to the coupled models. This large positive bias in simulated rainfall in the East Pacific in GC3 AMIP corresponds to the larger than observed second peak observed in Figure 4.1a.

Composites prior to the onset of the MSD, during the MSD and after the MSDE were computed for several diagnostic variables. The periods were separated using the WT method to determine the dates of the MSDO and MSDE in ERA5 and the climate model output. Figure 4.5 shows the composite differences between the period of the MSD and of the two peaks in out-going longwave radiation (OLR) and vertical velocity ( $\omega - 500$ ) at 500 hPa. The positive OLR and  $\omega$  anomalies in the MSD-MSDO panels in southern Mexico and northern Central America are indicative of decreased height of convection and decreased ascent, in agreement with the MSD being the drier period. These positive anomalies in the continent are accompanied by negative OLR and  $\omega - 500$  anomalies west of the continent, around 125°W.


The MSDE-MSD panels show the difference between the second peak of rainfall and the drier MSD period. Negative OLR and  $\omega$  anomalies indicate stronger and higher convection over a wide region including the easternmost Pacific Ocean, southern Mexico, northern Central America Cuba



**Figure 4.5:** Out-going longwave radiation (OLR) [ $\text{W m}^{-2}$ ] (shaded) and  $\omega$  500-hPa [ $10^{-2} \text{ Pa s}^{-1}$ ] (line contours) differences between the MSD and MSDO and the MSDE and MSD.

and the Caribbean Sea. Note also the region of the North American Monsoon, on the northwest corner of Mexico and the southernmost US, as the MSD-MSDO difference suggests increased convective activity in the North American Monsoon region and MSDE-MSD the opposite.


Similarly, Figure 4.6 shows the low-level wind field during the three stages of the MSD. In ERA-5, prior to the MSD the wind flow in the Caribbean shows strong easterlies that flow into the Gulf of Mexico and southeastern US but very weak winds in the EP (see Figure 4.4f). During the MSD, the winds in the EP become modestly strong easterlies associated the easterly flow from the Caribbean Sea that crosses over Costa Rica and Nicaragua from the Caribbean Sea to the East Pacific. Note that the easterlies converge towards the region at 125°W where OLR and  $\omega$  anomalies suggest increased ascent.



**Figure 4.6:** As in Figure 4.5 but showing wind vectors at the 850 hPa level.

By the end of the MSD the easterlies in ERA5 weaken substantially on the western coast of Central America and in the Caribbean Sea. The simulations seem to generally reproduce the characteristics of the wind field with some differences worth mentioning. For instance, prior to the onset of the MSD, all the simulations show a modest westerly wind flow in the east Pacific at 10°N, which can also be seen in Figure 4.4f, which is not observed in ERA5. After the MSD ends, most simulations show a very weak westerly flow in the East Pacific, close to ERA5; however, GC3 AMIP shows a modest westerly wind converging towards the west coast of Nicaragua. This low-level convergence may be forcing the increased convective activity and precipitation during this time in GC3 AMIP.


The SSTs and incoming shortwave radiation are key elements for explain the seasonal cycle of the MSD, according to previous theories summarised in section 2.3. Figure 4.7 shows the corresponding SST and incoming shortwave anomalies during the different stages of the seasonal cycle. From the first peak to the MSD, a positive SST difference of +1.5 K in the Gulf of California and the western coast of the Baja California Peninsula is observed in reanalysis



**Figure 4.7:** As in Figure 4.5 but the anomalies are shown for SSTs [K] (contours) and incoming shortwave radiation [ $\text{W m}^{-2}$ ] at the surface (line-contours). Incoming shortwave is defined such as negative differences imply less incoming shortwave and positive anomalies represent more incoming shortwave at the surface.

and the models. The differences appear as a sharp SST meridional gradient pattern around  $115^{\circ}\text{W}$ . During this stage, the incoming shortwave increases in Central America, which agrees with Figure 4.4d. Note the negative incoming shortwave differences west of Central America at  $125^{\circ}\text{W}$ , the region of negative OLR and  $\omega$ -500 hPa anomalies where low-level winds converge, all of which supports the notion of increased convective activity that reduces incoming shortwave west of the continent. This feature was noted by Herrera et al. (2015).

After the MSD, the western coast of the Baja California Peninsula continues to warm and the East Pacific continues to cool, in contrast to previous suggestions (Magaña et al., 1999; Magaña and Caetano, 2005; Herrera et al., 2015). Meanwhile, the Caribbean Sea warms by 1



**Figure 4.8:** As in Figure 4.5 but showing in shading, moisture flux divergence  $\nabla \cdot \vec{u}q$  at the 850 hPa level with units of  $10^{-7} \text{ s}^{-1} \text{ kg / kg}$  and zonal wind anomalies (line contours) in  $\text{m s}^{-1}$ .

K and the northern Gulf of Mexico slightly cools down. The incoming shortwave differences show a regional-scale decrease in incoming shortwave, as the summer draws to an end. These SST differences indicate that the meridional SST gradient in both the EP and Caribbean Sea and Gulf of Mexico is greatly modified during the stages of the MSD.

The main dynamical argument put forth to explain the MSD is centred around variations in the moisture flux convergence (MFC), argued to be driven by the Caribbean-Low Level Jet (see e.g. Gamble et al., 2008; Herrera et al., 2015; Martinez et al., 2019). The MFC and zonal wind variations in each stage of the MSD is shown in Figure 4.8 for ERA-5 and two simulations. The low-level MFC increases from monsoon onset (MO) to the first peak period (MSDO) in the EP. This anomaly in MFC corresponds to a region of positive zonal wind anomalies indicative of weaker easterly flow. This zonal wind anomaly from MSD to MSDO is much stronger in the models. The MSD-MSDO difference shows a strong positive MFC anomaly across southern Mexico and most of Central America.

In turn, the MFC anomalies associated with the end of the drier period, observed as the MSDE-MSD anomalies, show negative values, suggesting increased moisture flux, over southern Mexico and northern Central America. Increased moisture flux during the transition from the MSD to the second peak agrees well with the precipitation differences during these periods. The


MSDE-MSD zonal wind anomalies in the EP show positive zonal wind anomalies, suggesting a weakened easterly wind flow (see also Fig. 4.6).

The MSD in Central America and southern Mexico has been strongly linked to the strengthening of the CLLJ (Herrera et al., 2015). The maximum zonal wind observed in the CLLJ is found at the very end of July (Fig. 4.4e), synchronized with the start of the MSD. The zonal wind anomalies in the MSD-MSDO panels in Figure 4.8 show that easterlies in the Caribbean Sea do not strengthen by more than  $2 \text{ m s}^{-1}$  from the first peak to the MSD. Only in the models is there a modest negative anomaly at the westernmost Caribbean Sea. In other words, while the peak of the climatological CLLJ coincides with the climatological timing of the onset of the MSD, these composite analyses constructed by more specifically separating the MSD periods does not show relevant variations in the zonal wind of the Caribbean Sea. The drier MSD period does coincide with stronger easterly flow over the eastern Pacific, which may be associated with the weaker MFC over land.

#### 4.1.2 Summary and discussion

The midsummer drought is a prominent feature of the seasonal cycle of rainfall of southern Mexico, northern Central America and the Caribbean. The average 20% decrease during the midsummer compared to the wetter periods of early and late summer is a rare feature of monsoon regions that has important implications for agriculture and water management (Hellin et al., 2017; de Sousa et al., 2018; Harvey et al., 2018).

Climate predictions of the MSD, particularly those concerning whether this "drought" will become more pronounced in the following years, are not trustworthy because of several reasons. One factor is the current limitation in the understanding of the physical processes that cause the MSD (section 2.3) as debate still exists over which large or regional-scale processes are most important to explain the increases and decreases of precipitation over intraseasonal time-scales. Secondly, methods used to diagnose the timing and strength of the MSD typically deal with monthly-scale metrics, which would obscure subtle trends and processes that have an effect on shorter time-scales. Also relevant is the fact that climate models used to produce the predictions show significant biases in the EP ITCZ and the seasonal cycle of rainfall in the region, in fact, most CMIP3 and CMIP5 models did not show a bimodal signature in the seasonal cycle. Models that do not have a climatological MSD cannot provide a prediction for this regime in future climate.



**Figure 4.9:** Scatter plot of the (a, b) area-averaged precipitation over land (Box in Figure ??) during the different stages of the MSD. (c) scatter of the East Pacific SSTs against the precipitation over land during the first peak period. (d-l) show the scatter differences in several variables between the different stages of onset of the MSD (MSDO), the drier MSD and the end of the MSDE. The differences are shown for area-averaged quantities in the East Pacific (EP), the Caribbean Sea (CSEA) and overland (MSD) as above. The units for  $\Delta U$  are  $[\text{m s}^{-1}]$ ,  $\Delta M\text{FC}$   $[10^{-11} \text{s}^{-1}]$ ,  $\Delta S\text{W}$  and  $\Delta L\text{HF}$   $[\text{W m}^{-2}]$  and  $\Delta p\text{r mm d}^{-1}$ . The Pearson correlation coefficient for the 38 yr of reanalysis or observations ( $r_{obs}$ ) is shown for each panel.

For these reasons, this section analysed the CMIP6 simulations from the Met Office models, UKESM1 and HadGEM3, aiming to understand the causes of the biases in the seasonal cycle.

Furthermore, these models are better compared to CMIP3 and CMIP5 cohorts since UKESM1 and HadGEM3 actually simulate a bimodal precipitation regime in these regions. The purpose of this investigation is to use these climate models to better diagnose the relevant biases for the representation of the MSD but also understand the processes that these models are capturing leading to the MSD, in order to, hopefully, also highlight the dynamics of the MSD in general.

The wavelet transform method was developed to determine the pentads of onset and end of the MSD. For instance, Figures 4.9a,b show the scatter of the mean precipitation during the first peak against second peak and first peak against MSD in all the simulations and ERA5. The magnitude of the first and second peaks appear to be unrelated in these models and in observations, which would suggest that the processes driving each peak are not exactly the same. Similarly, composite analysis of various diagnostics during the different stages of the seasonal cycle was done, for instance, OLR composites showed that the MSD is not a local feature in a small region of southern Mexico but extends throughout a wide range of North America, from central Mexico through Belize, Guatemala, El Salvador, Honduras, Nicaragua, and northern Costa Rica.

This composite approach also allowed to test previously proposed hypotheses by analysing the differences between model experiments and the observed variability in the characteristics of the precipitation at each stage of the MSD. For example, Magaña et al. (1999) proposed a mechanism that explains the MSD through SST-cloud feedbacks. In this hypothesis, shortwave, SSTs and precipitation are strongly coupled in the EP Ocean. The first peak of precipitation in southern Mexico and Central America would then be associated with the EP SSTs prior to the onset of rainfall. Figure 4.9c shows that EP SSTs prior to onset do not explain the inter-model differences in the magnitude of the first peak nor do they show a strong relationship in the observed interannual variability of the first peak mean precipitation. Similarly, Figure 4.9d shows that surface incoming shortwave variations are only weakly related to SSTs variations in the EP, in both models and reanalysis, during the first peak period.

The feedback mechanism also suggests that the second peak is a result of a second increase in surface incoming shortwave that occurs as cloud cover decreases during the drier MSD. This increase in incoming shortwave then increases EP SSTs and thus increasing convective activity. Although the incoming shortwave does show a bimodal behaviour (Figure 4.4d), the SSTs in the East Pacific do not increase during the MSD period, but in fact cool during the end of the MSD. Furthermore, as in Figure 4.9d, variations in incoming shortwave were not strongly

related to SST changes in any of the stages of the MSD (not shown). This suggests that the SSTs are not only dependent on the incoming shortwave in both models and reanalysis.

The low-level winds (Figure 4.6) show notable changes between the onset of the MSD (MSDO), the MSD and the end of the MSD (MSDE). Weak westerlies in the EP are found during the wetter periods but the zonal wind becomes a modest easterly flow during the drier MSD period. The MSDO appears to be synchronized with the strengthening of the Caribbean Low-Level Jet (Fig. 4.4e). During the MSD, the strong zonal flow in the Caribbean crosses Central America into the central-eastern Pacific. This easterly flow during the MSD converges to 125°W in the EP Ocean, a region that also shows increased ascent during the MSD.

Figure 4.9e, f show the relationships between the zonal flow in the EP Ocean and precipitation in southern Mexico and Central America. The changes in the wind flow between the first and the MSD are not related to the drying response over land during the same period. However, the differences between the second peak and the MSD in the wind flow and precipitation show a strong relationship both in observed interannual variability as well as in the model spread. Simulations with a stronger EP zonal wind anomaly show the strongest increment in precipitation over land. The zonal wind change in the EP from the MSD to the second peak period is also modestly related to the MFC over the continent (Fig. 4.9g) with weaker easterly winds in the EP associated with more convergence over land in the models and reanalysis.

The easterly flow in the EP has been associated with the strength of the CLLJ (Herrera et al., 2015). The zonal wind changes in the MSDE-MSD difference in the EP shows a modest linear relationship with the zonal flow in the Caribbean Sea (Fig. 4.9h). During the other periods, the relationship between the CLLJ and the EP zonal component of the wind is even weaker in both models and observations (not shown).

A potentially relevant bias found in the models was stronger-than-observed surface latent heat fluxes (LHF) (Figure 4.4g, h) compared to the reanalysis. Changes in the surface energy balance and the surface temperature in historical versus pre industrial control simulations may also be responsible for the precipitation differences between these experiments. However, the variations in the LHF, both MSD-MSDO and MSDE-MSD either in the Caribbean Sea or over land (Figure 4.9i,j) are not related to precipitation over land.

The main factor associated with the precipitation variations in the seasonal cycle appears to be the low-level moisture flux convergence (MFC) (Figure 4.9k, l). The variations in the MFC over land explain intermodel differences and observed interannual variability in precipitation,

particularly in the positive rainfall increment from the MSD to the second peak. From the first peak to the MSD, moisture flux decreases and increases again from the MSD to the second peak.

# Appendices

*Cor animalium, fundamentum est vitæ, princeps omnium, Microcosmi Sol, a quo omnis vegetatio dependet, vigor omnis & robur emanat.*

*The heart of animals is the foundation of their life, the sovereign of everything within them, the sun of their microcosm, that upon which all growth depends, from which all power proceeds.*

— William Harvey ?

# A

## Review of Cardiac Physiology and Electrophysiology

### Contents

---

<b>A.1 Anatomy . . . . .</b>	<b>33</b>
<b>A.2 Mechanical Cycle . . . . .</b>	<b>36</b>
<b>A.3 Electrical Cycle . . . . .</b>	<b>37</b>
<b>A.4 Cellular Electromechanical Coupling . . . . .</b>	<b>38</b>

---

Appendices are just like chapters. Their sections and subsections get numbered and included in the table of contents; figures and equations and tables added up, etc. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Sed et dui sem. Aliquam dictum et ante ut semper. Donec sollicitudin sed quam at aliquet. Sed maximus diam elementum justo auctor, eget volutpat elit eleifend. Curabitur hendrerit ligula in erat feugiat, at rutrum risus suscipit. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Integer risus nulla, facilisis eget lacinia a, pretium mattis metus. Vestibulum aliquam varius ligula nec consectetur. Maecenas ac ipsum odio. Cras ac elit consequat, eleifend ipsum sodales, euismod nunc. Nam vitae tempor enim, sit amet eleifend nisi. Etiam at erat vel neque consequat.

### A.1 Anatomy

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Donec accumsan cursus neque. Pellentesque eget tempor turpis, quis malesuada dui. Proin egestas, sapien sit amet feugiat

vulputate, nunc nibh mollis nunc, nec auctor turpis purus sed metus. Aenean consequat leo congue volutpat euismod. Vestibulum et vulputate nisl, at ultrices ligula. Cras pulvinar lacinia ipsum at bibendum. In ac augue ut ante mollis molestie in a arcu.

Etiam vitae quam sollicitudin, luctus tortor eu, efficitur nunc. Vestibulum maximus, ante quis consequat sagittis, augue velit luctus odio, in scelerisque arcu magna id diam. Proin et mauris congue magna auctor pretium id sit amet felis. Maecenas sit amet lorem ipsum. Proin a risus diam. Integer tempus eget est condimentum faucibus. Suspendisse sem metus, consequat vel ante eget, porttitor maximus dui. Nunc dapibus tincidunt enim, non aliquam diam vehicula sed. Proin vel felis ut quam porta tempor. Vestibulum elit mi, dictum eget augue non, volutpat imperdiet eros. Praesent ac egestas neque, et vehicula felis.

Pellentesque malesuada volutpat justo, id eleifend leo pharetra at. Pellentesque feugiat rutrum lobortis. Curabitur hendrerit erat porta massa tincidunt rutrum. Donec tincidunt facilisis luctus. Aliquam dapibus sodales consectetur. Suspendisse lacinia, ipsum sit amet elementum fermentum, nulla urna mattis erat, eu porta metus ipsum vel purus. Fusce eget sem nisl. Pellentesque dapibus, urna vitae tristique aliquam, purus leo gravida nunc, id faucibus ipsum magna aliquet ligula. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Proin sem lacus, rutrum eget efficitur sed, aliquam vel augue. Aliquam ut eros vitae sem cursus ultrices ut ornare urna. Nullam tempor porta enim, in pellentesque arcu commodo quis. Interdum et malesuada fames ac ante ipsum primis in faucibus. Curabitur maximus orci purus, ut molestie turpis pellentesque ut.

Donec lacinia tristique ultricies. Proin dignissim risus ut dolor pulvinar mollis. Proin ac turpis vitae nibh finibus ullamcorper viverra quis felis. Mauris pellentesque neque diam, id feugiat diam vestibulum vitae. In suscipit dui eu libero ultrices, et sagittis nunc blandit. Aliquam at aliquet ex. Nullam molestie pulvinar ex vitae interdum. Praesent purus nunc, gravida id est consectetur, convallis elementum nulla. Praesent ex dolor, maximus eu facilisis at, viverra eget nulla. Donec ullamcorper ante nisi. Sed volutpat diam eros. Nullam egestas neque non tortor aliquet, sed pretium velit tincidunt. Aenean condimentum, est ac vestibulum mattis, quam augue congue augue, mattis ultrices nibh libero non ante. Lorem ipsum dolor sit amet, consectetur adipiscing elit.

Aenean volutpat eros tortor, non convallis sapien blandit et. Maecenas faucibus nulla a magna posuere commodo. Nullam laoreet ante a turpis laoreet malesuada. Phasellus in varius sem. Vestibulum sagittis nibh sed tincidunt blandit. Donec aliquam accumsan odio sit amet lacinia. Integer in tellus diam. Vivamus varius massa leo, vitae ullamcorper metus pulvinar

sed. Maecenas nec lorem ornare, elementum est quis, gravida massa. Suspendisse volutpat odio ex, ac ultrices leo ultrices vel. Sed sed convallis ipsum. Pellentesque euismod a nulla sed rhoncus. Sed vehicula urna vitae mi aliquet, non sodales lacus ullamcorper. Duis mattis justo turpis, id tempus est tempus eu. Curabitur vitae hendrerit ligula.

Curabitur non pretium enim, in commodo ligula. Etiam commodo eget ligula a lacinia. Vestibulum laoreet ante tellus, vel congue sapien ornare in. Donec venenatis cursus velit vitae pulvinar. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Suspendisse in metus lectus. Pellentesque gravida dolor eget finibus imperdiet. Duis id molestie tortor. Mauris laoreet faucibus facilisis. Aliquam vitae dictum massa, sit amet dignissim lacus.

Fusce eleifend tellus id ex consequat maximus. Donec ultrices ex ut turpis ornare, non molestie mi placerat. Nulla sit amet auctor nunc, sit amet euismod elit. Phasellus risus tellus, condimentum a metus et, venenatis tristique urna. Cras mattis felis eget ipsum fermentum egestas. Ut augue odio, venenatis id convallis vel, congue quis augue. Maecenas sed maximus est, posuere aliquet tortor. Ut condimentum egestas nisi eu porttitor. Ut mi turpis, posuere id lorem vel, elementum tempor arcu.

Morbi nisl arcu, venenatis non metus ac, ullamcorper scelerisque justo. Nulla et accumsan lorem. Mauris aliquet dui sit amet libero aliquet, in ornare metus porttitor. Integer ultricies urna eu consequat ultrices. Maecenas a justo id purus ultricies posuere sed et quam. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Sed eleifend risus quis aliquet gravida. Nullam ac erat porta est bibendum dictum in a dolor. Nam eget turpis viverra, vulputate lectus eget, mattis ligula. Nam at tellus eget dui lobortis sodales et ut augue. In vestibulum diam eget mi cursus, ut tincidunt nulla pellentesque.

Aliquam erat volutpat. Sed ultrices massa id ex mattis bibendum. Nunc augue magna, ornare at aliquet gravida, vehicula sed lorem. Quisque lobortis ipsum eu posuere eleifend. Duis bibendum cursus viverra. Nam venenatis elit leo, vitae feugiat quam aliquet sed. Cras velit est, tempus ac lorem sed, pharetra lobortis ipsum. Donec suscipit gravida interdum. Nunc non finibus est. Nullam turpis elit, tempus non ante.

## A.2 Mechanical Cycle

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean tellus est, suscipit sed facilisis quis, malesuada at ipsum. Nam tristique urna quis quam iaculis, et mattis orci pretium. Praesent euismod elit vel metus commodo ultrices. Vestibulum et tincidunt ex, in molestie ex. Donec ullamcorper sollicitudin accumsan. Etiam ac leo turpis. Duis a tortor felis. Nullam sollicitudin eu purus ac hendrerit. Nam hendrerit ligula libero, eget finibus orci bibendum a. Aenean ut ipsum magna.

Ut viverra, sapien sed accumsan blandit, nisi sem tempus tellus, at suscipit magna erat ornare nunc. Proin lacinia, nisi ut rutrum malesuada, nibh quam pellentesque nunc, sit amet consectetur purus felis ac tortor. Suspendisse lacinia ipsum eu sapien pellentesque mattis. Mauris ipsum nunc, placerat non diam vel, efficitur laoreet nunc. Sed lobortis, ipsum eget gravida facilisis, sem nulla viverra mi, in placerat eros sem viverra lacus. Aliquam porta aliquet diam vel commodo. Nulla facilisi. Duis erat libero, lobortis vel hendrerit vitae, sagittis id dui. Nulla pretium eros nec quam tincidunt, vel luctus mi aliquam. Integer imperdiet purus in est tristique venenatis. Ut pellentesque, nunc vitae iaculis ultricies, urna turpis dignissim risus, a laoreet felis magna nec erat.

Quisque sollicitudin faucibus ligula, et egestas nibh dictum sit amet. Proin eu mi a lectus congue pretium eu quis arcu. Suspendisse vehicula libero eu ipsum aliquam, vel elementum nibh mattis. Sed sed sapien vitae turpis tristique pulvinar a ut metus. Etiam semper gravida est, mollis gravida est porta ac. Proin eget tincidunt erat. Maecenas ultrices erat eget purus ultricies, ut lacinia arcu dictum. Nam et nisi sit amet ex congue mattis vel eget lorem. Aliquam erat volutpat. Pellentesque porttitor nibh vitae elementum consectetur. Aenean et est lobortis, congue sapien non, ullamcorper sapien. Ut facilisis sem non dapibus vehicula.

Mauris euismod odio dolor, sit amet gravida mauris placerat et. Curabitur nec dolor non nibh molestie lobortis dignissim non ante. Nullam rutrum lobortis ultrices. Aenean ex erat, elementum sed maximus id, posuere id quam. Proin rutrum ex elit, pretium aliquam risus finibus at. Aenean egestas orci velit, sed aliquet sapien condimentum a. Duis consequat, arcu eu viverra venenatis, dolor lorem gravida lectus, non aliquet nisi sem at augue. Donec laoreet blandit luctus. Aenean vehicula nisl vel faucibus luctus. Sed ut semper velit, vitae laoreet magna. Sed at interdum magna.

Sed iaculis faucibus odio, eu aliquam purus efficitur vel. Cras at nulla ac enim congue varius ut et nulla. Integer blandit mattis augue.

### A.3 Electrical Cycle

Lorem ipsum dolor sit amet, consectetur adipiscing elit. In faucibus condimentum rhoncus. Ut dictum nisl id risus gravida lobortis. Sed vehicula mollis tellus ut varius. Fusce eget egestas dui, et commodo dui. Proin sollicitudin interdum tempus. Nullam in elit a enim fringilla bibendum. Vestibulum sodales pellentesque condimentum. Nulla facilisi. Nunc et dolor in nulla eleifend dictum at vel ligula. Aliquam ut velit non elit ullamcorper porta ac et ex. Fusce ornare magna non nunc vestibulum, eget molestie quam dictum. In interdum aliquam odio, in posuere tellus convallis quis. Curabitur non diam elit. Proin vulputate orci diam, a tincidunt ante luctus eu. Ut a viverra ligula. Curabitur pulvinar tempus tellus eget suscipit.

Aliquam posuere massa at ante dapibus congue. Curabitur ullamcorper tortor eget consectetur aliquet. Mauris tempor magna id mauris fringilla, a varius erat blandit. Nam eleifend ullamcorper placerat. Phasellus augue tortor, volutpat bibendum lorem nec, fringilla volutpat nisl. Mauris cursus urna metus, vel eleifend orci iaculis ut. Sed sit amet scelerisque massa, quis consequat dui. Donec semper sem dui, ac placerat velit egestas vel. Nulla facilisi. Quisque tellus eros, sagittis malesuada augue ut, faucibus dictum nulla. Vestibulum non dapibus erat, ut consequat libero. Ut turpis mi, dapibus commodo libero lobortis, maximus vestibulum lectus. Vestibulum sit amet sapien dapibus, tincidunt leo in, suscipit arcu. Sed in erat bibendum, laoreet eros eu, pellentesque justo. Nulla sodales purus neque, eget maximus ipsum consequat at. Maecenas a nisl sagittis, tempus ipsum sed, dictum mauris.

Suspendisse posuere odio lacus, at auctor tortor vehicula sed. Phasellus suscipit ornare enim vitae placerat. Sed viverra purus vel sapien tempor, quis iaculis erat laoreet. Aenean vel nunc vestibulum, ornare nunc ac, mollis urna. Aenean ultrices felis ipsum, ac semper est ullamcorper in. Donec in justo varius, egestas tortor ut, venenatis augue. Duis mattis, ligula quis lacinia fringilla, tellus neque accumsan ipsum, vitae tempor metus elit vel nibh. Curabitur porttitor urna nec sapien tempor, et porttitor velit malesuada.

Suspendisse aliquam nisl quis placerat vulputate. Proin dapibus ipsum ac ante sagittis, volutpat auctor sem dapibus. Nam in facilisis odio. Integer ante mauris, eleifend et pulvinar in, venenatis quis ligula. Phasellus posuere sollicitudin tortor eget euismod. Maecenas mollis tortor eget justo vulputate sagittis. Etiam hendrerit massa quis ex molestie sodales. Quisque facilisis erat lacus, id convallis sem suscipit bibendum. Integer dui urna, pharetra sed porta sed, bibendum ut odio. Donec placerat at lectus egestas consequat. Sed id rhoncus est, vitae

vulputate sapien. Fusce tempus quam lorem, id ornare turpis sodales sed. Integer aliquet urna eget condimentum consequat. Vestibulum quis dui vel ligula posuere luctus id nec turpis.

Nam vitae placerat lacus. Mauris scelerisque interdum volutpat. Nunc aliquet tristique enim, sit amet molestie felis ullamcorper vitae. Nullam sollicitudin orci orci, in condimentum tellus consectetur in. Nam id justo justo. Fusce eget finibus est. Proin id tortor nec quam cursus vehicula. Aliquam ultrices eros eros, a tincidunt elit eleifend auctor.

Nullam consectetur dapibus ligula sit amet efficitur. Nunc non posuere sapien. Vivamus dui nisl, aliquam id ipsum non, pulvinar ornare neque. Nunc rhoncus pretium congue. Fusce id laoreet enim. Cras sed massa in eros bibendum auctor in nec sem. Nam commodo, velit id porta consequat, mi arcu gravida lorem, ut aliquam elit ante quis dui. Quisque in massa sed nibh blandit dictum.

Vestibulum molestie consectetur porttitor. Donec tincidunt vel orci at pharetra. Nullam id felis sit amet nulla tempus lacinia. Integer egestas ullamcorper massa, ut ultricies diam congue sit amet. Cras sit amet velit at nibh vehicula finibus a et lorem. Cras odio metus, venenatis ut ultrices non, ornare ac orci. Morbi et nulla dui. Mauris dictum molestie nibh, eu efficitur lorem accumsan quis.

## A.4 Cellular Electromechanical Coupling

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Nullam vitae consectetur metus, ac maximus ex. Quisque vitae ex eu lectus ultricies consequat vel non lorem. Etiam odio ipsum, tempus ut lobortis in, molestie ac leo. Vivamus mollis feugiat bibendum. Vestibulum eget venenatis quam. Aenean faucibus, massa sed ullamcorper porta, arcu nunc iaculis velit, quis consectetur purus neque placerat nibh. Vestibulum elit nunc, dignissim vulputate venenatis et, sodales non massa. Proin leo ligula, vehicula eu aliquam varius, posuere a dolor. Donec iaculis auctor neque, sit amet gravida libero porta vel. Vivamus consequat elementum lacus, at bibendum mauris egestas nec. Fusce fermentum diam eu dolor ornare, vitae vestibulum leo interdum. Morbi luctus libero quis dictum laoreet. Etiam semper porta ante, vel ullamcorper enim sodales quis.

Nullam eu nisi faucibus, fermentum ex auctor, tempor arcu. Phasellus condimentum erat mi, condimentum malesuada ligula congue venenatis. Nullam gravida imperdiet urna quis cursus. Ut tempus nec purus eget posuere. Cras non nulla sit amet justo aliquet pellentesque

nec sed eros. Nam aliquam nisl urna, in placerat magna gravida venenatis. Donec interdum vel magna ullamcorper molestie. Nunc felis neque, rhoncus fringilla faucibus sit amet, ultrices sed magna. Maecenas malesuada hendrerit diam in ultrices. Nam libero urna, volutpat ut auctor eget, interdum sed odio. Vestibulum suscipit mauris nec augue ornare, ut eleifend nulla gravida. Proin imperdiet, mauris quis consectetur porta, leo dui convallis leo, id lobortis massa diam eu libero. Aenean hendrerit vel ante aliquam venenatis. Pellentesque bibendum pretium odio, ut sagittis lectus feugiat a. Donec porttitor vulputate lacus.

Nunc volutpat efficitur lacus in aliquet. Nullam non iaculis diam, at ultrices diam. Proin vehicula vulputate cursus. Morbi tempus sapien id urna lobortis interdum. Maecenas elementum sagittis elementum. Donec at sodales velit, a posuere tortor. Nulla id hendrerit tortor. Sed semper velit in magna sagittis pulvinar. Nulla nec arcu molestie, ultricies sapien sit amet, sollicitudin nisi. Donec nisi massa, suscipit ut dignissim quis, lacinia id leo.

Suspendisse ut mi metus. Morbi tincidunt ligula in porttitor consectetur. Integer eu urna urna. Suspendisse potenti. Mauris sit amet felis eu diam auctor ullamcorper. Morbi in porta nisi. Nam ante tortor, venenatis vitae tempor sed, sagittis vitae velit. In semper orci sit amet nisi ullamcorper varius. Aenean dignissim ultrices imperdiet. Maecenas lacinia enim id neque porttitor iaculis. Curabitur laoreet ante ut urna dignissim, id sollicitudin metus consectetur. Aenean massa ipsum, auctor vel ante vel, blandit dignissim libero. Fusce interdum ac magna et interdum.

*The first kind of intellectual and artistic personality  
belongs to the hedgehogs, the second to the foxes ...*

— Sir Isaiah Berlin ?

## Bibliography

- Adams, D. K. and Comrie, A. C. (1997), ‘The north American monsoon’, *Bulletin of the American Meteorological Society* **78**(10), 2197–2214.
- Adams, D. K., Gutman, S. I., Holub, K. L. and Pereira, D. S. (2013), ‘Gnss observations of deep convective time scales in the amazon’, *Geophysical Research Letters* **40**(11), 2818–2823.  
**URL:** <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/grl.50573>
- Adler, R. F., Huffman, G. J., Chang, A., Ferraro, R., Xie, P.-P., Janowiak, J., Rudolf, B., Schneider, U., Curtis, S., Bolvin, D., Gruber, A., Susskind, J., Arkin, P. and Nelkin, E. (2003), ‘The version-2 global precipitation climatology project (gpcp) monthly precipitation analysis (1979–present)’, *Journal of hydrometeorology* **4**(6), 1147–1167.
- Amador, J. A. (2008), ‘The intra-americas sea low-level jet: Overview and future research’, *Annals of the New York Academy of Sciences* **1146**(1), 153–188.
- Anderson, T. G., Anchukaitis, K. J., Pons, D. and Taylor, M. (2019), ‘Multiscale trends and precipitation extremes in the central american midsummer drought’, *Environmental Research Letters* **14**(12), 124016.
- Andrews, M., Ridley, J., Wood, R., Andrews, T., Blockley, E., Booth, B., Burke, E., Dittus, A., Florek, P., Gray, L., Haddad, S., Hardiman, S., Hermanson, L., Hodson, D., Hogan, E., Jones, G., Knight, J., Kuhlbrodt, T., Misios, S. and Sutton, R. (2020), ‘Historical simulations with hadgem3-gc3.1 for cmip6’, *Journal of Advances in Modeling Earth Systems* .
- Andrews, T., Andrews, M. B., Bodas-Salcedo, A., Jones, G. S., Kuhlbrodt, T., Manners, J., Menary, M. B., Ridley, J., Ringer, M. A., Sellar, A. A., Senior, C. A. and Tang, Y. (2019), ‘Forcings, feedbacks, and climate sensitivity in hadgem3-gc3. 1 and ukesm1’, *Journal of Advances in Modeling Earth Systems* **11**(12), 4377–4394.
- Angeles, M. E., González, J. E., Ramírez-Beltrán, N. D., Tepley, C. A. and Comarazamy, D. E. (2010), ‘Origins of the caribbean rainfall bimodal behavior’, *Journal of Geophysical Research: Atmospheres* **115**(D11).
- Baldwin, M., Gray, L., Dunkerton, T., Hamilton, K., Haynes, P., Randel, W., Holton, J., Alexander, M., Hirota, I., Horinouchi, T. et al. (2001), ‘The quasi-biennial oscillation’, *Reviews of Geophysics* **39**(2), 179–229.
- Becker, A., Finger, P., Meyer-Christoffer, A., Rudolf, B. and Ziese, M. (2011), ‘Gpcc full data reanalysis version 6.0 at 1.0: monthly land-surface precipitation from rain-gauges built on gts-based and historic data’, *Global Precipitation Climatology Centre (GPCC): Berlin, Germany* .
- Bellucci, A., Gualdi, S. and Navarra, A. (2010), ‘The double-itcz syndrome in coupled general circulation models: The role of large-scale vertical circulation regimes’, *Journal of Climate*

- 23**(5), 1127–1145.
- Bischoff, T. and Schneider, T. (2016), ‘The Equatorial Energy Balance, ITCZ Position, and Double-ITCZ Bifurcations’, *Journal of Climate* **29**(8), 2997–3013.
- URL:** <https://doi.org/10.1175/JCLI-D-15-0328.1>
- Bjerknes, J. (1969), ‘Atmospheric teleconnections from the equatorial pacific’, *Mon. Wea. Rev* **97**(3), 163–172.
- C3S (2017), ‘ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate.’, *Copernicus Climate Change Service Climate Data Store (CDS)*. date of access: 18 March 2019, <https://climate.copernicus.eu/climate-reanalysis>.
- Cai, W., Santoso, A., Wang, G., Yeh, S.-W., An, S.-I., Cobb, K. M., Collins, M., Guilyardi, E., Jin, F.-F., Kug, J.-S. et al. (2015a), ‘Enso and greenhouse warming’, *Nature Climate Change* **5**(9), 849.
- Cai, W., Wu, L., Lengaigne, M., Li, T., McGregor, S., Kug, J.-S., Yu, J.-Y., Stuecker, M. F., Santoso, A., Li, X., Ham, Y.-G., Chikamoto, Y., Ng, B., McPhaden, M. J., Du, Y., Dommenech, D., Jia, F., Kajtar, J. B., Keenlyside, N., Lin, X., Luo, J.-J., Martín-Rey, M., Ruprich-Robert, Y., Wang, G., Xie, S.-P., Yang, Y., Kang, S. M., Choi, J.-Y., Gan, B., Kim, G.-I., Kim, C.-E., Kim, S., Kim, J.-H. and Chang, P. (2019), ‘Pantropical climate interactions’, *Science* **363**(6430), eaav4236.
- Carvalho, L. M. V., Jones, C., Posadas, A. N. D., Quiroz, R., Bookhagen, B. and Liebmann, B. (2012), ‘Precipitation characteristics of the south american monsoon system derived from multiple datasets’, *Journal of Climate* **25**(13), 4600–4620.
- URL:** <https://doi.org/10.1175/JCLI-D-11-00335.1>
- Chan, J. C. (1995), ‘Tropical cyclone activity in the western north pacific in relation to the stratospheric quasi-biennial oscillation’, *Monthly Weather Review* **123**(8), 2567–2571.
- Christensen, H., Berner, J., Coleman, D. R. and Palmer, T. (2017), ‘Stochastic parameterization and el niño–southern oscillation’, *Journal of Climate* **30**(1), 17–38.
- Chuvieco, E., Opazo, S., Sione, W., Valle, H. d., Anaya, J., Bella, C. D., Cruz, I., Manzo, L., López, G., Mari, N., González-Alonso, F., Morelli, F., Setzer, A., Csiszar, I., Kanpandegi, J. A., Bastarrika, A. and Libonati, R. (2008), ‘Global burned-land estimation in latin-america using modis composite data’, *Ecological Applications* **18**(1), 64–79.
- URL:** <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/06-2148.1>
- Collimore, C. C., Martin, D. W., Hitchman, M. H., Huesmann, A. and Waliser, D. E. (2003), ‘On the relationship between the qbo and tropical deep convection’, *Journal of climate* **16**(15), 2552–2568.
- Corrales-Suastegui, A., Fuentes-Franco, R. and Pavia, E. G. (2020), ‘The mid-summer drought over mexico and central america in the 21st century’, *International Journal of Climatology* **40**(3), 1703–1715.
- Curtis, S. and Gamble, D. W. (2008), ‘Regional variations of the caribbean mid-summer drought’, *Theoretical and Applied Climatology* **94**, 25–34.
- de Sousa, K., Casanoves, F., Sellare, J., Ospina, A., Suchini, J. G., Aguilar, A. and Mercado, L. (2018), ‘How climate awareness influences farmers’ adaptation decisions in central america?’, *Journal of Rural Studies* **64**, 11–19.
- Dilley, M. (1996), ‘Synoptic controls on precipitation in the valley of oaxaca, mexico’, *International Journal of Climatology: A Journal of the Royal Meteorological Society* **16**(9), 1019–1031.

- Dinku, T., Ruiz, F., Connor, S. J. and Ceccato, P. (2010), ‘Validation and intercomparison of satellite rainfall estimates over colombia’, *Journal of Applied Meteorology and Climatology* **49**(5), 1004–1014.
- Donohoe, A., Marshall, J., Ferreira, D. and Mcgee, D. (2013), ‘The Relationship between ITCZ Location and Cross-Equatorial Atmospheric Heat Transport: From the Seasonal Cycle to the Last Glacial Maximum’, *Journal of Climate* **26**(11), 3597–3618.
- URL:** <https://doi.org/10.1175/JCLI-D-12-00467.1>
- Durán-Quesada, A. M., Gimeno, L. and Amador, J. (2017), ‘Role of moisture transport for central american precipitation’, *Earth System Dynamics* **8**(1), 147–161.
- URL:** <https://www.earth-syst-dynam.net/8/147/2017/>
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J. and Taylor, K. E. (2016), ‘Overview of the coupled model intercomparison project phase 6 (cmip6) experimental design and organization’, *Geoscientific Model Development (Online)* **9**.
- Franchito, S. H., Rao, V. B., Vasques, A. C., Santo, C. M. and Conforte, J. C. (2009), ‘Validation of trmm precipitation radar monthly rainfall estimates over brazil’, *Journal of Geophysical Research: Atmospheres* **114**(D2).
- Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A. and Michaelsen, J. (2015), ‘The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes’, *Scientific data* **2**, 150066.
- Gadgil, S. (2018), ‘The monsoon system: Land–sea breeze or the itcz?’, *Journal of Earth System Science* **127**(1), 1.
- Gamble, D. W., Parnell, D. B. and Curtis, S. (2008), ‘Spatial variability of the caribbean mid-summer drought and relation to north atlantic high circulation’, *International Journal of Climatology* **28**(3), 343–350.
- Gan, M., Kousky, V. and Ropelewski, C. (2004), ‘The South America monsoon circulation and its relationship to rainfall over west-central Brazil’, *Journal of climate* **17**, 47–66.
- García-Martínez, I. M. and Bollasina, M. A. (2020), ‘Sub-monthly evolution of the caribbean low-level jet and its relationship with regional precipitation and atmospheric circulation’, *Climate Dynamics* pp. 1–18.
- Geil, K. L., Serra, Y. L. and Zeng, X. (2013), ‘Assessment of cmip5 model simulations of the north american monsoon system’, *Journal of Climate* **26**(22), 8787–8801.
- Giannini, A., Kushnir, Y. and Cane, M. A. (2000), ‘Interannual variability of Caribbean rainfall, ENSO, and the Atlantic Ocean’, *Journal of Climate* **13**, 297–311.
- Gill, A. E. (1980), ‘Some simple solutions for heat-induced tropical circulation’, *Quarterly Journal of the Royal Meteorological Society* **106**(449), 447–462.
- Giorgetta, M. A., Bengtsson, L. and Arpe, K. (1999), ‘An investigation of qbo signals in the east asian and indian monsoon in gcm experiments’, *Climate dynamics* **15**(6), 435–450.
- Guilyardi, E., Wittenberg, A., Fedorov, A., Collins, M., Wang, C., Capotondi, A., Van Oldenborgh, G. J. and Stockdale, T. (2009), ‘Understanding el niño in ocean–atmosphere general circulation models: Progress and challenges’, *Bulletin of the American Meteorological Society* **90**(3), 325–340.
- Harris, I., Jones, P. D., Osborn, T. J. and Lister, D. H. (2014), ‘Updated high-resolution grids of monthly climatic observations—the cru ts3. 10 dataset’, *International journal of climatology*

- 34(3), 623–642.
- Harvey, C. A., Saborio-Rodríguez, M., Martínez-Rodríguez, M. R., Viguera, B., Chain-Guadarrama, A., Vignola, R. and Alpizar, F. (2018), ‘Climate change impacts and adaptation among smallholder farmers in central america’, *Agriculture & Food Security* 7(1), 57.
- Hellin, J., Cox, R. and López-Ridaura, S. (2017), ‘Maize Diversity, Market Access, and Poverty Reduction in the Western Highlands of Guatemala’, *Mountain Research and Development* 37(2), 188 – 197.
- Herrera, E., Magaña, V. and Caetano, E. (2015), ‘Air–sea interactions and dynamical processes associated with the midsummer drought’, *International Journal of Climatology* 35(7), 1569–1578.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S. and Thépaut, J.-N. (2020), ‘The era5 global reanalysis’, *Quarterly Journal of the Royal Meteorological Society* 1(n/a).
- URL:** <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3803>
- Hidalgo, H. G., Durán-Quesada, A. M., Amador, J. A. and Alfaro, E. J. (2015), ‘The caribbean low-level jet, the inter-tropical convergence zone and precipitation patterns in the intra-americas sea: a proposed dynamical mechanism’, *Geografiska Annaler: Series A, Physical Geography* 97(1), 41–59.
- URL:** <https://doi.org/10.1111/geoa.12085>
- Hoskins, B., Yang, G.-Y. and Fonseca, R. (2020), ‘The detailed dynamics of the june–august hadley cell’, *Quarterly Journal of the Royal Meteorological Society* 146(727), 557–575.
- Huffman, G. J., Adler, R. F., Bolvin, D. T. and Nelkin, E. J. (2010), The trmm multi-satellite precipitation analysis (tmpa), in ‘Satellite rainfall applications for surface hydrology’, Springer, pp. 3–22.
- Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P. and Stocker, E. F. (2007), ‘The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales’, *Journal of Hydrometeorology* 8(1), 38–55.
- URL:** <https://doi.org/10.1175/JHM560.1>
- Jones, C. and Carvalho, L. M. (2013), ‘Climate change in the South American monsoon system: present climate and CMIP5 projections’, *Journal of Climate* 26(17), 6660–6678.
- Karnauskas, K. B., Seager, R., Giannini, A. and Busalacchi, A. J. (2013), ‘A simple mechanism for the climatological midsummer drought along the Pacific coast of Central America’, *Atmosfera* 26, 261–281.
- Kennedy, J., Rayner, N., Smith, R., Parker, D. and Saunby, M. (2011), ‘Reassessing biases and other uncertainties in sea surface temperature observations measured in situ since 1850: 2. biases and homogenization’, *Journal of Geophysical Research: Atmospheres* 116(D14).
- Kim, H., Caron, J. M., Richter, J. H. and Simpson, I. R. (2020), ‘The lack of qbo-mjo connection in cmip6 models’, *Geophysical Research Letters* 47(11), e2020GL087295. e2020GL087295

- 2020GL087295.
- URL:** <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020GL087295>
- Kuhlbrodt, T., Jones, C. G., Sellar, A., Storkey, D., Blockley, E., Stringer, M., Hill, R., Graham, T., Ridley, J., Blaker, A., Calvert, D., Copsey, D., Ellis, R., Hewitt, H., Hyder, P., Ineson, S., Mulcahy, J., Siahaan, A. and Walton, J. (2018), 'The low-resolution version of hadgem3 gc3. 1: Development and evaluation for global climate', *Journal of advances in modeling earth systems* **10**(11), 2865–2888.
- Lawrence, P. J., Feddema, J. J., Bonan, G. B., Meehl, G. A., O'Neill, B. C., Oleson, K. W., Levis, S., Lawrence, D. M., Kluzek, E., Lindsay, K. and Thornton, P. E. (2012), 'Simulating the biogeochemical and biogeophysical impacts of transient land cover change and wood harvest in the community climate system model (ccsm4) from 1850 to 2100', *Journal of Climate* **25**(9), 3071–3095.
- Lee, J. C. K. and Klingaman, N. P. (2018), 'The effect of the quasi-biennial oscillation on the madden-julian oscillation in the met office unified model global ocean mixed layer configuration', *Atmospheric Science Letters* **19**(5), e816.
- URL:** <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/asl.816>
- Li, G. and Xie, S.-P. (2014), 'Tropical biases in cmip5 multimodel ensemble: The excessive equatorial pacific cold tongue and double itcz problems', *Journal of Climate* **27**(4), 1765–1780.
- Lorenz, E. N. (1967), *The nature and theory of the general circulation of the atmosphere*, Vol. 218, World Meteorological Organization Geneva.
- Machado, L., Laurent, H., Dessay, N. and Miranda, I. (2004), 'Seasonal and diurnal variability of convection over the amazonia: A comparison of different vegetation types and large scale forcing', *Theoretical and Applied Climatology* **78**(1-3), 61–77.
- Magaña, V., Amador, J. A. and Medina, S. (1999), 'The midsummer drought over Mexico and Central America', *Journal of Climate* **12**, 1577–1588.
- Magaña, V. and Caetano, E. (2005), 'Temporal evolution of summer convective activity over the Americas warm pools', *Geophysical Research Letters* **32**(2).
- Maldonado, T., Alfaro, E., Rutgersson, A. and Amador, J. A. (2017), 'The early rainy season in central america: the role of the tropical north atlantic ssts', *International Journal of Climatology* **37**(9), 3731–3742.
- Maldonado, T., Rutgersson, A., Amador, J., Alfaro, E. and Claremar, B. (2016), 'Variability of the caribbean low-level jet during boreal winter: large-scale forcings', *International Journal of Climatology* **36**, 1954–1969.
- Mapes, B. E., Liu, P. and Buennning, N. (2005), 'Indian monsoon onset and the americas midsummer drought: Out-of-equilibrium responses to smooth seasonal forcing', *Journal of Climate* **18**(7), 1109–1115.
- Marengo, J. A., Liebmann, B., Grimm, A. M., Misra, V., Silva Dias, P. L., Cavalcanti, I. F. A., Carvalho, L. M. V., Berbery, E. H., Ambrizzi, T., Vera, C. S., Saulo, A. C., Nogues-Paegle, J., Zipser, E., Seth, A. and Alves, L. M. (2012), 'Recent developments on the South American monsoon system', *International Journal of Climatology* **32**, 1–21.
- Marengo, J. A., Liebmann, B., Kousky, V. E., Filizola, N. P. and Wainer, I. C. (2001), 'Onset and end of the rainy season in the brazilian amazon basin', *Journal of Climate* **14**(5), 833–852.
- Martin, Z., Vitart, F., Wang, S. and Sobel, A. (2020), 'The impact of the stratosphere on the mjo in a

- forecast model', *Journal of Geophysical Research: Atmospheres* **125**(4), e2019JD032106. e2019JD032106 2019JD032106.
- URL:** <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD032106>
- Martin, Z., Wang, S., Nie, J. and Sobel, A. (2019), 'The Impact of the QBO on MJO Convection in Cloud-Resolving Simulations', *Journal of the Atmospheric Sciences* **76**(3), 669–688.
- URL:** <https://doi.org/10.1175/JAS-D-18-0179.1>
- Martinez, C., Goddard, L., Kushnir, Y. and Ting, M. (2019), 'Seasonal climatology and dynamical mechanisms of rainfall in the caribbean', *Climate dynamics* **53**(1-2), 825–846.
- Martinez, C., Kushnir, Y., Goddard, L. and Ting, M. (2020), 'Interannual variability of the early and late-rainy seasons in the caribbean', *Climate Dynamics* **55**(5), 1563–1583.
- Menary, M. B., Kuhlbrodt, T., Ridley, J., Andrews, M. B., Dimdore-Miles, O. B., Deshayes, J., Eade, R., Gray, L., Ineson, S., Mignot, J., Roberts, C. D., Robson, J., Wood, R. A. and Xavier, P. (2018), 'Preindustrial control simulations with hadgem3-gc3. 1 for cmip6', *Journal of Advances in Modeling Earth Systems* **10**(12), 3049–3075.
- Mosiño, A. and García, E. (1966), Evaluación de la sequía intraestival en la república mexicana, in 'Proc. Conf. Reg. Latinoamericana Unión Geogr. Int', Vol. 3, pp. 500–516.
- Mulcahy, J. P., Jones, C., Sellar, A., Johnson, B., Boutle, I. A., Jones, A., Andrews, T., Rumbold, S. T., Molland, J., Bellouin, N., Johnson, C. E., Williams, K. D., Grosvenor, D. P. and McCoy, D. T. (2018), 'Improved aerosol processes and effective radiative forcing in hadgem3 and ukesm1', *Journal of Advances in Modeling Earth Systems* **10**(11), 2786–2805.
- Muñoz, E., Busalacchi, A. J., Nigam, S. and Ruiz-Barradas, A. (2008), 'Winter and summer structure of the caribbean low-level jet', *Journal of Climate* **21**(6), 1260–1276.
- Neelin, J. D., Battisti, D. S., Hirst, A. C., Jin, F.-F., Wakata, Y., Yamagata, T. and Zebiak, S. E. (1998), 'ENSO theory', *Journal of Geophysical Research: Oceans* **103**(C7), 14261–14290.
- Nie, J. and Sobel, A. H. (2015), 'Responses of tropical deep convection to the qbo: Cloud-resolving simulations', *Journal of the Atmospheric Sciences* **72**(9), 3625–3638.
- Nieto-Ferreira, R. and Rickenbach, T. M. (2011), 'Regionality of monsoon onset in South America: a three-stage conceptual model', *International Journal of Climatology* **31**(9), 1309–1321.
- Ordoñez, P., Nieto, R., Gimeno, L., Ribera, P., Gallego, D., Ochoa-Moya, C. A. and Quintanar, A. I. (2019), 'Climatological moisture sources for the western north american monsoon through a lagrangian approach: their influence on precipitation intensity', *Earth System Dynamics* **10**(1), 59–72.
- Oueslati, B. and Bellon, G. (2013), 'Convective entrainment and large-scale organization of tropical precipitation: Sensitivity of the cnrm-cm5 hierarchy of models', *Journal of climate* **26**(9), 2931–2946.
- Pascale, S. and Bordoni, S. (2016), 'Tropical and extratropical controls of gulf of california surges and summertime precipitation over the southwestern united states', *Monthly Weather Review* **144**(7), 2695–2718.
- Perdigón-Morales, J., Romero-Centeno, R., Barrett, B. S. and Ordoñez, P. (2019), 'Intraseasonal variability of summer precipitation in mexico: Mjo influence on the midsummer drought', *Journal of Climate* (2019).
- Perdigón-Morales, J., Romero-Centeno, R., Ordóñez, P. and Barrett, B. S. (2018), 'The midsummer

- drought in mexico: perspectives on duration and intensity from the chirps precipitation database', *International Journal of Climatology* **38**, 2174–2186.
- Rauscher, S. A., Giorgi, F., Diffenbaugh, N. S. and Seth, A. (2008), 'Extension and intensification of the Meso-American mid-summer drought in the twenty-first century', *Climate Dynamics* **31**, 551–571.
- Richter, J. H., Anstey, J. A., Butchart, N., Kawatani, Y., Meehl, G. A., Osprey, S. and Simpson, I. R. (2020), 'Progress in simulating the quasi-biennial oscillation in cmip models', *Journal of Geophysical Research: Atmospheres* **125**(8), e2019JD032362. e2019JD032362 10.1029/2019JD032362.  
**URL:** <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JD032362>
- Ridley, J., Menary, M., Kuhlbrodt, T., Andrews, M. and Andrews, T. (2018), 'Mohc hadgem3-gc31-ll model output prepared for cmip6 cmip picontrol'.  
**URL:** <https://doi.org/10.22033/ESGF/CMIP6.6294>
- Ridley, J., Menary, M., Kuhlbrodt, T., Andrews, M. and Andrews, T. (2019a), 'Mohc hadgem3-gc31-ll model output prepared for cmip6 cmip amip'.  
**URL:** <https://doi.org/10.22033/ESGF/CMIP6.5853>
- Ridley, J., Menary, M., Kuhlbrodt, T., Andrews, M. and Andrews, T. (2019b), 'Mohc hadgem3-gc31-ll model output prepared for cmip6 cmip historical'.  
**URL:** <https://doi.org/10.22033/ESGF/CMIP6.6109>
- Ridley, J., Menary, M., Kuhlbrodt, T., Andrews, M. and Andrews, T. (2019c), 'Mohc hadgem3-gc31-mm model output prepared for cmip6 cmip picontrol'.  
**URL:** <https://doi.org/10.22033/ESGF/CMIP6.6297>
- Ropelewski, C. F. and Halpert, M. S. (1986), 'North american precipitation and temperature patterns associated with the el niño/southern oscillation (enso)', *Monthly Weather Review* **114**(12), 2352–2362.
- Ropelewski, C. F. and Halpert, M. S. (1987), 'Global and regional scale precipitation patterns associated with the el niño/southern oscillation', *Monthly weather review* **115**(8), 1606–1626.
- Ryu, J.-H. and Hayhoe, K. (2014), 'Understanding the sources of Caribbean precipitation biases in CMIP3 and CMIP5 simulations', *Climate dynamics* **42**(11-12), 3233–3252.
- Santoso, A., Mcphaden, M. J. and Cai, W. (2017), 'The defining characteristics of enso extremes and the strong 2015/2016 el niño', *Reviews of Geophysics* **55**(4), 1079–1129.
- Schneider, T., Bischoff, T. and Haug, G. H. (2014), 'Migrations and dynamics of the Intertropical Convergence Zone', *Nature* **513**(7516), 45.
- Sellar, A. A., Jones, C. G., Mulcahy, J., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahaan, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G. and Zerroukat, M. (2019), 'Ukesm1: Description and evaluation of the uk earth system model', *Journal of Advances in Modeling Earth Systems* **0(ja)**.

- URL:** <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019MS001739>
- Son, S.-W., Lim, Y., Yoo, C., Hendon, H. H. and Kim, J. (2017), ‘Stratospheric control of the Madden–Julian oscillation’, *Journal of Climate* **30**(6), 1909–1922.
- Stensrud, D. J., Gall, R. L. and Nordquist, M. K. (1997), ‘Surges over the gulf of California during the Mexican monsoon’, *Monthly Weather Review* **125**(4), 417–437.
- Straffon, A., Zavala-Hidalgo, J. and Estrada, F. (2019), ‘Preconditioning of the precipitation interannual variability in southern Mexico and Central America by oceanic and atmospheric anomalies’, *International Journal of Climatology* .
- Sulca, J., Takahashi, K., Espinoza, J.-C., Vuille, M. and Lavado-Casimiro, W. (2018), ‘Impacts of different ENSO flavors and tropical Pacific convection variability (ITCZ, SPCZ) on austral summer rainfall in South America, with a focus on Peru’, *International Journal of Climatology* **38**(1), 420–435.
- Tang, Y., Rumbold, S., Ellis, R., Kelley, D., Mulcahy, J., Sellar, A., Walton, J. and Jones, C. (2019a), ‘Mohc ukcsm1.0-ll model output prepared for cmip6 cmip historical’.
- URL:** <https://doi.org/10.22033/ESGF/CMIP6.6113>
- Tang, Y., Rumbold, S., Ellis, R., Kelley, D., Mulcahy, J., Sellar, A., Walton, J. and Jones, C. (2019b), ‘Mohc ukcsm1.0-ll model output prepared for cmip6 cmip picontrol’.
- URL:** <https://doi.org/10.22033/ESGF/CMIP6.6298>
- Tegtmeier, S., Anstey, J., Davis, S., Dragani, R., Harada, Y., Ivanciu, I., Pilch Kedzierski, R., Krüger, K., Legras, B., Long, C., Wang, J. S., Wargan, K. and Wright, J. S. (2020), ‘Temperature and tropopause characteristics from reanalyses data in the tropical tropopause layer’, *Atmospheric Chemistry and Physics* **20**(2), 753–770.
- URL:** <https://www.atmos-chem-phys.net/20/753/2020/>
- Trejo, F. J. P., Barbosa, H. A., Peñaloza-Murillo, M. A., Moreno, M. A. and Farías, A. (2016), ‘Intercomparison of improved satellite rainfall estimation with chirps gridded product and rain gauge data over Venezuela’, *Atmósfera* **29**(4), 323–342.
- Trenberth, K. E. (1997), ‘The definition of El Niño’, *Bulletin of the American Meteorological Society* **78**(12), 2771–2778.
- Vera, C., Higgins, W., Amador, J., Ambrizzi, T., Garreaud, R., Gochis, D., Gutzler, D., Lettenmaier, D., Marengo, J., Mechoso, C. R., Nogues-Paegle, J., Dias, P. L. S. and Zhang, C. (2006), ‘Toward a unified view of the American monsoon systems’, *Journal of Climate* **19**(20), 4977–5000.
- Walters, D., Boutle, I., Brooks, M., Melvin, T., Stratton, R., Vosper, S., Wells, H., Williams, K., Wood, N., Allen, T., Bushell, A., Copsey, D., Earnshaw, P., Edwards, J., Gross, M., Hardiman, S., Harris, C., Heming, J., Klingaman, N., Levine, R., Manners, J., Martin, G., Milton, S., Mittermaier, M., Morcrette, C., Riddick, T., Roberts, M., Sanchez, C., Selwood, P., Stirling, A., Smith, C., Suri, D., Tennant, W., Vidale, P. L., Wilkinson, J., Willett, M., Woolnough, S. and Xavier, P. (2019), ‘The Met Office unified model global atmosphere 7.0/7.1 and JULES global land 7.0 configurations’, *Geoscientific Model Development* **12**(5), 1909–1963.
- Wang, C. and Picaut, J. (2004), ‘Understanding ENSO physics—A review’, *Earth’s Climate: The Ocean–Atmosphere Interaction, Geophys. Monogr.* **147**, 21–48.
- Wang, P. X., Wang, B., Cheng, H., Fasullo, J., Guo, Z., Kiefer, T. and Liu, Z. (2017), ‘The global

- monsoon across time scales: Mechanisms and outstanding issues', *Earth-Science Reviews* **174**, 84–121.
- Wang, S., Tippett, M. K., Sobel, A. H., Martin, Z. and Vitart, F. (2019), 'Impact of the qbo on prediction and predictability of the mjo convection', *J. Geophys. Res. Atmos.*
- Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder, P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann, R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T. and Xavier, P. K. (2018), 'The met office global coupled model 3.0 and 3.1 (gc3. 0 and gc3. 1) configurations', *Journal of Advances in Modeling Earth Systems* **10**(2), 357–380.
- Xie, P. and Arkin, P. A. (1997), 'Global precipitation: A 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs', *Bulletin of the American Meteorological Society* **78**, 2539–2558.
- Xie, P., Arkin, P. A. and Janowiak, J. E. (2007), *CMAP: The CPC Merged Analysis of Precipitation*, Springer Netherlands, Dordrecht, pp. 319–328.
- Zhou, T., Turner, A. G., Kinter, J. L., Wang, B., Qian, Y., Chen, X., Wu, B., Liu, B., Zou, L. and Bian, H. (2016), 'GMMIP (v1. 0) contribution to CMIP6: global monsoons model inter-comparison project', *Geoscientific Model Development* **9**, 3589–3604.