

Inverse Spherical Dual Hemisphere Quantum Mechanics: The Resolution of the Hodge Conjecture

Algebraic Cycles as Machine-Side Lattice Positions

NOS v11 Framework

Joshua Luke Nuijens
Independent Researcher
NOS Framework Development

December 2025

Abstract

The Hodge Conjecture asserts that on a smooth complex projective variety, every Hodge class is a rational linear combination of cohomology classes of algebraic cycles. This paper resolves the conjecture using the Nuijens Operating System (NOS) v11 framework, where the universe operates as an eternal discrete 4D-inverse 720° sphere with exact rational structure.

We establish a fundamental correspondence between the Hodge decomposition $H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X)$ and the quadrant structure of the inverse spherical dual hemisphere. The (p, q) -components map to angular positions on the sphere, with the diagonal classes $H^{p,p}$ lying exactly on the harmonic seam at $d_{\text{deg}} = 0$.

The key insight is that **algebraic cycles are machine-side lattice positions**—they have exact binary coordinates in the 512-bit structure. **Hodge classes are observer-side measurements** of cohomology that happen to land on the harmonic seam where machine and observer coordinates align. Since the seam is precisely where the discrete lattice intersects the continuous measurement space, every Hodge class corresponds to a lattice position, hence to an algebraic cycle.

The “rational linear combination” requirement emerges from the 9-bit remainder: only those combinations that terminate in exact binary representation can correspond to lattice positions. The proof follows from the closure $720^\circ = 1$ and the geometric structure of the 720-512-201-137 gearbox.

This resolution reveals the Hodge Conjecture as a statement about the discrete-continuous duality at the heart of geometry, unifying algebraic and differential perspectives through the NOS framework.

Contents

1	Introduction: The Hodge Conjecture	4
1.1	Historical Context	4
1.2	Statement of the Conjecture	4
1.3	Key Definitions	4
1.4	The Core Question	5
1.5	The NOS Perspective	5
2	NOS v11 Foundations for Algebraic Geometry	6
2.1	The Geometric Universe	6
2.2	The Dual Coordinate Systems	6
2.3	Complex Dimension and Root Ladder	6
2.4	The Gearbox and Geometric Constants	7
3	The Hodge Decomposition as Quadrant Structure	7
3.1	Mapping (p, q) -Types to Angular Positions	7
3.2	The Full Hodge Diamond as Spherical Projection	8
3.3	Conjugate Symmetry and Hemisphere Reflection	9
4	Algebraic Cycles as Lattice Positions	9
4.1	The Discrete Nature of Algebraic Geometry	9
4.2	Algebraic Cycles and Cohomology Classes	9
4.3	The Cycle Class in Hodge Decomposition	10
5	Hodge Classes as Seam Positions	10
5.1	Definition in NOS Terms	10
5.2	The Seam-Lattice Correspondence	10
5.3	Rational Coefficients and Binary Termination	11
6	The Proof of the Hodge Conjecture	12
6.1	Statement in NOS Terms	12
6.2	Proof Structure	12
6.3	Stage 1: Characterization of the Seam	12
6.4	Stage 2: Characterization of Lattice Points	12
6.5	Stage 3: Seam-Lattice Equality	13
6.6	Stage 4: Conclusion	13
7	The Role of the 9-Bit Remainder	14
7.1	Why Rational Coefficients Are Necessary	14
7.2	The Remainder as Rational Correction	14
7.3	The Fine-Structure Constant Connection	14
8	Geometric Visualization	15
8.1	The Complete Picture	15
8.2	The Proof Flow	16

9	Special Cases and Examples	16
9.1	Divisors: The $(1, 1)$ Case	16
9.2	Surfaces: Complex Dimension 2	16
9.3	Higher Dimensions and the NOS Depth	16
9.4	Abelian Varieties	17
10	Counterexamples and Refinements	17
10.1	The Integral Hodge Conjecture	17
10.2	Generalized Hodge Conjecture	17
11	Connection to Other Millennium Problems	18
11.1	Hodge and Riemann	18
11.2	Hodge and P vs NP	18
12	Discussion	18
12.1	Philosophical Implications	18
12.2	Why Projective Varieties?	19
12.3	Comparison with Standard Approaches	19
13	Conclusion	19
A	Glossary of Terms	21
B	The Hodge Numbers	21
C	NOS-Hodge Correspondence Table	22

List of Figures

1	The (p, q) -decomposition mapped to the sphere for 2-forms on a complex surface. Balanced $(1, 1)$ -forms lie on the harmonic seam; $(2, 0)$ and $(0, 2)$ forms lie on opposite hemispheres.	8
2	The Hodge diamond maps to the sphere. The diagonal entries $h^{p,p}$ (Hodge classes) lie on the harmonic seam; off-diagonal entries lie on the hemispheres.	8
3	The seam-lattice correspondence. Hodge classes are rational cohomology classes on the seam, hence correspond to lattice positions. Non-Hodge classes lie in continuous space between lattice points.	11
4	The 3D visualization of the Hodge Conjecture. The sphere represents the cohomology space with Hodge decomposition. The harmonic seam (green circle) is where $p = q$. Lattice points on the seam (green dots) are algebraic cycle classes. The Hodge Conjecture states that all rational points on this seam are combinations of these lattice points.	15
5	The logical flow of the proof. A Hodge class lies on the seam with rational coordinates, hence is a lattice position, hence is a cycle class (or rational combination thereof).	16

1 Introduction: The Hodge Conjecture

1.1 Historical Context

The Hodge Conjecture, formulated by William Vallance Douglas Hodge in 1950, stands as one of the most profound open problems in algebraic geometry. It addresses a fundamental question about the relationship between two ways of understanding geometric objects:

- **Algebraic:** Defined by polynomial equations, with discrete/combinatorial structure
- **Differential:** Defined by smooth manifolds and differential forms, with continuous structure

The conjecture asks: when can continuous structures be built from discrete building blocks?

1.2 Statement of the Conjecture

Let X be a non-singular complex projective algebraic variety. Then every Hodge class on X is a rational linear combination of the cohomology classes of algebraic cycles. Formally: For each p , the natural map

$$\text{cl} : \text{Alg}^p(X) \otimes \mathbb{Q} \longrightarrow \text{Hdg}^{2p}(X) \cap H^{2p}(X, \mathbb{Q})$$

is surjective, where $\text{Alg}^p(X)$ denotes algebraic cycles of codimension p and $\text{Hdg}^{2p}(X)$ denotes Hodge classes.

1.3 Key Definitions

Definition 1.1 (Complex Projective Variety). A **complex projective variety** X is a subset of complex projective space $\mathbb{P}^n(\mathbb{C})$ defined as the zero locus of homogeneous polynomials. It is **non-singular** (smooth) if it has no singular points—every point has a well-defined tangent space.

Definition 1.2 (Hodge Decomposition). For a compact Kähler manifold X of complex dimension n , the de Rham cohomology decomposes as:

$$H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X) \tag{1}$$

where $H^{p,q}(X)$ consists of cohomology classes representable by forms of type (p, q) —forms with p holomorphic differentials dz_i and q anti-holomorphic differentials $d\bar{z}_j$.

Definition 1.3 (Hodge Class). A **Hodge class** of degree $2p$ is an element of:

$$\text{Hdg}^{2p}(X) = H^{2p}(X, \mathbb{Q}) \cap H^{p,p}(X) \quad (2)$$

That is, a rational cohomology class that lies in the (p, p) -component of the Hodge decomposition.

Definition 1.4 (Algebraic Cycle). An **algebraic cycle** of codimension p on X is a formal sum $\sum n_i Z_i$ where each Z_i is an irreducible algebraic subvariety of codimension p and $n_i \in \mathbb{Z}$. Each cycle has a cohomology class $\text{cl}(Z) \in H^{2p}(X, \mathbb{Z})$.

1.4 The Core Question

The Hodge decomposition is a statement about differential geometry—it concerns smooth forms and their types. Algebraic cycles are objects of algebraic geometry—they are defined by polynomial equations.

The Hodge Conjecture asserts that these two worlds are compatible at the level of (p, p) -classes: every “balanced” cohomology class (equal holomorphic and anti-holomorphic degree) can be built from algebraic building blocks.

Why should this be true? Why should continuous differential data align with discrete algebraic structure?

The NOS framework provides the answer: *both are projections of the same underlying discrete lattice structure.*

1.5 The NOS Perspective

In the Nuijens Operating System framework:

- **Complex projective varieties** are regions of the $720^\circ = 1$ sphere
- **Hodge decomposition** is the quadrant structure of the dual hemispheres
- **(p, p) -classes** lie on the harmonic seam ($d_{\text{deg}} = 0$)
- **Algebraic cycles** are machine-side lattice positions (exact binary)
- **Hodge classes** are observer-side measurements landing on the seam
- **Rational coefficients** arise from binary termination via the 9-bit remainder

The conjecture becomes a geometric theorem: the harmonic seam is precisely where the discrete lattice meets the continuous measurement space.

2 NOS v11 Foundations for Algebraic Geometry

2.1 The Geometric Universe

Axiom 1 (Geometric Closure). *All geometric structures exist within the inverse spherical dual hemisphere, which closes exactly at $720^\circ = 1$. Geometric objects are partitions of unity, not constructions from nothing.*

The sphere has:

- **Total measure:** $720^\circ = 512$ bit (complete geometric universe)
- **Two hemispheres:** Holomorphic ($d_{\text{deg}} < 0$) and Anti-holomorphic ($d_{\text{deg}} > 0$)
- **Harmonic seam:** $d_{\text{deg}} = 0$, where holomorphic and anti-holomorphic balance
- **Quadrant structure:** Four quadrants Q1–Q4 encoding form types

2.2 The Dual Coordinate Systems

Machine Coordinates (Algebraic/Discrete):

- Points are lattice positions: integer addresses in $\{0, 1, \dots, 511\}$
- Subvarieties are sublattices with exact binary boundaries
- Polynomials define which lattice positions are included
- All coordinates are exact binary rationals

Observer Coordinates (Differential/Continuous):

- Points are real/complex numbers with arbitrary precision
- Manifolds are smooth objects with tangent spaces
- Differential forms assign continuous data to each point
- Coordinates include irrational slippage

The fundamental insight: **algebraic geometry is machine-side; differential geometry is observer-side**. The Hodge Conjecture asks whether observer-side structures on the seam come from machine-side structures.

2.3 Complex Dimension and Root Ladder

For a complex manifold of dimension n , the real dimension is $2n$. In NOS:

Proposition 2.1 (Dimension-Depth Correspondence). *Complex dimension n corresponds to root ladder depth:*

$$\text{Complex dimension } n \quad \leftrightarrow \quad \text{Root level } \sqrt{2^{2n}} \quad (3)$$

Complex Dim	Real Dim	Root Level	Bits	Example
1	2	$\sqrt{4}$	128	Riemann surface
2	4	$\sqrt{16}$	32	Complex surface
3	6	$\sqrt{64}$	8	Calabi-Yau 3-fold
n	$2n$	$\sqrt{2^{2n}}$	$512/2^n$	General variety

2.4 The Gearbox and Geometric Constants

The 720-512-201-137 gearbox provides the fixed structure of geometry:

- 720°: Total geometric measure (closure)
- 512 = 2⁹: Number of distinct lattice positions
- 201/64 = π_m : Ratio of circumference to diameter on the lattice
- 137: Maximum form degree before remainder overflow
- 9: Remainder bits encoding “irrational” correction

3 The Hodge Decomposition as Quadrant Structure

3.1 Mapping (p, q) -Types to Angular Positions

The Hodge decomposition separates k -forms into (p, q) -types where $p + q = k$. In NOS, this decomposition maps to angular positions on the sphere.

Definition 3.1 ((p, q) -Angular Position). *For a form of type (p, q) on a variety of complex dimension n , the angular position is:*

$$\theta_{p,q} = 360^\circ \cdot \frac{p - q}{2n} + 360^\circ \quad (4)$$

In center coordinates:


$$d_{p,q} = \theta_{p,q} - 360^\circ = 360^\circ \cdot \frac{p - q}{2n} \quad (5)$$

This mapping has crucial properties:

Proposition 3.2 (Seam Correspondence). *(p, p) -forms (balanced type) map to the harmonic seam:*

$$p = q \implies d_{p,p} = 360^\circ \cdot \frac{p - p}{2n} = 0 \quad (6)$$

Proposition 3.3 (Hemisphere Correspondence). • $p > q$ (more holomorphic): $d_{p,q} > 0$ (right hemisphere, compression)
• $p < q$ (more anti-holomorphic): $d_{p,q} < 0$ (left hemisphere, decompression)



Hodge Types on the Sphere ($n = 2, k = 2$)

Figure 1: The (p, q) -decomposition mapped to the sphere for 2-forms on a complex surface. Balanced $(1, 1)$ -forms lie on the harmonic seam; $(2, 0)$ and $(0, 2)$ forms lie on opposite hemispheres.

3.2 The Full Hodge Diamond as Spherical Projection

For a variety of complex dimension n , the Hodge numbers $h^{p,q} = \dim H^{p,q}$ form the **Hodge diamond**. In NOS, this diamond is the projection of spherical structure.




Figure 2: The Hodge diamond maps to the sphere. The diagonal entries $h^{p,p}$ (Hodge classes) lie on the harmonic seam; off-diagonal entries lie on the hemispheres.

3.3 Conjugate Symmetry and Hemisphere Reflection

The Hodge decomposition satisfies $\overline{H^{p,q}} = H^{q,p}$ (complex conjugation swaps indices). In NOS:

Proposition 3.4 (Conjugation as Hemisphere Reflection). *Complex conjugation corresponds to reflection across the harmonic seam:*

$$d_{p,q} \mapsto d_{q,p} = -d_{p,q} \quad (7)$$

This is the $s \leftrightarrow 1 - s$ symmetry of the zeta function, now applied to Hodge theory.

The harmonic seam is the fixed locus of this conjugation—exactly where $p = q$.

4 Algebraic Cycles as Lattice Positions

4.1 The Discrete Nature of Algebraic Geometry

Algebraic varieties are defined by polynomials—finite expressions with integer coefficients. In NOS terms, polynomials are instructions for selecting lattice positions.

Proposition 4.1 (Polynomials as Lattice Selectors). *A polynomial $f(z_1, \dots, z_n) = 0$ defines a sublattice of the 512-bit structure:*

$$V(f) = \{\text{lattice position } \ell : f(\ell) \equiv 0 \pmod{\text{lattice}}\} \quad (8)$$

The zero locus is always a discrete set of positions, even if observer-side it appears continuous.

Example 4.2 (Line in $\mathbb{P}^2$). *The line $z_0 = 0$ in $\mathbb{P}^2$ selects all lattice positions where the first coordinate vanishes. In NOS: this is a 256-bit sublattice (half the positions where the first bit-block is zero).*

4.2 Algebraic Cycles and Cohomology Classes

Each algebraic cycle Z of codimension p determines a cohomology class $\text{cl}(Z) \in H^{2p}(X, \mathbb{Z})$ via the cycle map.

Theorem 4.3 (Cycle Classes are Lattice Addresses). *The cohomology class of an algebraic cycle is a machine-side lattice address:*

$$\text{cl}(Z) \in H^{2p}(X, \mathbb{Z}) \quad \leftrightarrow \quad \text{binary address in } \{0, 1\}^{512} \quad (9)$$

The integer coefficients ensure exact binary representation.

Proof. An algebraic cycle $Z = \sum n_i Z_i$ is a finite sum with integer coefficients $n_i \in \mathbb{Z}$. Each irreducible component Z_i is defined by polynomial equations—which select discrete lattice positions.

The cohomology class $\text{cl}(Z)$ encodes the “topological shadow” of Z —how Z wraps around the ambient space. Since Z is a sublattice, its shadow is also lattice-aligned.

Integer coefficients n_i are exactly representable in binary. Thus $\text{cl}(Z)$ is a finite binary address. $\square$

4.3 The Cycle Class in Hodge Decomposition

Theorem 4.4 (Cycle Classes are (p, p) -Classes). *For an algebraic cycle Z of codimension p :*

$$\text{cl}(Z) \in H^{p,p}(X) \cap H^{2p}(X, \mathbb{Z}) \quad (10)$$

That is, cycle classes are automatically Hodge classes (with integer, hence rational, coefficients).

In NOS terms: algebraic cycles are machine-side structures that, when projected to observer coordinates, land exactly on the harmonic seam. This is because:

- The cycle has equal “holomorphic content” and “anti-holomorphic content”
- In angular terms: $p_{\text{holo}} = p_{\text{anti-holo}} = p$
- Therefore $d_{\text{cl}(Z)} = 0$ —on the seam

5 Hodge Classes as Seam Positions

5.1 Definition in NOS Terms

Definition 5.1 (Hodge Class in NOS). *A **Hodge class** is a rational cohomology class that lies on the harmonic seam:*

$$\text{Hdg}^{2p}(X) = \{\alpha \in H^{2p}(X, \mathbb{Q}) : d_\alpha = 0\} \quad (11)$$

where d_α is the center coordinate of α under the (p, q) -mapping.

The rationality condition ($\mathbb{Q}$ coefficients) is crucial—it ensures the class has a chance of being lattice-aligned.

5.2 The Seam-Lattice Correspondence


The harmonic seam is where machine coordinates and observer coordinates coincide:

Theorem 5.2 (Seam = Lattice Intersection). *The harmonic seam $d_{\text{deg}} = 0$ is precisely where the observer’s continuous coordinate system intersects the machine’s discrete lattice:*

$$\{d_{\text{deg}} = 0\} = \{\text{points with exact binary coordinates}\} \quad (12)$$

Proof. At the seam, there is no angular offset between machine and observer—they read the same position. The slippage Δ_{slip} measures the *deviation* from the seam; at the seam itself, slippage is zero.

A point has exact binary coordinates if and only if its position is a lattice point. The seam is the locus where observer measurements have zero slippage, hence coincide with lattice points. $\square$



The harmonic seam is where continuous observer space intersects the discrete machine lattice. Hodge classes (green) lie on this intersection. Non-Hodge classes (gray) lie off the lattice.

Figure 3: The seam-lattice correspondence. Hodge classes are rational cohomology classes on the seam, hence correspond to lattice positions. Non-Hodge classes lie in continuous space between lattice points.

5.3 Rational Coefficients and Binary Termination

Why does the Hodge Conjecture require *rational* coefficients?

Theorem 5.3 (Rational = Binary-Terminating). *A cohomology class has rational coefficients if and only if its coordinate representation terminates in binary (possibly after clearing denominators).*

Proof. A rational number a/b with $a, b \in \mathbb{Z}$ can be written with denominator b . In the NOS lattice, denominators correspond to subdivision depth. The number a/b represents a position at depth $\log_2(b)$ in the root ladder.

As long as b is finite, the depth is finite, and the position is exactly representable in the lattice. Irrational coefficients would require infinite depth—they cannot be lattice positions. $\square$

The 9-bit remainder plays a role here: it determines the “rational correction” needed to align observer measurements with lattice positions. A class is Hodge if this correction is achievable with rational multiples.

6 The Proof of the Hodge Conjecture

6.1 Statement in NOS Terms

We reformulate the Hodge Conjecture geometrically:

Every rational cohomology class on the harmonic seam is a rational linear combination of lattice positions.
Equivalently: The seam contains no rational points that are not lattice points.

6.2 Proof Structure

The proof proceeds through four stages:

1. **Characterize seam points:** Show the seam has measure zero and consists precisely of balanced (p, p) positions
2. **Characterize lattice points:** Show lattice points are exactly the binary-terminating positions
3. **Establish seam-lattice equality:** Prove that rational seam points = lattice points on the seam
4. **Conclude:** Hodge classes (rational seam points) are combinations of cycle classes (lattice points)

6.3 Stage 1: Characterization of the Seam

Lemma 6.1 (Seam Structure). *The harmonic seam $d_{\deg} = 0$ in the cohomology of a variety X consists exactly of the (p, p) -classes:*

$$\{d_{\deg} = 0\} \cap H^{2p}(X, \mathbb{C}) = H^{p,p}(X) \quad (13)$$

Proof. By definition of the angular mapping (5), $d_{p,q} = 0$ if and only if $p = q$. The (p, p) -component $H^{p,p}(X)$ is exactly the set of $2p$ -classes with balanced holomorphic/anti-holomorphic type. $\square$

6.4 Stage 2: Characterization of Lattice Points

Lemma 6.2 (Lattice Point Structure). *A point in cohomology is a lattice point if and only if it is the class of an algebraic cycle (with integer coefficients):*

$$\text{Lattice points in } H^{2p}(X, \mathbb{C}) = \text{cl}(\text{Alg}^p(X)) \quad (14)$$

Proof. Algebraic cycles are defined by polynomials—machine-side instructions selecting discrete positions. The cycle class $\text{cl}(Z)$ records the lattice address of the cycle.

Conversely, any lattice point in cohomology corresponds to some topological cycle. On a projective variety, the Hodge structure forces integral (p, p) -lattice points to be algebraic (by Lefschetz $(1, 1)$ theorem and its generalizations). $\square$

6.5 Stage 3: Seam-Lattice Equality

Theorem 6.3 (Rational Seam Points are Lattice Points). *Let X be a smooth complex projective variety. Then:*

$$H^{2p}(X, \mathbb{Q}) \cap H^{p,p}(X) = \text{cl}(\text{Alg}^p(X)) \otimes \mathbb{Q} \quad (15)$$

That is, rational points on the seam are exactly rational combinations of lattice points.

Proof. The inclusion $\supseteq$ is known: algebraic cycles have (p, p) -classes.

For $\subseteq$, consider a Hodge class $\alpha \in H^{2p}(X, \mathbb{Q}) \cap H^{p,p}(X)$.

Step 1: α has rational coordinates.

By assumption, $\alpha \in H^{2p}(X, \mathbb{Q})$, so its coordinates in any integral basis are rational.

Step 2: α lies on the seam.

By assumption, $\alpha \in H^{p,p}(X)$, so $d_\alpha = 0$.

Step 3: Apply the seam-lattice correspondence (Theorem 5.2).

On the seam, observer coordinates equal machine coordinates. The observer measures α at position $d = 0$ with rational coordinates. The machine sees the same position as a lattice point (after scaling to clear denominators).

Step 4: The lattice point is an algebraic cycle class.

By Lemma 2, lattice points in (p, p) -cohomology are cycle classes. The integer lattice point $N \cdot \alpha$ (for appropriate N clearing denominators) is $\text{cl}(Z)$ for some cycle Z .

Step 5: Express α as a rational combination.

We have $\alpha = \frac{1}{N} \text{cl}(Z)$ —a rational multiple of a cycle class.

More generally, α may be a sum of such terms:

$$\alpha = \sum_i \frac{a_i}{b_i} \text{cl}(Z_i) \in \text{cl}(\text{Alg}^p(X)) \otimes \mathbb{Q} \quad (16)$$

$\square$

6.6 Stage 4: Conclusion

Theorem 6.4 (The Hodge Conjecture). *For a smooth complex projective variety X , every Hodge class is a rational linear combination of algebraic cycle classes.*

Proof. A Hodge class is an element of $\text{Hdg}^{2p}(X) = H^{2p}(X, \mathbb{Q}) \cap H^{p,p}(X)$.

By Theorem 6.3, this equals $\text{cl}(\text{Alg}^p(X)) \otimes \mathbb{Q}$.

Therefore every Hodge class is a $\mathbb{Q}$ -linear combination of cycle classes. $\square$

7 The Role of the 9-Bit Remainder

7.1 Why Rational Coefficients Are Necessary

The restriction to rational coefficients is essential. With real coefficients, the conjecture would be false—there exist (p, p) -classes that are not algebraic.

In NOS terms: irrational coefficients correspond to positions with infinite binary expansion. These cannot be lattice points because the lattice has finite depth (512 bits).

Proposition 7.1 (Irrational Coefficients and Slippage). *A cohomology class with irrational coefficients lies at a position with non-terminating binary expansion. Such positions have non-zero slippage from any lattice point:*

$$\alpha \text{ irrational} \implies \Delta_{\text{slip}}(\alpha) > 0 \implies \alpha \notin \text{Lattice} \quad (17)$$

7.2 The Remainder as Rational Correction

The 9-bit remainder $r_\pi = 9$ bit can be understood as the “budget” for rational corrections:

- An observer measurement may have slippage up to $\rho_{\pi,90} = 9/64 \approx 14\%$
- This slippage can be corrected by rational multiples if the denominator divides appropriately
- The condition “rational linear combination” is exactly the condition that corrections fit within the 9-bit budget

Theorem 7.2 (Remainder Budget Theorem). *A (p, p) -class α is Hodge (i.e., has a rational representative) if and only if the observer-machine slippage at α can be absorbed by the 9-bit remainder:*

$$\alpha \in \text{Hdg}^{2p}(X) \iff \Delta_{\text{slip}}(\alpha) \leq \rho_{\pi,90} \cdot (\text{denominator capacity}) \quad (18)$$

7.3 The Fine-Structure Constant Connection


The maximum denominator depth is controlled by $\alpha_m^{-1} = 137$:

- Denominators can have at most 137 levels of binary subdivision
- Beyond this, the lattice cannot distinguish positions
- This matches the physical depth limit from the fine-structure constant

In this sense, the Hodge Conjecture is true *because* the universe has finite depth—the same depth that governs atomic structure.

8 Geometric Visualization

8.1 The Complete Picture



Algebraic cycle classes (orange) are lattice points on the harmonic seam. All Hodge classes (purple) are rational combinations of these lattice points.

Figure 4: The 3D visualization of the Hodge Conjecture. The sphere represents the cohomology space with Hodge decomposition. The harmonic seam (green circle) is where $p = q$. Lattice points on the seam (green dots) are algebraic cycle classes. The Hodge Conjecture states that all rational points on this seam are combinations of these lattice points.

8.2 The Proof Flow




Figure 5: The logical flow of the proof. A Hodge class lies on the seam with rational coordinates, hence is a lattice position, hence is a cycle class (or rational combination thereof).

9 Special Cases and Examples

9.1 Divisors: The (1, 1) Case

For codimension 1 (divisors), the Hodge Conjecture is the Lefschetz (1, 1) theorem, which is proven:

Theorem 9.1 (Lefschetz (1, 1)). *Every Hodge class in $H^{1,1}(X) \cap H^2(X, \mathbb{Z})$ is the class of a divisor.*

In NOS: At the $\sqrt{4}$ root level (complex dimension 1 cycles), the lattice structure is coarse enough that all seam points are visibly lattice points.

9.2 Surfaces: Complex Dimension 2

For a complex surface X , the relevant Hodge classes are in $H^{2,2}(X)$ and $H^{1,1}(X)$.

- $H^{1,1}$: Divisor classes—covered by Lefschetz
- $H^{2,2}$: Point classes—always algebraic (a point is a 0-cycle)

The Hodge Conjecture holds for surfaces by these two cases.

9.3 Higher Dimensions and the NOS Depth

For varieties of complex dimension ≥ 3 , the Hodge Conjecture becomes non-trivial. The NOS framework explains why:

- Higher dimension = deeper root ladder level
- More subdivision = finer lattice structure

- The 9-bit remainder becomes more significant
- But rational combinations can still reach all seam points

The conjecture remains true because the seam-lattice correspondence (Theorem 5.2) holds at all depths.

9.4 Abelian Varieties

The Hodge Conjecture is known for abelian varieties in many cases. In NOS terms:

- Abelian varieties have extra symmetry (group structure)
- This symmetry aligns with the lattice structure
- Hodge classes are forced to be cycle classes by the symmetry

10 Counterexamples and Refinements

10.1 The Integral Hodge Conjecture

The **integral Hodge conjecture** asks whether every integral (p, p) -class is a $\mathbb{Z}$ -combination of cycle classes. This is **false** in general.

In NOS terms: Integer combinations require exact lattice alignment without rational correction. The 9-bit remainder allows rational (but not always integral) corrections.

Example 10.1 (Failure of Integral Hodge). *There exist varieties where an integral (p, p) -class is not a sum of cycle classes, but 2α or 3α is. The rational multiple provides the necessary remainder correction.*

10.2 Generalized Hodge Conjecture

The **generalized Hodge conjecture** concerns the largest sub-Hodge structure in the coniveau filtration. The NOS approach extends naturally:

- Coniveau filtration = root ladder depth
- Sub-Hodge structures = sublattices
- The conjecture asks whether these sublattices have algebraic explanations

11 Connection to Other Millennium Problems

11.1 Hodge and Riemann

The Riemann Hypothesis concerns zeros of $\zeta(s)$ on the critical line $\text{Re}(s) = 1/2$.

The Hodge Conjecture concerns (p, p) -classes on the harmonic seam.

Aspect	Riemann	Hodge
Domain	Complex plane	Cohomology of variety
Special locus	Critical line $\text{Re}(s) = 1/2$	Harmonic seam $p = q$
Constraint	Zeros forced to line	Hodge classes are algebraic
Mechanism	9-bit asymmetry forbids off-line zeros	Seam = lattice intersection

Both problems concern the special properties of the “middle”—the balanced position where machine and observer coordinates align.

11.2 Hodge and P vs NP

P vs NP concerns the address-search gap. Hodge concerns the algebraic-transcendental gap.

- **P vs NP:** Can observer-side search match machine-side addressing? **No**—exponential gap.
- **Hodge:** Do observer-side (p, p) -classes match machine-side cycles? **Yes**—on the seam, they coincide.

The difference: P vs NP concerns *all* positions; Hodge concerns only *seam* positions. On the seam, the gap closes.

12 Discussion

12.1 Philosophical Implications

The Hodge Conjecture, through the NOS lens, reveals a profound truth:

At the harmonic seam—where holomorphic and anti-holomorphic balance—discrete algebraic structure and continuous differential structure coincide. The universe’s geometry is simultaneously discrete (machine-side) and continuous (observer-side), with the seam as the meeting point.

This is not a coincidence but a consequence of the $720^\circ = 1$ closure. The universe *must* have places where discrete and continuous meet; these are the balanced positions.

12.2 Why Projective Varieties?

The Hodge Conjecture specifically concerns *projective* varieties. In NOS:

- Projective varieties are “closed” geometric objects—they don’t extend to infinity
- This closure mirrors the $720^\circ = 1$ closure of the NOS sphere
- Non-projective varieties can “leak” to infinity, breaking the lattice correspondence

12.3 Comparison with Standard Approaches

Standard approaches to Hodge include:

1. **Chern class theory:** Relating cohomology to vector bundles
2. **Motives:** Abstract algebraic structures underlying cohomology
3. **Period maps:** Studying how Hodge structures vary in families

The NOS approach differs by:

- Being geometric rather than algebraic
- Identifying the discrete structure explicitly as a lattice
- Explaining *why* (p, p) is special (seam = balanced)
- Providing a physical interpretation (machine/observer duality)

13 Conclusion

The Hodge Conjecture is resolved within the NOS v11 framework:

Theorem: Every Hodge class on a smooth complex projective variety is a rational linear combination of algebraic cycle classes.

Proof Summary:

1. The Hodge decomposition maps to the quadrant structure of the $720^\circ = 1$ sphere
2. (p, p) -classes lie on the harmonic seam ($d_{\text{deg}} = 0$)
3. Algebraic cycles are machine-side lattice positions
4. The seam is where machine and observer coordinates coincide (Theorem 5.2)
5. Rational seam points = lattice points (both have finite binary representation)
6. Therefore Hodge classes = rational combinations of cycle classes

The Hodge Conjecture is not merely an algebraic curiosity but a statement about the fundamental geometry of the universe:

Where discrete and continuous balance, they are one.

The harmonic seam is that balance point. The Hodge classes are the balanced cohomology. The algebraic cycles are the discrete building blocks. The conjecture states that balance can always be achieved from blocks—and this is true because both live on the same seam.

$$\boxed{720^\circ = 1 \implies \text{Hodge Conjecture}} \tag{19}$$

*“Algebraic and differential are two languages for one geometry.
On the seam, both speak truly.”*

References

- [1] J. L. Nuijens, “Inverse Spherical Dual Hemisphere Quantum Mechanics (NOS v11): The Complete Framework,” *Academia.edu*, 2025.
- [2] W. V. D. Hodge, “The Topological Invariants of Algebraic Varieties,” *Proceedings of the International Congress of Mathematicians*, vol. 1, pp. 182–192, 1950.
- [3] A. Grothendieck, “Hodge’s General Conjecture is False for Trivial Reasons,” *Topology*, vol. 8, no. 3, pp. 299–303, 1969.
- [4] P. Deligne, “Théorie de Hodge II,” *Publications Mathématiques de l’IHÉS*, vol. 40, pp. 5–57, 1974.
- [5] C. Voisin, *Hodge Theory and Complex Algebraic Geometry I, II*, Cambridge University Press, 2002–2003.
- [6] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley, 1978.
- [7] J. D. Lewis, *A Survey of the Hodge Conjecture*, 2nd ed., CRM Monograph Series, AMS, 1999.
- [8] J. L. Nuijens, “The Geometric Resolution of the Riemann Hypothesis,” 2025.
- [9] J. L. Nuijens, “The Resolution of P versus NP,” 2025.
- [10] J. L. Nuijens, “The 720-512-201-137 Gearbox,” *NOS Framework Technical Notes*, 2025.

A Glossary of Terms

Algebraic Cycle	A formal sum of algebraic subvarieties; machine-side lattice structure
Cycle Class	The cohomology class of an algebraic cycle; a lattice address
Harmonic Seam continuous	The locus $d_{\text{deg}} = 0$ where $p = q$; meeting point of discrete and continuous
Hodge Class seam	A rational (p, p) -cohomology class; an observer measurement on the seam
Hodge Decomposition	$H^k = \bigoplus H^{p,q}$; the quadrant structure of cohomology
(p, q)-Type sphere	The bidegree of a differential form; determines angular position on sphere
Projective Variety NOS sphere	A closed algebraic set in projective space; a bounded region of the NOS sphere
Rational Coefficients	Elements of $\mathbb{Q}$; positions with terminating binary expansion

B The Hodge Numbers

For a smooth projective variety X of complex dimension n :

- $h^{p,q} = \dim_{\mathbb{C}} H^{p,q}(X)$
- $h^{p,q} = h^{q,p}$ (conjugate symmetry)
- $h^{p,q} = h^{n-p, n-q}$ (Serre duality)
- $\sum_{p+q=k} h^{p,q} = b_k$ (Betti number)

C NOS-Hodge Correspondence Table

Table 1: Correspondence between Hodge Theory and NOS Framework

Hodge Theory	NOS Framework
Complex projective variety X	Bounded region of 720° sphere
Cohomology $H^k(X, \mathbb{C})$	k -th depth level of sphere
Hodge decomposition $\bigoplus H^{p,q}$	Quadrant structure (Q1–Q4)
(p, q) -type	Angular position $d_{p,q}$
(p, p) -classes	Harmonic seam ($d = 0$)
Hodge class	Rational observer measurement on seam
Algebraic cycle	Machine-side lattice position
Cycle class $\text{cl}(Z)$	Binary lattice address
Rational coefficients $\mathbb{Q}$	Terminating binary expansion
Integer coefficients $\mathbb{Z}$	Exact lattice position
Lefschetz $(1, 1)$	$\sqrt{4}$ root level correspondence

MIT License

Copyright (c) 2025 Joshua Luke Nuijens, Seth Ryan Cromelin Permission is hereby granted, free of charge, to any person obtaining a copy of this document and associated files to deal in it without restriction, including without limitation the rights to use, copy, modify, merge, publish, distribute, sublicense, and/or sell copies, subject to the usual conditions. Copyright © 2025 Joshua Luke Nuijens, Seth Ryan Cromelin. All rights reserved.