


How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 50% regional coverage per fungal protein; direction margin 0.0 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 50% regional coverage per fungal protein; direction margin 0.01 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 50% regional coverage per fungal protein; direction margin 0.1 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 70% regional coverage per fungal protein; direction margin 0.0 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 70% regional coverage per fungal protein; direction margin 0.01 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 70% regional coverage per fungal protein; direction margin 0.1 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 90% regional coverage per fungal protein; direction margin 0.0 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 90% regional coverage per fungal protein; direction margin 0.01 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 30 common residues and 90% regional coverage per fungal protein; direction margin 0.1 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 50% regional coverage per fungal protein; direction margin 0.0 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 50% regional coverage per fungal protein; direction margin 0.01 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 50% regional coverage per fungal protein; direction margin 0.1 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 70% regional coverage per fungal protein; direction margin 0.0 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 70% regional coverage per fungal protein; direction margin 0.01 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 70% regional coverage per fungal protein; direction margin 0.1 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 90% regional coverage per fungal protein; direction margin 0.0 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 90% regional coverage per fungal protein; direction margin 0.01 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.


Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.

How experimental reference choice affects duplicate structural contrasts

Minimum 50 common residues and 90% regional coverage per fungal protein; direction margin 0.1 Å

All two-boundary × full/pLDDT ≥70 variants must qualify. Labels: qualified units / entities / dependence components.



Directions compare A-minus-B RMSD using fungal versus experimental references on the same sequence-anchored residue maps.

Units share chains, sequences and publications; proportions are descriptive, not independent replication, significance or experimental validation.

Selected 13 cases; zero coverage is not evidence of no structural difference. Domain orientation and prediction uncertainty remain relevant.