

Example Data Preparation Pipeline for Training MedGuideX



Clinical Practice Guideline

ADA recommends the use of any of the following four criteria for diagnosing diabetes:
1) glycated hemoglobin (A1C) value of 6.5% or higher...



Executable Function

```
1 def diagnose_diabetes(A1C, random_glucose, classic_symptoms, hyperglycemic_crisis, ...):
2     results = []
3     if A1C >= 6.5:
4         results.append("diagnose_diabetes_by_A1C")
5         # other steps...
6     if classic_symptoms or hyperglycemic_crisis and random_glucose >= 200:
7         results.append("diagnose_diabetes_by_random_glucose")
8     if not results:
9         results.append("diabetes_not_diagnosed")
10    return results
```



Factual QA

Question: A 52-year-old woman presents for evaluation of abnormal glucose testing ... Laboratory testing shows a hemoglobin A1c of 7.9%, ... According to the guideline, what would be recommended regarding the diagnosis of diabetes mellitus?

Answer: diagnose diabetes



Counterfactual QA

Question: A 52-year-old patient has a random plasma glucose of 156 mg/dL The HbA1c is 5.7%. The diagnosis is: `diabetes\_not\_diagnosed`. What if the HbA1c were 6.8% instead, while all other patient and program facts remained fixed?

Answer: Diagnose diabetes based on $A1C \geq 6.5\%$