

HHS Public Access

Author manuscript

J Aging Health. Author manuscript; available in PMC 2016 September 14.

Published in final edited form as:

J Aging Health. 2013 April; 25(3): 459–477. doi:10.1177/0898264312474039.

Depressive Symptomatology, Exercise Adherence and Fitness are Associated with Reduced Cognitive Performance in Heart Failure

Michael L. Alosco, B.A.¹, Mary Beth Spitznagel, Ph.D.^{1,2}, Manfred van Dulmen, Ph.D.¹, Naftali Raz, Ph.D.³, Ronald Cohen, Ph.D.⁴, Lawrence H. Sweet, Ph.D.⁵, Lisa H. Colbert, Ph.D.⁶, Richard Josephson, M.S., M.D.^{7,8,9}, Joel Hughes, Ph.D.^{1,2}, Jim Rosneck, M.S.², and John Gunstad, Ph.D.^{1,2}

¹Department of Psychology, Kent State University, Kent, OH, USA

²Department of Psychiatry, Summa Health System, Akron City Hospital, Akron, OH, USA

³Institute of Gerontology, Wayne State University, Detroit, MI, USA

⁴Departments of Neurology Psychiatry and the Institute on Aging, Center for Cognitive Aging and Memory, University of Florida

⁵Department of Psychology, University of Georgia, Atlanta, USA

⁶Department of Kinesiology, University of Wisconsin, Madison, WI, USA

⁷University Hospitals Case Medical Center and Department of Medicine, Cleveland, OH, USA

⁸Harrington Heart & Vascular Institute, Cleveland, OH, USA

⁹Case Western Reserve University School of Medicine, Cleveland, OH, USA

Abstract

Objectives—Depression is common in heart failure (HF) and associated with reduced cognitive function. The current study used Structrual Equation Modeling to examine whether depression adversely impacts cognitive function in HF through its adverse affects on exercise adherence and cardiovascular fitness.

Methods—158 HF patients completed neuropsychological testing, physical fitness test, Beck Depression Inventory-II (BDI-II), and measures assessing exercise adherence, and physical exertion.

Results—The model demonstrated excellent model fit and increased scores on the BDI-II negatively affected exercise adherence and cardiovascular fitness. There was a strong inverse association between cardiovascular fitness and cognitive function. Sobel test showed a significant indirect pathway between the BDI-II and cognitive function through cardiovascular fitness.

Discussion—This study suggests depression in HF may adversely impact cognitive function through reduced cardiovascular fitness. Prospective studies are needed to determine whether

treatment of depression can lead to better lifestyle behaviors and ultimately improve neurocognitive outcomes in HF.

Keywords

Cognitive function; cardiovascular fitness; depression; exercise adherence; heart failure

1. Introduction

Heart failure (HF) affects nearly 6 million Americans and is a major contributor to hospitalizations, elevated rates of mortality, and economic burden (American Heart Association 2010; Roger et al., 2010; Jencks, Williams & Coleman, 2009). HF is associated with poor quality of life and reduced functional independence (Norberg, Bowman, & Lofgren, 2008; Bennet, et al., 2003), and vast evidence indicates cognitive impairment leads to poor outcomes in this population (Alosco, et al., 2012; Zuccala, et al., 2003; Reigel & Weaver, 2009; Kindermann, et al., 2010). Relative to healthy controls, HF patients are at a 4-fold increased risk for cognitive impairment, including deficits in attention, executive function, and memory (Pressler, et al., 2010). The exact mechanisms of cognitive impairment in this population are unclear, though they likely involve cerebral hypoperfusion secondary to cardiac dysfunction combined with the pathological effects of medical comorbidity (i.e., hypertension, diabetes) (Suave, Lewis, Blankenbiller, Rickabaugh, & Pressler, 2009; Alosco, et al., 2012; Jefferson, et al., 2007; Vogels, et al., 2007; Zuccala, et al., 2005).

Depression has also recently been shown to be an important contributor to cognitive impairment in this population. Depression is found in nearly 30% of HF patients (Diez-Quevdeo, et al., 2012) and associated with elevated mortality risk, repeated hospitalizations, and adverse cardiac events (Rutledge, Reis, Linke, Greenberg, & Mill, 2006; Kato, et al., 2012). Recent work in this population links increased depressive symptomatology with deficits in attention, executive function, psychomotor speed, and language (Garcia, et al., 2011). Pathophysiological changes (i.e., reduced cerebral blood flow) have also been noted among HF patients with depression relative to their non-depressed counterparts (Alves, et al., 2006).

The mechanisms underlying the association between depression and cognitive impairment in persons with HF may involve reduced cardiovascular fitness. Past work has shown that depression contributes to increased dropout rates of exercise training cardiac rehabilitation programs and reduced physical activity and cardiovascular fitness in people with HF and other cardiovascular diseases (Casey, Hughes, Waechter, Josephson, & Rosneck, 2008; McGrady, McGinnis, Badenhop, Bentle, & Rajput, 2009; Spaderna, et al., 2010; Lysy, Da Costa, & Dasqupta, 2008; Gottlieb, et al., 2009). The links between depression and vascular risk have important consequences, as recent work links reduced cardiovascular fitness to poorer cognitive function among older adults with HF (Alosco, et al., 2011; Steinberg, et al., 2011).

Although past work has examined independent effects of depression, exercise adherence, and cardiovascular fitness on cognitive function in HF, no study to date has explored how

these factors may interact in a framework of a hypothesis driven multivariate model. Based on the reviewed findings, we hypothesized that reduced cardiovascular fitness stemming from exercise non-adherence would mediate the relationship between depression and cognitive dysfunction in HF. Our hypothetical relationships among the variables of interest were formalized in a Structural Equations Model (SEM) presented in Figure 1.

2. Methods

2.1 Participants

The sample consisted of 158 consecutively enrolled persons with HF from a large-scale project examining neurocognitive function in older adults with HF. All participants were between the ages of 50-85 years of age, English-speaking, and had an established diagnosis of HF at New York Heart Association (NYHA) class II or III at the time of enrollment. Exclusion criteria included history of significant neurological disorder (e.g. dementia, stroke, multiple sclerosis), head injury with more than 10 minutes loss of consciousness, severe psychiatric disorder, substance abuse and/or dependence, and renal failure. Participants averaged 68.96±9.13 years of age, were 36.7% female, and 82.3% Caucasian. Medical record review indicated the current sample demonstrated an average left ventricular ejection fraction (LVEF) of 40.70 (SD = 14.39). See Table 1 for sample demographic and clinical characteristics.

2.2 Measures

- **2.2.1. Depressive Symptomatology**—The Beck Depression Inventory-II (BDI-II) was administered to assess self-reported depressive symptomatology. The BDI-II is a commonly used checklist of depressive symptoms that exhibits strong psychometric properties in persons with medical conditions (i.e., test-re-test reliability of r= .93 to r= .96, and an internal consistency of r= .54 to r= .74) (Amau, Meagher, Norris, & Bramson, 2001; Beck, Steer, & Brown, 1996). BDI-II scores range from 0-63 with increased score indicative of increased symptomatology. The BDI-II score was used as a single manifest predictor in the proposed model.
- **2.2.2. Exercise Adherence**—The Heart Failure Compliance Questionnaire (Evangelista, Berg, & Dracup, 2001) was used to assess treatment adherence in the current sample. This questionnaire asks participants to rate their adherence to different health behaviors (i.e., maintaining doctor appointments, medication management, substance abstinence, following dietary and exercise recommendations). The current study examined participants' responses to the item of the questionnaire assessing exercise adherence. Participants rated their adherence to exercise behaviors based on a scale of 0 (none of the time) to 4 (all of the time). Scores were converted to a 0 to 100 scale with a score of 75% or greater reflecting adherent behaviors (i.e., participants responded adhering to exercise recommendations most or all of the time) (Evangelista, et al., 2001). This exercise adherence item served as a single manifest mediator variable in the proposed model.
- **2.2.3. Cardiovascular Fitness**—Cardiovascular fitness served as a latent construct in the proposed model and was assessed using the 2-Minute Step Test (2MST) and the Duke

Activity Status Index (DASI). The 2MST is an assessment of cardiovascular endurance (Jones & Rikli, 2002). The 2MST has been suggested to be an alternative to the 6-minute walk test, which has been linked with functional work capacity, maximal oxygen uptake, and poor prognosis in patients with HF (Nixon, Joswiak, & Fricker, 1996; Bittner, et al., 1993). Past work has also shown that the 2MST is linked with poorer cognitive function among older adults with heart failure (Alosco, et al., 2011). The 2MST requires the patient to march in place for 2 minutes. The patient is asked to bring each knee up to a marked target set on the wall at the individual's own midpoint between the kneecap and crest of the iliac. The number of times the right knee met the marked target was counted. Increased step count within the 2-minutes was reflective of greater cardiovascular fitness. Average step count for females between the ages of 50-85 ranges from 71-115, while males for this age group range from 60-107 (Jones & Rikli, 2002).

Maximal physical exertion was assessed using the Duke Activity Status Index (DASI), a check-list, in which participants were asked to identify activities they were able to complete, ranging from basic tasks (i.e. walking around the house) to vigorous exercise (i.e. basketball, skiing). The score was a sum of the weighted values for each checked activity. From the composite score, a proxy estimate of peak oxygen uptake in mL/min was computed to serve as an indicator of maximal physical exertion (Hlatky, et al., 1989).

2.2.4. Cognitive Function—A brief neuropsychological battery was administered to all participants to assess multiple domains of cognitive function, including attention, executive function, memory, and language. All tests demonstrate strong psychometric properties including excellent reliability and discriminant and construct validity. Neuropsychological tests administered and used as indicators of the latent construct cognitive function in the proposed model include:

- Trail Making Test A (Spreen & Strauss, 1991): Completion time of Trail
 Making Test A is a reliable and valid measure of attention and complex
 visual scanning and psychomotor speed (Spreen & Strauss, 1991).
 Participants are asked to connect a series of letters in sequential order as
 quickly as possible. Longer time of completion is reflective of worse
 performance.
- Trail Making Test B (Dikmen, Grant, & Tekmin, 2000): Trail Making Test B is a widely used measure of executive function and assesses the ability to shift and maintain cognitive set. Participants are asked to connect a series of numbers and letters in alternating sequential order as quickly as possible. Longer time of completion is indicative of worse performance.
- Frontal Assessment Battery (Dubois, Slachevsky, Litvan, & Pillon, 2000): The Frontal Assessment Battery is comprised of six subtests that assess aspects of executive function, including lexical fluency, inhibitory control, sensitivity to interference, higher-ordered motor programming, and abstract reasoning. Scores range from 0-18 with higher scores reflective of better performance.

California Verbal Learning Test Delayed Recall (Delis, Kramer, Kaplan, & Ober, 2000): The California Verbal Learning Test-II (CVLT-II) long delay free recall was used to assess memory functioning in the current sample. Individual are asked to learn a 16-item word list and recall these words after a delay period. This widely used test of verbal memory. Scores on the long delay free recall range from 0-16 and increased score is reflective of better performance.

Animals Fluency Test (Morris, et al., 1989): The animal fluency test was
used to assess language abilities. This test has individuals generate as
many animal names as possible within a given time limit. Greater number
of animals named reflects better performance.

2.2.5. Demographic Characteristics—Medical and demographic characteristics were ascertained through participant self-report and medical record review.

2.3 Procedures

The local Institutional Review Board (IRB) approved the study procedures and all participants provided written informed consent prior to study enrollment. During a single assessment, participants completed demographic and psychosocial self-report measures, including the BDI-II, DASI, and the Heart Failure Compliance Questionnaire. Participants were also administered a brief neuropsychological test battery to assess cognitive function. Finally, all participants completed the 2MST.

2.4 Statistical Analyses

To test the hypothesis driven model depicted in Figure 1, we used a structural equation modeling (SEM) approach. The model consists of two latent factors: cardiovascular fitness and cognitive function. The 2MST and a proxy measure of peak oxygen uptake served as indicators of cardiovascular fitness. The five neuropsychological measures that assess multiple aspects of cognition contributed to cognitive function. To minimize discrepancy among the scales and maintain directionality, all raw scores of the neuropsychological measures were converted to T-scores (a distribution with a mean of 50 and a standard deviation of 10) using existing normative data adjusting for age, and gender in the case of the California Verbal Learning Test. Additionally, a single indicator variable for each latent construct (i.e., cognitive function and cardiovascular fitness) was fixed at 1 to correct for scaling. The BDI-II and exercise adherence served as a manifest predictor and mediator variable in the current model, respectively. A measurement model was first conducted to examine model fit among the latent variables and their respective indicators. A structural model was then fitted to the data to examine the relationships among the BDI-II, exercise adherence, cardiovascular fitness, and cognitive function. To maintain model parsimony and preserve statistical power, covariates were not included in the model.

The SEM was conducted using EQS software using maximum likelihood approach. Goodness of fit was evaluated by the χ^2 value, comparative fit index (CFI) and the root mean-square error of approximation (RMSEA), recommended as most useful indices of

model fit (Thompson, 2000). A non-significant χ^2 , CFI .95, and an RMSEA .06 indicated good model fit (Hu & Bentler, 1999).

3. Results

Descriptive Statistics

Self-reported depressive symptomatology was prevalent, in this sample of HF patients, with a mean BDI-II score of 7.67 (SD = 7.57). Many of the HF participants reported high rates of exercise non-adherence. Specifically, the sample averaged 56.33 (SD = 33.97) on the exercise adherence composite and 50.6% of the sample reported being non-adherent (i.e., score < 75). Not surprisingly, when compared to normative data accounting for age and gender, both male and female HF patients also performed below the average range on the 2MST, suggesting poorer cardiovascular fitness. See Table 1 for a summary of descriptive statistics.

Consistent with clinical convention, a T-score cutoff of 35 (1.5 SD below the mean) was used to define impairment on neuropsychological measures assessing cognitive function. Many HF participants fell below this cutoff on several of the cognitive tests. Specifically, 26.6% exhibited impairments on the Frontal Assessment Battery, 18.4% on Trail-Making Test B, 10.1% on Trail-Making Test A, and 7.6% on the California Verbal Learning Test Delayed Recall. Impairment on the Animals Fluency Test was less common. See Table 1 for cognitive test T-score means and standard deviations.

Measurement Model

A measurement model was first evaluated to examine fit of the indicators of the latent constructs cardiovascular fitness and cognitive function. The latent factors cardiovascular fitness and cognitive function were allowed to covary. Animals and estimated peak oxygen uptake (as assessed by the DASI) were fixed at 1.0 for cognitive function and cardiovascular fitness, respectively. The measurement model demonstrated excellent fit: χ^2 (13, N= 158) = 13.29, p= .43, CFI= 1.00, RMSEA = .01 (90% CI = .00, .08). All neuropsychological measures significantly loaded on to cognitive function latent factor (β = .33 to .81, p< .05) and the 2MST significantly loaded on to cardiovascular fitness factor (β = .68, p< .05). In addition, cognitive function and cardiovascular fitness factors significantly covaried (β = .53, p< .05), with greater cardiovascular fitness associated with better cognitive function.

Structural Model

Refer to Table 2 for the observed covariance matrix among the variables analyzed using SEM. The model demonstrated excellent fit: χ^2 (23, N= 158) = 31.39, p= .11, CFI= .97: RMSEA = .05 (90% CI = .00, .09). Examination of structural pathways revealed that increased depressive symptomatology was associated with reduced exercise adherence and poorer cardiovascular fitness (p< .05). Decreased exercise adherence was also significantly associated with worse cardiovascular fitness (p< .05).

Finally, the pathways connecting the BDI-II and cardiovascular fitness with cognitive function were also significant with a stronger association noted between cardiovascular

fitness and cognitive function (p<.05). In each case, increased depressive symptomatology and poorer cardiovascular fitness were associated with decreased cognitive function. The direct pathway between exercise adherence and cognitive function was not significant (p>.05). To determine whether cardiovascular fitness demonstrated a statistically stronger relationship with cognitive function relative to the BDI-II, the model was evaluated again after constraining the pathways between the BDI-II and cardiovascular fitness with cognitive function to be equal. This model fit the data significantly worse than the original one (χ^2 (1, N=158) = 30.90, p<.001), suggesting that cardiovascular fitness exerts greater influence on cognitive function than depressive symptomatology. Sobel test also revealed that there was a significant indirect effect of the BDI-II on cognitive function through cardiovascular fitness (p<.05), though not through exercise adherence (p>.05). Cardiovascular fitness demonstrated significant indirect effects with all of the individual cognitive variables (p<.05 for all). See Figure 1 for standardized parameter estimates.

4. Discussion

Consistent with extant research, HF patients evidenced poor cognitive performance and high prevalence of self-reported depressive symptomatology. In HF patients, depression is associated with poor psychosocial outcomes (Rutledge, et al., 2006; Kato, et al., 2012) and may contribute to cognitive impairment (Garcia, et al., 2011). This study suggests this relationship may stem in part from poor adherence to exercise regimen and reduced cardiovascular fitness.

More than half of the current sample of HF patients reportedly did not follow exercise recommendations, and our findings suggest that depressive symptomatology may contribute to poor adherence. The negative impact of depression on exercise adherence is common among patients with other cardiac problems (Casey, et. al, 2008). There are several potential explanations for the adverse effects of depression on exercise adherence. First, depressed patients may have exaggerated perception of physical symptoms associated with HF (i.e., fatigue, shortness of breath) (Sullivan, LaCroix, & Russo, 1999; Falk, Patel, Swedberg, & Ekman, 2009). Heightened aversion of these symptoms that also accompany normal physical exertion may act as stronger barriers to exercise. Second, motivational factors (i.e., increase self-confidence, enjoyment of exercise) play an important role in exercise adherence among cardiac patients (Marzolini, Mertens, Oh, & Pyley, 2010; Farley, Wade, & Birchmore, 2003; Coghill & Cooper, 2009; Fernandez, Salamonson, Griffiths, Juergens, & Davidson, 2008; Perry & Bennett, 2006) and depression that reduces hedonic value of many activities likely impairs motivational capacities. Lastly, poor self-efficacy is common in depression (Maeda, Shen, Schwarz, Farrell, & Mallon, 2012) and this finding extends to HF patients, particularly regarding their functional ability (Skotzko, et al., 2000). Indeed, poor self-efficacy is a known contributor to treatment adherence (Maeda, et al., 2012) and poor exercise adherence in HF and other cardiovascular disease populations (Maeda, et al., 2012; Oka, DeMarco, & Haskell, 1999; Gardner, et al., 2003). Prospective studies should examine whether treatment of depression (i.e., psychotherapy, anti-depressant medications) improves adherence to exercise regimens in HF patients.

Despite the adverse impact of depression on exercise adherence, a strong relationship also emerged between depressive symptomatology and cardiovascular fitness. The exact reason for this relationship is not entirely clear. Past work among cardiac patients has shown depressive symptoms are associated with reduced exercise capacity (Ruo, Rumsfeld, Pipkin, & Whooley, 2004). It is possible that reduced cardiovascular fitness in depressed HF patients may be a manifestation of a longstanding sedentary lifestyle rather than current exercise adherence (as was assessed in the current study) (Roshanaei-Moghaddam, Katon, & Russo, 2009). In fact, compared to short-term involvement, long-term participation in physical activity is associated with greater and gradual improvements in cardiovascular fitness over time (Bulmenthal, et al., 1991). Depression is also associated with other poor health behaviors that adversely affect cardiovascular fitness, including smoking and obesity (Katon, et al., 2004; Katon W. J., 2003). Indeed, many HF patients are obese (Kapoor & Heidenreich, 2010) and engage in behaviors such as smoking and alcohol use (Kravitz, et al., 1993). Future studies should prospectively examine the association among depression, exercise habits, and cardiovascular fitness in HF patients, as such studies may provide greater insight into the current findings and identify possible intervention strategies.

Our findings indicate that poor cardiovascular fitness may explain the relationship between depression and reduced cognitive function in HF patients. Cardiovascular fitness is strongly influenced by exercise activity and reflective of increased ability of the heart to deliver oxygen throughout the body (American College of Sports Medicine, 2006; Netz, Dwolatzky, Zinker, Argov, & Agmon, 2011). Our findings, in agreement with current literature support the argument that physical fitness rather than current physical activity to be more closely related to cognitive function (Etnier, Sibley, Pomeroy, & Kao, 2003; Netz, Argov, & Inbar, Fitness's moderation of the facilitative effect of acute exercise on cognitive flexibility in older women, 2009). Indeed, cardiovascular fitness is associated with better cognitive test performance in patients with cardiovascular disease (Alosco, et al., 2011; Steinberg, et al., 2011; Gunstad, et al., 2005) and with greater structural and functional integrity of the brain among older adults (Colcombe, et al., 2003; Erickson, et al., 2009). A likely explanation for these findings may involve the beneficial effects of cardiovascular fitness on the vascular system, including endothelial and cardiac functioning (Corvera-Tindel, Doering, Woo, Khan, & Dracup, 2004; Chicco, 2008; Papathanasiou, Tsamis, Georgiadou, & Adamopoulos, 2008; Davison, et al., 2010; Moser, et al., 2004; Jefferson A. L., 2010). Cerebral perfusion also likely plays a critical role, as hypoperfusion is commonly theorized to underlie cognitive deficits in HF patients (Jefferson, et al., 2007; Jefferson, Poppas, Paul, & Cohen, 2007) and this deficiency may be alleviated through improved physical fitness (Stanek, et al., 2011; Rogers, Meyer, & Mortel, 1990). Depressed HF patients appear to be a greater risk for reduced cerebral perfusion (Alves, et al., 2006) and prospective studies should examine whether exercise interventions in HF patients with depression improves cognitive function.

The current study is limited in several ways. First, the present analyses consisted of cross-sectional data and longitudinal studies are needed to clarify the directionality of the relationship between depression and exercise performance and subsequent outcomes. Second, depressive symptomatology, exercise adherence, and estimated oxygen uptake were examined using self-report and future work should examine the relationships among these variables using objective measures (i.e., actigraphy, stress testing). Additionally, the current

study also did not examine potential improvements in cognitive function following increased cardiovascular fitness or underlying mechanisms involved. Future studies may benefit from use of functional magnetic resonance imaging (fMRI) to examine the pathophysiological effects of cardiovascular fitness among depressed HF patients. Consistent with this notion, depression has also been associated with increased obesity and diabetes (Ehlert, Gaab, & Heinrichs, 2001; Akil, et al., 1993), and elevations in inflammatory markers (i.e., C-reactive protein) (Panagiotakos, et al., 2004) and future studies should also explore these factors as possible mechanisms for reduced cognitive function as well as poor cardiovascular fitness.

In summary, the current study suggests that reduced cardiovascular fitness may partially explain decreased cognitive function among HF patients with depression. As a frequent morbidity in persons with HF, depression is a potentially important contributor to poor exercise adherence and reduced cardiovascular fitness in this population and thus may impede rehabilitation. Longitudinal studies are needed to elucidate underlying mechanisms and determine whether treatment of depression leads to better neurocognitive outcomes through modification of factors negatively impacting cardiovascular fitness (i.e., physical inactivity).

Acknowledgments

Support for this work included National Institutes of Health (NIH) grants DK075119 and HLO89311. Dr. Naftali Raz is also supported by National Institutes of Health (NIH) grant R37 AG011230. The authors have no competing interests to report.

References

- Alosco ML, Spitznagel MB, Cohen R, Sweet LH, Josephson R, Waechter D, Hughes J, Rosneck J, Gunstad J. Cognitive impairment is independently associated with reduced instrumental activities of daily living in persons with heart failure. Journal of Cardiovascular Nursing. 2012; 27:44–50. [PubMed: 21558863]
- Alosco ML, Spitznagel MB, Raz N, Cohen R, Sweet LH, Colbert LH, Gunstad J. The 2-Minute Step Test is independently associated with cognitive function in older adults with heart failure. Aging Clinical and Experimental Research. 2011
- Alosco ML, Spitznagel MB, van Dulmen M, Raz N, Cohen R, Sweet LH, Gunstad J. The Additive Effects of Type-2 Diabetes on Cognitive Function in Older Adults with Heart Failure. Cardiology Research and Practice. 2012; 2012 Article ID 2012-348054. doi:10.1155/2012/348054.
- Alves TC, Rays J, Fraquas RJ, Wajngarten M, Telles RM, Duran FL, Busatto GF. Association between major depressive symptoms in heart failure and impaired regional cerebral blood flow in the medial temporal region: a study using 99m Tc-HMPAO single photon emission computerized tomography (SPECT). Psychological Medicine. 2006; 36(5):597–608. [PubMed: 16515732]
- Amau RC, Meagher MW, Norris MP, Bramson R. Psychometric evaluation of the Beck Depression Inventory-II with primary care medical patients. Health Psychology. 2001; 20(2):112–119. [PubMed: 11315728]
- American College of Sports Medicine. American College of Sports Medicines Guidelines for Exercise Testing and Prescription. Lippincott Williams & Wilkins; Baltimore: 2006.
- American Heart Association. Heart Disease and Stroke Statistics 2010 Update. American Heart Association; Dallas, TX: 2010.
- Beck, AT.; Steer, RA.; Brown, GK. Beck Depression Inventory. The Psychological Corporation; San Antonio, TX: 1996.
- Bennet SJ, Oldridge NB, Eckert GJ, Embree JL, Browning S, Hou N, Murray M. Comparison of quality of life measures in heart failure. Nursing Research. 2003; 52:207–216. [PubMed: 12867777]

Bittner V, Weiner DH, Yusuf S, Rogers WJ, McIntyre KM, Bangdiwala SI, Bourassa MG. Prediction of Mortality and Morbidity With a 6-Minute Walk Test in Patients With Left Ventricular Dysfunction. SOLVD Investigators. The Journal of the American Medical Association. 1993; 270(14):1702–1707. [PubMed: 8411500]

- Bulmenthal JA, Emery CF, Madden DJ, Coleman RE, Riddle MW, Schniebolk S, Higginbotham MB. Effects of exercise training on cardiorespiratory function in men and women older than 60 years of age. The American Journal of Cardiology. 1991; 67(6):633–639. [PubMed: 2000798]
- Casey E, Hughes JW, Waechter D, Josephson R, Rosneck J. Depression predicts failure to complete phase-II cardiac rehabilitatio. Journal of Behavioral Medicine. 2008; 31(5):421–431. [PubMed: 18719990]
- Caulin-Glaster T, Maciejewski PK, Snow R, Mazure C. Depressive symptoms and sex affect completion rates and clinical outcomes in cardiac rehabilitation. Preventive Cardiology. 2007; 10(1):15–21. [PubMed: 17215628]
- Chicco AJ. Exercise training in prevention and rehabilitation: which training mode is best? Minerva Cardioangiologica. 2008; 56(5):557–570. [PubMed: 18813189]
- Coghill N, Cooper AR. Motivators and de-motivators for adherence to a program of sustained walking. Preventive Medicine. 2009; 49(1):24–27. [PubMed: 19426757]
- Colcombe SJ, Erickson KI, Raz N, Webb AG, Cohen NJ, McAuley E, Kramer AF. Aerobic fitness reduces brain tissue loss in aging humans. The Journals of Gerontology. 2003; 58(2):176–180. [PubMed: 12586857]
- Corvera-Tindel T, Doering LV, Woo MA, Khan S, Dracup K. Effects of a home walking exercise program on functional status and symptoms in heart failure. American Heart Journal. 2004; 147(2):339–346. [PubMed: 14760334]
- Davison K, Bircher S, Hill A, Coates AM, Howe PR, Buckley JD. Relationships between Obesity, Cardiorespiratory Fitness, and Cardiovascular Function. Journal of Obesity. 2010 Article ID 191263. doi:10.1155/2010/191253.
- Delis, D.; Kramer, J.; Kaplan, E.; Ober, B. California Verbal Learning Test-Second Edition: Adult Version. Manual. Psychological Corporation; San Antonio, TX: 2000.
- Diez-Quevdeo C, Lupon J, Gonzalez B, Urrutia A, Cano L, Cabanes R, Bayes-Genis A. Depression, antidepressants, and long-term mortality in heart failure [published online ahead of print April 14 2012]. International Journal of Cardiology. 2012 doi: 10.1093/eurjhf/hfs133.
- Dikmen SH, Grant I, Tekmin N. Test-retest reliability and practice effects of the Expanded Halstead-Reitan Neuropsychological Test Battery. Journal of the International Neuropsychological Society. 2000:346–356.
- Dubois B, Slachevsky A, Litvan I, Pillon B. The FAB: a Frontal Assessment Battery at bedside. Neurology. 2000; 55(11):1621–1626. [PubMed: 11113214]
- Ehlert U, Gaab J, Heinrichs M. Psychoneuroendocrinological contributions to the etiology of depression, posttraumatic stress disorder, and stress-related bodily disorders: the role of the hypothalamus-pituitary-adrenal axis. Biological Psychology. 2001; 57(1-3):141–152. [PubMed: 11454437]
- Erickson KI, Parkash RS, Voss MW, Chaddock L, Hu L, Morris KS, Kramer AF. Aerobic fitness is associated with hippocampal volume in elderly humans. Hippocampus. 2009:1030–1090. [PubMed: 19123237]
- Evangelista LS, Berg J, Dracup K. Relationship between psychosocial variables and compliance in patients with heart failure. Heart & Lung. 2001; 30(4):294–301. [PubMed: 11449216]
- Falk K, Patel H, Swedberg K, Ekman I. Fatigue in patients with chronic heart failure a burden associated with emotional and symptom distress. European Journal of Cardiovascular Nursing. 2009; 8(2):91–96. [PubMed: 18715830]
- Farley RL, Wade TD, Birchmore L. Factors influencing attendance at cardiac rehabilitation among coronary heart disease patients. European Journal of Cardiovascular Nursing. 2003; 2(3):205–212. [PubMed: 14622628]
- Fernandez RS, Salamonson Y, Griffiths R, Juergens C, Davidson P. Sociodemographic predictors and reasons for participation in an outpatient cardiac rehabilitation programme following percutaneous

- coronary intervention. International Journal of Nursing Practice. 2008; 14(3):237–242. [PubMed: 18460065]
- Garcia S, Spitznagel MB, Cohen R, Raz N, Sweet L, Colbert L, Gunsad J. Depression Is Associated with Cognitive Dysfunction in Older Adults with Heart Failure. Cardiovascular Psychiatry and Neurology. 2011; 2011 Article ID 368324. doi:10.1155/2011/368324.
- Gardner JK, McConnell TR, Klinger TA, Herman CP, Hauck CA, Laubach CA Jr. Quality of life and self-efficacy: gender and diagnoses considerations for management during cardiac rehabilitation. Journal of Cardiopulmonary Rehabilitation. 2003; 23(4):299–306. [PubMed: 12894004]
- Gottlieb SS, Kop WJ, Ellis SJ, Binkley P, Howlett J, O'Connor C, Cooper L. Relation of depression to severity of illness in heart failure (from Heart Failure And a Controlled Trial Investigating Outcomes of Exercise Training [HF-ACTION]). The American Journal of Cardiology. 2009; 103(9):1285–1289. [PubMed: 19406273]
- Gunstad J, Macgregor KL, Paul RH, Poppas A, Jefferson AL, Todaro JF, Cohen RA. Cardiac rehabilitation improves cognitive performance in older adults with cardiovascular disease. Journal of Cardiopulmonary Rehabilitation. 2005; 25(3):173–176. [PubMed: 15931022]
- Hlatky MA, Boineau RE, Higginbotham MB, Lee KL, Mark DB, Califf RM, Pryor DB. A brief self-administered questionnaire to determine functional capacity (The Duke Activity Status Index). The American Journal of Cardiology. 1989; 64(10):651–654. [PubMed: 2782256]
- Hu, L.-t.; Bentler, PM. Cutoff Criteria for Fit Indexes in Covariance Structure Analysis: Conventional Criteria Versus New Alternatives. Structural Equation Modeling. 1999; 6(1):1–55.
- Jefferson AL. Cardiac output as a potential risk factor for abnormal brain aging. Journal of Alzheimer's Disease. 2010; 20(3):813–821.
- Jefferson AL, Poppas A, Paul RH, Cohen RA. Systemic hypoperfusion is associated with executive dysfunction in geriatric cardiac patients. Neurobiology of Aging. 2007; 28(3):477–483. [PubMed: 16469418]
- Jefferson AL, Tate DF, Poppas A, Brickman AM, Paul RH, Gunstad J, Cohen RA. Lower Cardiac Output Is Associated with Greater White Matter Hyperintensities in Older Adults with Cardiovascular Disease. Journal of the American Geriatrics Society. 2007; 55(7):1044–1048. [PubMed: 17608877]
- Jencks S, Williams M, Coleman E. Rehospitalization among patients in Medicare fee-for-service program. New England Journal of Medicine. 2009; 360:1418–1428. [PubMed: 19339721]
- Jones CJ, Rikli RE. Measuring functional fitness of older adults. The Journal on Active Aging. 2002:24–30.
- Kapoor JR, Heidenreich PA. Obesity and survival in patients with heart failure and preserved systolic function: a U-shaped relationship. American Heart Journal. 2010; 159(1):75–80. [PubMed: 20102870]
- Kato N, Kinugawa K, Shiga T, Hatano M, Takeda N, Imai Y, Nagai R. Depressive symptoms are common and associated with adverse clinical outcomes in heart failure with reduced and preserved ejection fraction. Journal of Cardiology. 2012; 60(1):23–30. [PubMed: 22445598]
- Katon WJ. Clinical and health services relationships between major depression, depressive symptoms, and general medical illness. Biological Psychiatry. 2003; 54(3):216–226. [PubMed: 12893098]
- Katon W, von Korff M, Ciechanowski P, Russo J, Lin E, Simon G, Young B. Behavioral and clinical factors associated with depression among individuals with diabetes. Diabetes Care. 2004; 27(4): 914–920. [PubMed: 15047648]
- Kindermann I, Fischer D, Karbach J, Link A, Walenta K, Barth C, Böhm M. Cognitive function in patients with decompensated heart failure: the Cognitive Impairment in Heart Failure (CogImpair-HF) study. European Journal of Heart Failure. 2012; 14:403–413.
- Kravitz RL, Hays RD, Sherbourne CD, DiMatteo MR, Rogers WH, Ordway L, Greenfield S. Recall of recommendations and adherence to advice among patients with chronic medical conditions. Archives of Internal Medicine. 1993; 153(16):1869–1878. [PubMed: 8250648]
- Lysy Z, Da Costa D, Dasqupta K. The association of physical activity and depression in Type 2 diabetes. Diabetic Medicine. 2008; 25(10):1133–1141. [PubMed: 19046190]
- Maeda U, Shen BJ, Schwarz ER, Farrell KA, Mallon S. Self-Efficacy Mediates the Associations of Social Support and Depression with Treatment Adherence in Heart Failure Patients. International

- Journal of Behavioral Medicine. 2012 Advance online publication. doi:10.1007/s12529-011-9215-0.
- Marzolini S, Mertens DJ, Oh PI, Pyley MJ. Self-reported compliance to home-based resistance training in cardiac patients. European Journal of Cardiovascular Prevention and Rehabilitation. 2010; 17(1):42–49.
- Morris JC, Heyman A, Mohs RC, Hughes JP, van Belle G, Fillnbaum G, Clark C. . The consortium to establish a registry for Alzheimer's disease (CERAD). Part I. Clinical and neuropsychological assessment of Alzheimers disease. Neurology. 1989; 39(9):1159–1165. [PubMed: 2771064]
- Moser DJ, Hoth KF, Robinson RG, Paulsen JS, Sinkey CA, Benjamin ML, Schultz SK. Blood Vessel Function and Cognition in Elderly Patients With Atherosclerosis. Stroke. 2004:e369–e372. Published online before print. doi:10.1161/?01.STR.0000145050.35039.51. [PubMed: 15472091]
- Netz Y, Dwolatzky T, Zinker Y, Argov E, Agmon R. Aerobic fitness and multidomain cognitive function in advanced age. International Psychogeriatrics. 2011; 23(1):114–124. [PubMed: 20566000]
- Nixon PA, Joswiak ML, Fricker FJ. A six-minute walk test for assessing exercise tolerance in severely ill children. The Journal of Pediatrics. 1996; 129(3):362–366. [PubMed: 8804324]
- Norberg E, Bowman K, Lofgren B. Activities of daily living for old persons in primary health care with chronic heart failure. Scandinavian Joural of Caring Sciences. 2008; 22:203–210.
- Oka RK, DeMarco T, Haskell WL. Perceptions of physical fitness in patients with heart failure. Progress in Cardiovascular Nursing. 1999; 14(3):97–102. [PubMed: 10549047]
- Panagiotakos DB, Pitsavos C, Chrysohoou C, Tsetsekou E, Papageorgiou C, Christodolou G, Stefanadis C. Inflammation, coagulation, and depressive symptomatology in cardiovascular disease-free people; the ATTICA study. European Heart Journal. 2004; 25:492–499. [PubMed: 15039129]
- Papathanasiou G, Tsamis N, Georgiadou P, Adamopoulos S. Beneficial effects of physical training and methodology of exercise prescription in patients with heart failure. Hellenic Journal of Cardiology. 2008; 49(4):267–277. [PubMed: 18935714]
- Perry CK, Bennett JA. Heart disease prevention in women: promoting exercise. Journal of the American Academy of Nurse Practitioners. 2006; 18(12):568–573. [PubMed: 17184309]
- Pressler SJ, Subramanian U, Kareken D, Perkins SM, Gradus-Pizlo I, Sauve MJ, Shaw RM. Cognitive deficits in chronic heart failure. Nursing Research. 2010; 59(2):127–139. [PubMed: 20216015]
- Riegel B, Weaver TE. Poor sleep and impaired self-care: towards a comprehensive model linking sleep, cognition, and heart failure outcomes. European Journal of Cardiovascular Nursing. 2009:337–344. [PubMed: 19679510]
- Roger VL, Go AS, Lloyd-Jones D, Benjamin EJ, Berry JD, Borden WB, Lichtman J. Heart disease and stroke statistics—2012 update: a report from the American Heart Association. Circulation. 2012; 125(1):2–e220. Published online before print. doi: 10.1161/?CIR.0b013e31823ac046.
- Rogers RL, Meyer JS, Mortel KF. After reaching retirement age physical activity sustains cerebral perfusion and cognition. Journal of the American Geriatrics Society. 1990; 38(2):123–128. [PubMed: 2299115]
- Roshanaei-Moghaddam B, Katon WJ, Russo J. The longitudinal effects of depression on physical activity. General Hospital Psychiatry. 2009; 31(4):306–315. [PubMed: 19555789]
- Ruo B, Rumsfeld JS, Pipkin S, Whooley MA. Relation between depressive symptoms and treadmill exercise capacity in the Heart and Soul Study. American Journal of Cardiology. 2004; 94(1):96–99. [PubMed: 15219515]
- Rutledge T, Reis VA, Linke SE, Greenberg BH, Mill PJ. Depression in heart failure a meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. Journal of the American College of Cardiology. 2006; 48(8):1527–1537. [PubMed: 17045884]
- Skotzko CE, Krichten C, Zietowski G, Alves L, Freudenberger R, Robinson S, Gottlieb SS. Depression is common and precludes accurate assessment of functional status in elderly patients with congestive heart failure. Journal of Cardiac Failure. 2000; 6(4):300–304. [PubMed: 11145754]

Spaderna H, Zahn D, Schulze Scheleithoff S, Stadlbauer T, Rupprecht L, Smits JM, Weidner G. Depression and disease severity as correlates of everyday physical activity in heart transplant candidates. Transplant International. 2010; 23(8):813–822. [PubMed: 20158693]

- Spreen, O.; Strauss, E. A Compendium of Neuropsychological Tests. Oxford University ress; New York: 1991.
- Stanek KM, Gunstad J, Spitznagel MB, Waechter D, Hughes JW, Luyster F, Rosneck J. Improvements in cognitive function following cardiac rehabilitation for older adults with cardiovascular disease. The International Journal of Neuroscience. 2011; 121(2):86–93. [PubMed: 21062215]
- Suave MJ, Lewis WR, Blankenbiller M, Rickabaugh B, Pressler SJ. Cognitive impairments in chronic heart failure: a case controlled study. Journal of Cardiac Failure. 2009; 15(1):1–10. [PubMed: 19181287]
- Sullivan, M.; LaCroix, A.; Russo, J. Disability and psychological illness in primary care. In: Tansella, M.; Thornicroft, G., editors. Common mental disorders in primary care. Routledge; London: 1999.
- Thompson, B. Ten commandments of structural equation modeling. In: Grimm, L.; Yarnell, PR., editors. Reading and understanding more multivariate statistics. American Psychological Association; Washington, DC: 2000.
- Vogels RL, Scheltens P, Schroeder-Tanka JM, Weinstein HC. Cognitive impairment in heart failure: A systematic review of the literature. European Journal of Heart Failure. 2007; 9(5):440–449. [PubMed: 17174152]
- Vogels RL, van der Flier WM, van Harten B, Gouw AA, Scheltens P, Schroeder-Tanka JM, Weinstein HC. Brain magnetic resonance imaging abnormalities in patients with heart failure. European Journal of Hearth Failure. 2007; 9(10):1003–1009.
- Zuccala G, Marzetti E, Cesari M, Lo Monaco MR, Antonica L, Cocchi A, Bernabei R. Correlates of cognitive impairment among patients with heart failure: results of a multicenter survey. American Journal of Medicine. 2005; 118(5):496–502. [PubMed: 15866252]
- Zuccala G, Pedone C, Cesari M, Onder G, Pahor M, Marzetti E, Bernabei R. The effects of cognitive impairment on mortality among hospitalized patients with heart failure. American Journal of Medicine. 2003; 115:97–103. [PubMed: 12893394]




Figure 1. Model Examining the Role of Cardiovascular Fitness and Exercise Adherence in the Relationship Between Depression and Cognitive Function in HF Patients (N=158)

Notes. The pathways between cognitive function and animal was fixed as was the pathway between CV Fitness and Peak VO₂. Standardized parameters estimates are presented in the model; significance levels for these paths are based on the unstandardized estimates. Abbreviations—BDI-II = Beck Depression Inventory-II; TMTA = Trail Making Test A; TMTB = Trail Making Test B; FAB = Frontal Assessment Battery; CVLT= California Verbal Learning Test Delayed Recall; Animal = Animals Fluency Test; 2MST = 2-minute step test; CV = Cardiovascular

* p < .05

Alosco et al. Page 15

Table 1Demographic, Clinical, and Cognitive Characteristics of 158 Older Adults with Heart Failure

Demographic Characteristics								
Age, mean (SD)	68.96 (9.13)							
Sex (% Women)	36.7							
Race (% Caucasian)	82.3							
Years of Education, mean (SD)	13.39 (2.74)							
Clinical Characteristics								
Left Ventricular Ejection Fraction, mean (SD)	40.70 (14.39)							
Beck Depression Inventory-II, mean (SD)	7.67 (7.57)							
Exercise adherence, % non-adherent	50.6							
2-Minute Step Test Males	62.90 (24.65)							
2-Minute Step Test Females	55.89 (21.14)							
2-Minute Step Test Overall Sample	60.32 (23.59)							
DASI Peak Oxygen Uptake (mL/min), mean (SD)	24.46 (6.53)							
Cognitive Test Performance, T-score mean (SD)								
Trail Making Test A	49.93 (10.56)							
Trail Making Test B	43.62 (18.63)							
Frontal Assessment Battery	43.08 (21.85)							
CVLT Long Delayed Recall	47.12 (10.30)							
Animals	54.72 (11.24)							

DASI = Duke Activity Status Index; CVLT = California Verbal Learning Test

Table 2

Bivariate Covariance Matrix (N=158)

	2MST	BDI	CVLT	TMTA	ТМТВ	FAB	Animal	Exercise	Peak
2MST	556.69								
BDI	-35.89	57.37							
CVLT	-1 .23	-3.82	106.15						
TMTA	66.54	-18.64	24.74	111.51					
TMTB	127.53	-56.41	49.65	116.47	346.94				
FAB	124 .13	-25.50	57.68	88.95	193.34	477.44			
Animal	40.05	-9.74	28.28	26.75	64.27	71.08	126.30		
Exercise	132 .82	-41.06	5.60	12.77	25.46	44.86	64.16	1153.96	
Peak	82 .73	-23.54	3.24	21.57	39.50	35.89	19.02	71.64	42.61

Note. 2MST = 2-minute step test; BDI= Beck Depression Inventory-II; CVLT = California Verbal Learning Test-II Long Delayed Recall; TMTA = Trail Making Test A; TMTB = Trail Making Test B; FAB = Frontal Assessment Battery; Animal = Animals Fluency Test; Peak = Estimate Peak Oxygen Uptake; Exercise = Exercise Adherence