


STUDY PROTOCOL

T-RAC: Study protocol of a randomised clinical trial for assessing the acceptability and preliminary efficacy of adding an exergame-augmented dynamic imagery intervention to the behavioural activation treatment of depression



Alexandru Tiba^{1*}, Marius Drugaș¹, Ioana Sârbu¹, Trip Simona¹, Carmen Bora¹, Daiana Miclăuș¹, Laura Voss², Ioana Sanislav¹, Daniel Ciurescu³

1 Department of Psychology, University of Oradea, Oradea, Romania, **2** The Hull York Medical School, University of York, York, United Kingdom, **3** Faculty of Medicine, Transilvania University, Brașov, Romania

* alexandrutiba@gmail.com

OPEN ACCESS

Citation: Tiba A, Drugaș M, Sârbu I, Simona T, Bora C, Miclăuș D, et al. (2023) T-RAC: Study protocol of a randomised clinical trial for assessing the acceptability and preliminary efficacy of adding an exergame-augmented dynamic imagery intervention to the behavioural activation treatment of depression. PLoS ONE 18(7): e0288910. <https://doi.org/10.1371/journal.pone.0288910>

Editor: Vanessa Carels, PLoS ONE, UNITED STATES

Received: June 16, 2023

Accepted: June 23, 2023

Published: July 31, 2023

Copyright: © 2023 Tiba et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: No datasets were generated or analysed during the current study. All relevant data from this study will be made available upon study completion.

Funding: A.T PN-III-P1-1.1-TE-2021-1090, within PNCDI III. Ministry of Research, Innovation and Digitization, CNCS - UEFISCDI, <https://uefiscdi-direct.ro/> The funders did not and will not have a role in study design, data collection and analysis,

Abstract

Background

Improving the existent effective treatments of depression is a promising way to optimise the effects of psychological treatments. Here we examine the effects of adding a rehabilitation type of imagery based on exergames and dynamic simulations to a short behavioural activation treatment of depression. We investigate the acceptability and the efficacy of an exergame-augmented dynamic imagery intervention added to behavioural activation treatment and associated mechanisms of change.

Methods and analyses

In a two-arm pilot randomised controlled trial, the acceptability and preliminary efficacy of an exergame-augmented dynamic imagery intervention added to behavioural activation treatment for depressed individuals will be assessed. Participants (age 18–65) meeting criteria for depression are recruited by media and local announcements. 110 participants will be randomly allocated to behavioural activation plus imagery group or to standard behavioural activation group. The primary outcome is depressive symptom severity (Beck Depression Inventory II) and secondary outcomes are anhedonia, apathy and behavioural activation and avoidance. The outcomes are assessed at baseline, mid treatment, post-treatment and 3-month follow-up. Moderation and mediation analyses will be explored. An intention-to-treat approach with additional per-protocol analysis will be used for data analysis.

decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Major depressive disorder (MDD) is a common mental health problem that results in severe impairment of the life of affected individuals [1]. Although there are several effective treatments of depression, their efficacy was challenged, only a moderate-small effect being evidenced after removing the existing research biases [2]. Estimates suggest that 59% of depressed patients do not even respond to psychological treatments (50% reduction in symptoms) and two thirds do not remit [2]. To this shortcoming it adds the high rates of relapse, recurrence, residual symptoms, and the fact that many affected individuals receive no treatment. It is estimated that after treatment, 30 to 50% of recovered patients are still troubled by residual symptoms and up to 75% of patients relapse [3]. Improving the treatment for depression has a major impact on global mental health and economy.

Finding efficient ways to treat depression has always been a preoccupation for both researchers and clinicians. Despite significant efforts, the efficacy of treatments for depression did not significantly improve in time [4]. Targeting new mechanisms and optimising the already effective interventions are promising solutions to advance the treatment of depression [5].

Mental rehearsal interventions are considered promising stand-alone or added-on treatments for depression [5–9]. Recently, the effect of imagery has been tested by adding imagery to behavioural activation treatment for depression. Behavioural activation treatment is one of the most effective and robust psychological treatments for depression [10, 11]. Although adding imagery interventions to behavioural activation brings great promise for improving the treatment of depression, there are several potential improvements for imagery interventions. Renner et al. [9] have recently pointed out the limitations of mental imagery interventions in depression: only individuals who vividly imagine actions benefit from intervention; depressed individuals have difficulties vividly simulating actions, not knowing the ingredients of a prospective mental rehearsal of actions.

We suggest that a major source of potential improvements for mental imagery interventions comes from paying more attention to the contribution of the ongoing states of the sensorimotor and affective brain systems (motor, affective, and visual) to mental simulations. Considering the characteristics of the sensorimotor and affective brain systems involved in the mental construction of those experiences results in several advantages for imagery interventions. First, the differences in the states of the sensorial, affective, and motor systems recruited for simulation should be considered during imagery interventions for depressed individuals. For instance, depressed individuals show large differences in the state of modality-specific brain systems, having attenuated excitation of the motor cortex [12, 13] and deficient recruitment of brain systems involved in positive affect [14], which may influence the effect of the interventions. Second, considering how particular imagery modalities (e.g., visual versus motor) are used during the imagery process in depressed individuals may show a significant advantage. Differences in using some modalities over others may result in differences in the effect of imagery over behaviour [15]. For instance, the recruitment of the motor brain in imagery and the effect of mental imagery on action largely depend on the success of using kinesthetic rather than visual imagery [16]. Depressed individuals have more impaired motor imagery than visual imagery and compensate for this deficit by using modalities, which may limit the impact of imagery interventions [17]. Third, adopting an understanding of the mental simulation of feelings as mental re-activation of body sensations in imagery intervention may also bring a significant advantage for intervention. Recent research showed that felt sensations in the body are core components of feelings [18, 19], they can be maintained as representations in working memory [20], they are attenuated in depressed individuals [21], and that

prospective simulations of sensations felt in the body during feelings in positive situations are attenuated in individuals with high levels of depressive symptomatology [22].

Here we propose that we may use these advantages and significantly improve imagery interventions by focusing on enhancing the embodiment of mental rehearsal in imagery-based treatments for depressed individuals. A large body of research suggests that the effect of mental simulations on emotion and behaviour heavily depends on their embodiments, or the efficient recruitment of specific sensorimotor and affective modalities in mental simulations by attention, verbal, and memory controls [23]. For instance, when simulating and paying attention to visual details rather than proprioceptive sensations, one will recruit the visual cortex for action simulation rather than the interoceptive and motor cortex, and vice versa [24]. Mental rehearsal of action is efficient for action and affective control when it adequately recruits motor and affective simulations [25]. Depressed individuals have difficulties recruiting motor and affective brain areas in their thinking. Thus, to optimise imagery treatment for depressed individuals, one should enhance the recruitment of modal (motor and affective) brain systems in the mental simulation process. When deficiencies of the modal brain areas required for imagery simulations are stable alterations (i.e., cognitive deficits of cognition), the intervention should follow a rehabilitation approach similar to motor rehabilitation imagery used in patients with brain damage (e.g., damages due to stroke). We focus on two potential solutions for improving imagery based on embodied processes: exergames and dynamic imagery.

A promising way to support the embodiment of simulations of action in depressed individuals is by using remote kinematics in the form of exergaming (XboxKinect™ games). Previous research showed that using exergaming results in the augmentation of motor (kinesthetic) imagery [26–28], increased motor rehabilitation in stroke patients [29], motor learning [30] and it is efficient as a stand-alone intervention for the reduction of depressive symptomatology [31–34]. Furthermore, exergaming has an increased acceptability and feasibility in various populations [29]. Based on these results, using exergaming as a tool for motor rehabilitation of action cognition in imagery interventions for depressed individuals seems to be worth investigating.

Another promising solution came from dynamic imagery. Dynamic imagery, as opposed to static imagery (mental rehearsal of movement without overt movement) requires mimicry of simulated movements [35]. Previous research has shown a *dynamic superiority effect*, individuals remembering better moving rather than static stimuli [36]. Similarly, a dynamic superiority effect has been evidenced for imagery. In comparison with static imagery, dynamic imagery results in higher vividness of mental imagery [37] motor learning [35] and behavioural performance [38]. Moreover, the imagery deficits which are evidenced in static motor imagery tasks are not evident when a dynamic strategy of motor imagery is applied for the same population [39].

Given that imagery methods added to behavioural activation treatments are promising for the treatment of depression [6, 9] and that dynamic imagery and stimulation by remote kinematic may help overcome limitations of existing imagery interventions, we intend to test the effect of adding a dynamic imagery intervention augmented by exergames to a short behavioural activation treatment [40].

Starting from the idea that action cognition and motor imagery deficits are stable alterations in depressed individuals [41–43] we use a rehabilitation type of motor imagery intervention based on principles of rehabilitation of motor imagery for individuals with neurological conditions. To this end, we integrate forward modelling of action and motor imagery [44], remote kinematics (Kinect) and embodied cognition account (action support of mental simulations, [25, 45]). This intervention is a new one that proved efficient in clinical work (Tiba, [Unpublished protocol]), [46]. There are several steps in the intervention. In the first session,

the therapists introduce the *dynamic simulation routine*: how to support mental imagery of activities through gestures, language, and attention. Then, the patients undergo an exergame for 5 to 10 minutes and an “actfulness” exercise focusing on feelings of movement and motor simulation. After the exercise, the participants dynamically imagine a planned activity. In session two, patients are thought to restructure action memories by focusing on movements. Deficient action simulations are rehabilitated by (a) partial movements (they alternate covert with dynamic-partial movements simulations in response to stimuli); (b) linguistic supports (training in gerundival perceptions that is to recognize and name stimuli by actions; for instance, a door is recognized as a door to open; [47]), (c) enhanced perceptual and affective simulations (PETTLEP system, [48]), and (d) episodic memory support (they have to form future memories of *action cores*-last sequence of movement before perception of desired environmental change, correct them by experience, and remember at the end of the day). Thus, it is a rehearsal training that mixes actual and mental practice [49, 50] with augmentation of the motor component of simulations in thinking by enhancing *gestures, language, and episodic memory* as controls of simulation, and is applied to promote the use of motor simulations in everyday life.

We expect that adding a dynamic imagery intervention will augment the effects of behavioural activation treatment (BA) on severity of depressive symptoms. Second, we investigate whether adding the dynamic imagery intervention to behavioural activation treatment has any effect on secondary outcomes such as anhedonia, apathy, and vividness of imagery. Third, we investigate the moderators and mediators of the effects of the intervention.

Methods and analysis

Study design

The study is a two-arm phase II randomised controlled trial (RCT) with a parallel design to compare a short behavioural activation (BA) treatment with BA plus dynamic imagery enhanced by exergames (XboxKinect™). We recruit participants by online media, advertisements, and by contacting local psychiatrists. After the online screening and baseline assessment, participants are randomised to either BA or BA plus T-RAC. Then the participants are enrolled in the treatment. Participants complete a weekly assessment, one assessment of mediators at week six, one assessment after treatment and one assessment at three months. The consort flow for the Study is presented in Fig 1.

Ethics and dissemination. This trial is approved by the ethics committee of the Faculty of Socio-Humanistic Sciences of University of Oradea (no: 2394/18.11.2022). Informed consent is required from participants before enrollment in the study. The results will be submitted for publication in peer-reviewed journals and presented at conferences.

Trial registration number. This trial was registered in the Clinical Trials Register: NCT05625230.

Recruitment of participants

The recruitment of the participants will follow the CONSORT chart that will guide the flow of the participants. 110 participants will be recruited so that the study will have sufficient power to detect the differences at a 20% attrition rate. Participants will be selected through announcements about the study both online and at the local clinics. Participants are invited to complete an online screening questionnaire, including the PHQ-9 scale and questions for checking the eligibility criteria. Participants with a PHQ-9 score of ten or higher are programmed for the SCID-5-CV interview administered by telephone or face-to-face. Participants who are eligible




Fig 1. The study flow.

<https://doi.org/10.1371/journal.pone.0288910.g001>

will be randomly assigned (using Redcap) to one of the two groups: (1) behavioural activation (BA) or (2) BA plus rehabilitation of action simulations (BA Plus T-RAC).

Participants allocated to the behavioural activation arm will undergo a standard BA treatment. A therapist will administer the treatment for 8 sessions based on a manual built on the behavioural treatment protocol described by Martell et al. [51] and used in the COBRA trial

[52]. Each session is focused on reviewing the activity log, discussing difficulties in the implementation of previously planned activities, planning “antidepressant” activities for the next week, and finding support for the implementation of the activity. Participants allocated to the BA plus T-RAC will undergo the same BA protocol plus an augmented imagery intervention. Each session is focused on reviewing the activity log, planning “antidepressant” activities for the next week, and finding support for the implementation of the activity. The difference is that after activity planning, the participant will follow Kinect for 10 minutes and will imagine one planned activity using actfulness. From session two, restructuring action memories are added.

Eligibility

The inclusion criteria were age from 18 to 65, a score of 10 or higher on the PHQ-9 and a diagnosis of depression based on clinical interview. The exclusion criteria were: significant intellectual impairment; meeting criteria for a current psychotic or substance abuse disorder; history of mania/hypomania; significant changes in the dose of antidepressant medication during the past month; receiving psychological therapy; and the presence of suicidal intent.

Management of suicidal ideation

We assess the suicidal ideation and intent at all assessments points of the study. Suicidal ideation is measured by the PHQ-9 item 9, BDI II, and clinical interview. If suicidal thoughts are reported, the C-SSRS interview is applied [53]. If suicidal intent (both unclear and clear) is identified the patient is not recruited in the study and contacted by phone by the clinical psychologist for proper recommendation to medical services. The participants with low risk suicidal ideation (suicidal ideation with no intent) are required to sign a non-suicide contract to continue the participation in the study. If suicide intent is identified, the participant is excluded from the study and necessary steps for medical assistance are followed (visit to psychiatrist, inpatient care). The suicide protocol is supervised by a licensed clinical psychologist.

Outcomes

Primary outcome. *Depressive symptom severity* is assessed with the Beck Depression Inventory-II (BDI-II) [54]. BDI-II comprises 21 items assessing the DSM symptoms of the major depression episode during the past 2 weeks. Each question is scored on a 4-point Likert scale ranging from 0 to 3, with higher scores indicating higher depression. The BDI-II manual states that scores between 0 and 13 indicate minimal depression, 14 to 19 indicate mild depression, 20 to 28 indicate moderate depression, and 29 to 63 indicate severe depression [55]. Psychometric strengths such as validity and reliability were supported by numerous studies across many populations and cultural backgrounds [54, 56].

Secondary outcomes. *Depressive symptoms severity* is assessed with the Patient Health Questionnaire- 9 [57]. PHQ-9 scale contains nine items that measure the severity of nine depressive symptoms on a 4-point scale during the past two weeks. Each item is rated on a 4-point scale ranging from 0 (not at all) to 3 (nearly every day).

Depression remission is measured using The Structured Clinical Interview for DSM-5 Clinician Version [SCID-5-CV; 58], affective module at post measurement and follow-up. SCID-5 is a structured clinical interview used to determine the core diagnoses according to DSM-5 guidelines. Participants are considered remitted when they no longer fulfil diagnostic criteria for a major depression episode.

Depression response is assessed based on BDI II scores. To identify responders to treatment we compute the reliable change index. A reliable change index of more than 1.96 will classify a participant as a responder [59].

Anhedonia severity is assessed with the Snaith-Hamilton Pleasure Scale [60] SHAPS is a 14-item scale assessing the degree to which a person is able to experience pleasure or the anticipation of a pleasurable experience. Participants respond on a four points scale from strongly disagree to strongly agree. Each "disagree" response receives one point, whereas each "agree" response receives zero points. Thus, the final score ranges from 0 to 14.

Apathy level is assessed with the Apathy-Motivation Index [61]. AMI is a 18-item questionnaire used to identify three dimensions of apathy: Behavioural Activation (propensity to initiate goal-directed behaviour), Social Motivation (level of engagement in social interactions), and Emotional Sensitivity (feelings of positive and negative affection). Higher scores indicate greater apathy. Item scores are averaged to get scores for subscales and a total score (range 0–4) [61].

Anxiety severity is assessed with the Generalised Anxiety Scale 7 [GAD-7; 62]. GAD-7 is a self-report scale measuring the severity of generalised anxiety disorder (GAD) during the past 2 weeks. It contains seven items related to DSM criteria for anxiety. Participants respond on a 4-point scale. Each item is rated on a 4-point scale ranging from 0 (not at all) to 3 (nearly every day).

Health and disability level outcome is assessed with The World Health Organisation Disability Assessment Schedule (WHODAS 2.0) 12 –Self-Report Version WHODAS 12), [63]. WHO-DAS 12 is a practical, general assessment instrument. It has 6 subscale, each subscale containing two questions from the corresponding subscale in the full version: cognition, mobility, self-care, getting along, life activities and participation. The response options for each item ranged from 0 (none), to 4 (extreme/cannot do). The individual scores in each subscale are calculated by adding the two item scores. The total score ranges from 0–48.

Adverse and unwanted effects of the experimental intervention are assessed using the Negative Effects Questionnaire [64]. NEQ is a 20-item scale used to assess the adverse and unwanted effects of psychological treatments. Participants respond on a 5-point Likert-scale about the presence of adverse events during treatment such as symptoms (e.g., stress, shame, anxiety), quality (e.g., confidence regarding treatment and therapy), dependency (e.g., addiction to treatment medicine or therapy), stigma (e.g., feeling ashamed in front of other people), hopelessness, and failure (e.g., when thinking that things could get better) [65].

The vividness of motor imagery is assessed using the Vividness of Motor Imagery Questionnaire-2 (VMIQ-2) [66]. The vividness of imagining twelve movements is measured using a Likert scale from 1 (perfectly clear and vivid as you would normally picture the image) to 5 (no image or sensation at all, you are only aware you are "thinking" of the movement). The VMIQ-2 version was found to be as a psychometrically valid questionnaire) [66, 67].

The level of rewards is assessed with "The environmental and reward observation scale" (EROS); [68]. EROS consists of 10 items (e.g., "It is easy for me to find enjoyment in my life"), where responses are given on a Likert-type scale ranging from "completely disagree" to "absolutely agree", on a 4-point scale. A higher score indicates a larger experience of reinforcing events from the environment.

The level of rumination is assessed with the Ruminative Response Scale-SV [69]. Participants respond on a 4-point Likert scale ranging from 1 ("almost never") to 4 ("almost always"). The scale describes different components of the rumination response to depressed mood such as self-focused rumination (e.g., "I can't stop thinking about some things"), symptom focused rumination (e.g., "I have never been able to distract myself from unwanted thoughts") or rumination focused on the possible consequences and causes of the mood (e.g., "When I have a

problem it will gnaw in my mind for a long time"). Brooding involves self-criticism, whereas reflection is related to more active problem solving.

Activation and avoidance is assessed using the Behavioral Activation for Depression Scale—Short Form (BADS-SF) [70]. BADS-SF is a 9-item scale used to assess the activation over the past week. It is organised in activation and avoidance subscales. BADS-SF is used to track when and how participants were activated during BA treatment.

Working memory is assessed with a Backward digit span task. Participants are given a series of digits and asked to repeat them backward. The score is based on how many numbers participants are able to repeat. A higher score indicates better working memory. Pairs of the sequences are read aloud to participants. The subsequent set of sequences will be one digit longer if both are successfully repeated and one digit shorter if none of the pair's sequences is correctly repeated. The following pair of sequences will have the same length if one of the pair's sequences is incorrectly repeated while the other sequence is correctly repeated [71].

Executive functioning is assessed with the verbal fluency test. In the verbal fluency test, participants are asked to say as many words as possible with a given letter (consonant F, A or S) for 1 minute.

Verb fluency. In the verb fluency task, participants are asked to produce as many verbs as possible for one minute. The participant's score in each task is the number of total correct words.

Acceptability ratings. Acceptability is measured as satisfaction with the intervention, intention to continue and the intention to recommend the intervention assessed on a 5-points Likert scale. Drop-out rates are analysed as well.

Affect and behaviour monitoring scale. Changes in functioning, emotions, behaviour profile, efficacy, and difficulty of simulation are assessed. For each item, the score ranges from 1 (not at all) to 7 (very much).

Proposed sample size

The sample size was established using G*Power and selecting the difference between two independent means. Forty-four participants per arm will be required to obtain a 90% statistical power. This estimation is based on previous studies targeting robust processes in depressed individuals (rumination) and between-treatment effect size of Cohen's $d = 0.7$. Assuming a dropout rate of 20%, 55 patients are required in each treatment arm, resulting in a total of 110 participants.

Statistical analyses

Multiple imputation methods will be selected to address the missing data. All randomised patients will be included in intention-to-treat analyses (ITT). Per-protocol analysis will be conducted separately. Analyses will be computed with an alpha significance level of 5% and two-tailed analyses. Analyses will be performed using IBM SPSS. Comparisons will be computed at a 5% significance level, and BDI II is the primary outcome variable considered as continuous variables. Differences between groups at the posttest will be analysed with a mixed ANOVA having group as the main effect and group *time as an interaction effect. Subgroup analyses will be conducted for anhedonia, apathy, motor imagery, and rumination. F-statistics and p values will be reported for main and interaction effects. The results will be considered as supporting evidence if the F test of the interaction effect is statistically significant (group*time interaction term). Potential mediators of treatment effects will be analysed exploratorily using Hayes model 4 analyses for the primary (depression severity) and secondary outcomes (anhedonia, and apathy). Hayes model 1 will be used to analyse the moderation. Between group

effect sizes will be calculated based on Cohen's *d*. Treatment completers will be analysed using per protocol analyses. Participants are considered "remitted" if they no longer fulfil the SCID-5 criteria for major depression episodes.

Patient and public involvement

Participants and the public are not involved in the development of the study design. The intervention was modified based on piloting the procedures with healthy individuals and case studies of patients with depression. Based on the feedback from initial testing, procedures were modified to increase acceptability. Both XboxKinect™ exergames and imagery interventions were well received by various clinical and non-clinical populations in previous RCTs. Similarly, behavioural activation treatment is a well-received and efficient treatment of depression. Study results are disseminated through presentations at conferences and publications in major outlets for psychology. Local therapists will be trained to deliver behavioural activation. The public will know about the results available on the website and in the media.

Discussion

The study is the first to evaluate an exergame-augmented dynamic imagery intervention added to behavioural activation treatment. It is expected that adding the exergame-augmented dynamic imagery intervention will increase the acceptability of the treatment and decrease the severity of depression, anhedonia, and apathy. We also expect, based on moderation analyses, that this effect to be more pronounced in participants with difficulties of motor imagery at baseline. We further investigate mediators of the treatment effect on the severity of depression and secondary outcomes.

This study is the first to investigate the added benefit of exergame-augmented dynamic motor imagery intervention to behavioural activation treatment. There is an expected advantage that this intervention will make the treatment more accepted (due to XboxKinect™) and will overcome the deficiencies in vividness of imagery and the recruitment of motor networks in imagery. If efficient it may work through different mechanisms than the standard BA treatment.

There are several strengths of the study: (1) it is a theory-driven intervention which could show an efficacy advantage, (2) it includes attractive exergames that may show additional benefits (exergame time may be increased and transformed into physical exercise) (3) we included a clinician administered semi-structured interview (SCID-5-CV), (4) we planned for a mid-treatment assessment to evaluate mechanisms of change. At session six, the participants will complete measures of the mechanisms of change, which will be carried out based on temporal precedence criteria [72], (5) assessors are blinded to participants' treatment conditions.

There are several limitations of the study: (1) it is not a large scale RCT some limitations being related to the small sample size, (2) as the sample size was determined based on post-treatment differences, moderation and mediation analyses being underpowered, (3) participants with high risk of suicide will be excluded based on a score C-SSRS scale [53], limiting the generalisation of the results to participants with high risk of suicide.

This study investigates the acceptability, feasibility, and preliminary efficacy of adding an exergame-augmented dynamic imagery rehabilitative intervention to behavioural activation treatment. Thus, an ecological rehabilitative intervention for action cognition is available. This study increases our knowledge of highly available treatments with increased accessibility for individuals with depression.

Supporting information

S1 Checklist. SPIRIT 2013 checklist: Recommended items to address in a clinical trial protocol and related documents*.

(DOC)

S1 Fig.

(DOC)

S1 File.

(PDF)

S2 File.

(PDF)

Author Contributions

Conceptualization: Alexandru Tiba, Laura Voss.

Formal analysis: Alexandru Tiba, Carmen Bora.

Funding acquisition: Alexandru Tiba.

Investigation: Alexandru Tiba, Ioana Sârbu, Daiana Miclăuș, Ioana Sanislav.

Methodology: Alexandru Tiba, Carmen Bora, Laura Voss.

Project administration: Alexandru Tiba, Marius Drugaș, Ioana Sârbu.

Resources: Alexandru Tiba.

Supervision: Alexandru Tiba, Ioana Sârbu, Carmen Bora.

Validation: Carmen Bora.

Writing – original draft: Alexandru Tiba, Laura Voss.

Writing – review & editing: Alexandru Tiba, Marius Drugaș, Ioana Sârbu, Trip Simona, Carmen Bora, Daiana Miclăuș, Laura Voss, Ioana Sanislav, Daniel Ciurescu.

References

1. World Health Organization (2017). Depression and Other Common Mental Disorders: Global Health Estimates. Geneva. Licence: CC BY-NC-SA 3.0 IGO.
2. Cuijpers P., Karyotaki E., Ciharova M., Miguel C., Noma H., & Furukawa T. A. (2021). The effects of psychotherapies for depression on response, remission, reliable change, and deterioration: A meta-analysis. *Acta psychiatrica Scandinavica*, 144(3), 288–299. <https://doi.org/10.1111/acps.13335> PMID: 34107050
3. Hollon S. D., DeRubeis R. J., Shelton R. C., Amsterdam J. D., Salomon R. M., O'Reardon J. P., et al. (2005). Prevention of relapse following cognitive therapy vs medications in moderate to severe depression. *Archives of General Psychiatry*, 62, 417–422. <https://doi.org/10.1001/archpsyc.62.4.417> PMID: 15809409
4. Cuijpers P., Noma H., Karyotaki E., Vinkers C. H., Cipriani A., & Furukawa T. A. (2020). A network meta-analysis of the effects of psychotherapies, pharmacotherapies and their combination in the treatment of adult depression. *World Psychiatry*, 19(1), 92–107. <https://doi.org/10.1002/wps.20701> PMID: 31922679
5. Holmes E. A., Ghaderi A., Harmer C. J., Ramchandani P. G., Cuijpers P., Morrison A. P., et al. (2018). The Lancet Psychiatry Commission on psychological treatments research in tomorrow's science. *The Lancet. Psychiatry*, 5(3), 237–286. [https://doi.org/10.1016/S2215-0366\(17\)30513-8](https://doi.org/10.1016/S2215-0366(17)30513-8) PMID: 29482764
6. Pellas J., Renner F., Ji J. L., & Damberg M. (2022). Telephone-based behavioral activation with mental imagery for depression: A pilot randomized clinical trial in isolated older adults during the Covid-19

- pandemic. *International journal of geriatric psychiatry*, 37(1), <https://doi.org/10.1002/gps.5646> PMID: 34729823
7. Renner F., Ji J. L., Pictet A., Holmes E. A., & Blackwell S. E. (2017). Effects of Engaging in Repeated Mental Imagery of Future Positive Events on Behavioural Activation in Individuals with Major Depressive Disorder. *Cognitive Therapy and Research*, 41(3), 369–380. <https://doi.org/10.1007/s10608-016-9776-y> PMID: 28515538
 8. Renner F., Murphy F. C., Ji J. L., Manly T., & Holmes E. A. (2019). Mental imagery as a “motivational amplifier” to promote activities. *Behaviour Research and Therapy*, 114, 51–59. <https://doi.org/10.1016/j.brat.2019.02.002> PMID: 30797989
 9. Renner F., Werthmann J., Paetsch A., Bär H. E., Heise M., & Bruijninks S. J. (2021). Prospective mental imagery in depression: impact on reward processing and reward-motivated behaviour. *Clinical Psychology in Europe*, 3(2), 1–16. <https://doi.org/10.32872/cpe.3013> PMID: 36397959
 10. Cuijpers P., van Straten A., & Warmerdam L. (2007). Behavioral treatment of depression: A metaanalysis of activity scheduling. *Clinical Psychology Review*, 27(3), 318–326.
 11. Ekers D., Webster L., Van Straten A., Cuijpers P., Richards D., & Gilbody S. (2014). Behavioural activation for depression; an update of meta-analysis of effectiveness and sub group analysis. *PLoS one*, 9(6), e100100. <https://doi.org/10.1371/journal.pone.0100100> PMID: 24936656
 12. Cotovio G., Rodrigues da Silva D., Real Lage E., Seybert C., & Oliveira-Maia A. J. (2022). Hemispheric asymmetry of motor cortex excitability in mood disorders—Evidence from a systematic review and meta-analysis. *Clinical neurophysiology: official journal of the International Federation of Clinical Neurophysiology*, 137, 25–37. <https://doi.org/10.1016/j.clinph.2022.01.137> PMID: 35240425
 13. Maeda F., Keenan J. P., & Pascual-Leone A. (2000). Interhemispheric asymmetry of motor cortical excitability in major depression as measured by transcranial magnetic stimulation. *The British journal of psychiatry: the journal of mental science*, 177, 169–173. <https://doi.org/10.1192/bjp.177.2.169> PMID: 11026958
 14. Heller A. S., Johnstone T., Shackman A. J., Light S. N., Peterson M. J., Kolden G. G., et al. (2009). Reduced capacity to sustain positive emotion in major depression reflects diminished maintenance of fronto-striatal brain activation. *Proceedings of the National Academy of Sciences of the United States of America*, 106(52), 22445–22450. <https://doi.org/10.1073/pnas.0910651106> PMID: 20080793
 15. Bonnet C., Bayram M., El Bouzaïdi Tiali S., Lebon F., Harquel S., Palluel-Germain R., et al. (2022). Kinesthetic motor-imagery training improves performance on lexical-semantic access. *PLoS one*, 17(6), e0270352. <https://doi.org/10.1371/journal.pone.0270352> PMID: 35749512
 16. Yang Y. J., Jeon E. J., Kim J. S., & Chung C. K. (2021). Characterization of kinesthetic motor imagery compared with visual motor imageries. *Scientific Reports*, 11(1), 1–11. <https://doi.org/10.1038/s41598-021-82241-0> PMID: 33580093
 17. Bennabi D., Monnin J., Haffen E., Carvalho N., Vandel P., Pozzo T., et al. (2014). Motor imagery in unipolar major depression. *Frontiers in Behavioral Neuroscience*, 8, Article 413. <https://doi.org/10.3389/fnbeh.2014.00413> PMID: 25538580
 18. Harrison N. A., Gray M. A., Gianaros P. J., & Critchley H. D. (2010). The embodiment of emotional feelings in the brain. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 30(38), 12878–12884. <https://doi.org/10.1523/JNEUROSCI.1725-10.2010> PMID: 20861391
 19. Nummenmaa L., Glerean E., Hari R., & Hietanen J. K. (2014). Bodily maps of emotions. *Proceedings of the National Academy of Sciences*, 111(2), 646–651. <https://doi.org/10.1073/pnas.1321664111> PMID: 24379370
 20. Mikels J. A., & Reuter-Lorenz P. A. (2019). Affective Working Memory: An Integrative Psychological Construct. *Perspectives on psychological science: a journal of the Association for Psychological Science*, 14(4), 543–559. <https://doi.org/10.1177/1745691619837597> PMID: 31059662
 21. Lyons N., Strasser A., Beitz B., Teismann T., Ostermann T., Anderle L., et al. (2021). Bodily maps of emotion in major depressive disorder. *Cognitive Therapy and Research*, 45(3), 508–516. <https://doi.org/10.1007/s10608-020-10195-0>
 22. Tiba A. I. & Manea L (2018a). The Vividness of Imagining Emotional Feelings is Attenuated in Dysphoria and Predicts the Experience of Positive Emotional Feelings. *Journal of Clinical Psychology*.
 23. Winkielman P., Niedenthal P. M., & Oberman L. (2008). The embodied emotional mind. In Semin G. R. & Smith E. R. (Eds.), *Embodied grounding: Social, cognitive, affective, and neuroscientific approaches* (pp. 263–288). Cambridge University Press.
 24. Moulton S. T., & Kosslyn S. M. (2009). Imagining predictions: mental imagery as mental emulation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1521), 1273–1280. <https://doi.org/10.1098/rstb.2008.0314> PMID: 19528008
 25. Glenberg A. M. (2010). Embodiment as a unifying perspective for psychology. *Wiley Interdisciplinary Reviews: Cognitive Science*, 1, 586–596. <https://doi.org/10.1002/wcs.55> PMID: 26271505

26. Wriessnegger S. C., Steyerl D., Koschutnig K., & Müller-Putz G. R. (2014). Short time sports exercise boosts motor imagery patterns: implications of mental practice in rehabilitation programs. *Frontiers in human neuroscience*, 8, 469. <https://doi.org/10.3389/fnhum.2014.00469> PMID: 25071505
27. Wriessnegger S. C., Brunner C., & Müller-Putz G. R. (2018). Frequency Specific Cortical Dynamics During Motor Imagery Are Influenced by Prior Physical Activity. *Frontiers in psychology*, 9, 1976. <https://doi.org/10.3389/fpsyg.2018.01976> PMID: 30410454
28. Wriessnegger S. C., Müller-Putz G. R., Brunner C., & Sburlea A. I. (2020). Inter- and Intra-individual Variability in Brain Oscillations During Sports Motor Imagery. *Frontiers in human neuroscience*, 14, 576241. <https://doi.org/10.3389/fnhum.2020.576241> PMID: 33192406
29. Almasi S., Ahmadi H., Asadi F., Shahmoradi L., Arji G., Alizadeh M., et al. (2022). Kinect-Based Rehabilitation Systems for Stroke Patients: A Scoping Review. *BioMed research international*, 2022, 4339054. <https://doi.org/10.1155/2022/4339054> PMID: 35386303
30. Sielużycki C., Maśliński J., Kaczmarczyk P., Kubacki R., Cieśliński W. B., & Witkowski K. (2019). Can Kinect aid motor learning in sportsmen? A study for three standing techniques in judo. *PloS one*, 14(2), e0210260. <https://doi.org/10.1371/journal.pone.0210260> PMID: 30726211
31. Abd-Alrazaq A., Al-Jafar E., Alajlani M., Toro C., Alhuwail D., Ahmed A., et al. (2022). The effectiveness of serious games for alleviating depression: systematic review and meta-analysis. *JMIR Serious Games*, 10(1), e32331. <https://doi.org/10.2196/32331> PMID: 35029530
32. Huang K., Zhao Y., He R., Zhong T., Yang H., Chen Y., et al. (2022). Exergame-based exercise training for depressive symptoms in adults: A systematic review and meta-analysis. *Psychology of Sport and Exercise*, 102266. <https://doi.org/10.1016/j.psychsport.2022.102266>
33. Rica R. L., Shimojo G. L., Gomes M. C., Alonso A. C., Pitta R. M., Santa-Rosa F. A., et al. (2020). Effects of a Kinect-based physical training program on body composition, functional fitness and depression in institutionalized older adults. *Geriatrics & gerontology international*, 20(3), 195–200. <https://doi.org/10.1111/ggi.13857> PMID: 31923924
34. W. Yunus F., Tan X. Z., & Romli M. H. (2020). Investigating the feasibility of exergame on sleep and emotion among university students. *Games for Health Journal*, 9(6), 415–424. <https://doi.org/10.1089/g4h.2019.0077> PMID: 33301386
35. Guillot A., Moschberger K., & Collet C. (2013). Coupling movement with imagery as a new perspective for motor imagery practice. *Behavioral and Brain Functions*, 9(1), 1–8. <https://doi.org/10.1186/1744-9081-9-8> PMID: 23425312
36. Matthews W. J., Benjamin C., Osborne C. (2007). Memory for moving and static images. *Psychonomic Bulletin & Review* 14, 989–993. <https://doi.org/10.3758/bf03194133> PMID: 18087971
37. Callow N., Roberts R., & Fawkes J. Z. (2006). Effects of dynamic and static imagery on vividness of imagery, skiing performance, and confidence. *Journal of Imagery Research in Sport and Physical Activity*, 1(1). <https://doi.org/10.2202/1932-0191.1001>
38. Guillot A., Di Rienzo F., Frank C., & Debarnot U., & MacIntyre T. (2021). From simulation to motor execution: a review of the impact of dynamic motor imagery on performance. *International Review of Sport and Exercise Psychology*, 1–20. <https://doi.org/10.1080/1750984X.2021.2007539>
39. Fusco A., Iasevoli L., Iosa M., Gallotta M. C., Padua L., Tucci L., et al. (2019). Dynamic motor imagery mentally simulates uncommon real locomotion better than static motor imagery both in young adults and elderly. *PloS one*, 14(6), e0218378. <https://doi.org/10.1371/journal.pone.0218378> PMID: 31242209
40. Jacobson N. S., Martell C. R., & Dimidjian S. (2001). Behavioral activation treatment for depression: returning to contextual roots. *Clinical Psychology: science and practice*, 8(3), 255. <https://doi.org/10.1093/clipsy.8.3.255>
41. Bortolato B., Miskowiak K. W., Köhler C. A., Maes M., Fernandes B. S., Berk M., et al. (2016). Cognitive remission: a novel objective for the treatment of major depression? *BMC Medicine*, 14, 9. <https://doi.org/10.1186/s12916-016-0560-3> PMID: 26801406
42. Chen J., Yang L., Zhang Z., Ma W., Xing-qu W., Zhang X., et al. (2013). The association between the disruption of motor imagery and the number of depressive episodes of major depression. *Journal of Affective Disorders*, 150(2), 337–343. <https://doi.org/10.1016/j.jad.2013.04.015> PMID: 23684121
43. Gorwood P., Richard-Devantoy S., Baylé F., & Cléry-Melun M. L. (2014). Psychomotor retardation is a scar of past depressive episodes, revealed by simple cognitive tests. *European Neuropsychopharmacology*, 24(10), 1630–1640. <https://doi.org/10.1016/j.euroneuro.2014.07.013> PMID: 25129432
44. Shadmehr R., Smith M. A., & Krakauer J. W. (2010). Error correction, sensory prediction, and adaptation in motor control. *Annual review of neuroscience*. <https://doi.org/10.1146/annurev-neuro-060909-153135> PMID: 20367317

45. Tiba A. I. & Manea L. (2018b). An Embodied Simulation Account of Cognition in Rational Emotive Behavior Therapy. *New Ideas in Psychology* 48 C pp. 12–20 <https://doi.org/10.1016/j.newideapsych.2017.08.003>
46. Tiba A. I., & Voss L. (2022). A motor imagery training for improving action cognition results in the reduction of residual symptoms after major depressive disorder: a single case study. *Journal of Evidence-Based Psychotherapies*, 22(1), 137–167. <https://doi.org/10.24193/jebp.2022.1.8>
47. Lambie J. A., & Marcel A. J. (2002). Consciousness and the Varieties of Emotion Experience A Theoretical Framework. *Psychological Review*, 109, 219–259. <https://doi.org/10.1037/0033-295x.109.2.219> PMID: 11990318
48. Holmes P. S., & Collins D. J. (2001). The PETTLEP approach to motor imagery: A functional equivalence model for sport psychologists. *Journal of applied sport psychology*, 13(1), 60–83. <https://doi.org/10.1080/10413200109339004>
49. Courtine G., Papaxanthis C., Gentili R., & Pozzo T. (2004). Gait-dependent motor memory facilitation in covert movement execution. *Cognitive Brain Research*, 22(1), 67–75. <https://doi.org/10.1016/j.cogbrainres.2004.07.008> PMID: 15561502
50. Malouin F., Jackson P. L., & Richards C. L. (2013). Towards the integration of mental practice in rehabilitation programs. A critical review. *Frontiers in human neuroscience*, 7, 576. <https://doi.org/10.3389/fnhum.2013.00576> PMID: 24065903
51. Martell C. R., Addis M. E., & Jacobson N. S. (2001). *Depression in context: Strategies for guided action*. WW Norton & Co.
52. Richards D. A., Rhodes S., Ekers D., McMillan D., Taylor R. S., Byford S., et al. (2017). Cost and Outcome of Behavioural Activation (COBRA): a randomised controlled trial of behavioural activation versus cognitive-behavioural therapy for depression. *Health Technology Assessment (Winchester, England)*, 21(46), 1–366. <https://doi.org/10.3310/hta21460> PMID: 28857042
53. Posner, K., Brent, D., Lucas, C., Gould, M., Stanley, B., Brown, G., et al. (2008). Columbia-suicide severity rating scale (C-SSRS). New York, NY: Columbia University Medical Center, 10. SUICIDAL BEHAVIOR (vtspc.org)
54. Beck A. T., Steer R. A., & Brown G. K. (1996). *BDI-II: Beck depression inventory*. Pearson.
55. Segal D. L., Coolidge F. L., Cahill B. S., & O'Riley A. A. (2008). Psychometric Properties of the Beck Depression Inventory—II (BDI-II) Among Community-Dwelling Older Adults. *Behavior modification*, 32(1), 3–20. <https://doi.org/10.1177/0145445507303833> PMID: 18096969
56. Beck A. T., Steer R. A., & Carbin M. G. (1988). Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review*, 8(1), 77–100. [https://doi.org/10.1016/0272-7358\(88\)90050-5](https://doi.org/10.1016/0272-7358(88)90050-5)
57. Kroenke K., Spitzer R. L., & Williams J. B. W. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16(9), 606–613. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x> PMID: 11556941
58. First M. B., Williams J. B., Karg R. S., & Spitzer R. L. (2016). User's guide for the SCID-5-CV Structured Clinical Interview for DSM-5® disorders: Clinical version. American Psychiatric Publishing, Inc.
59. Jacobson N. S., & Truax P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59(1), 12–19. <https://doi.org/10.1037//0022-006x.59.1.12> PMID: 2002127
60. Snaith R. P., Hamilton M., Morley S., Humayan A., Hargreaves D., & Trigwell P. (1995). A scale for the assessment of hedonic tone the Snaith–Hamilton Pleasure Scale. *The British Journal of Psychiatry*, 167(1), 99–103. <https://doi.org/10.1192/bjp.167.1.99> PMID: 7551619
61. Ang Y. S., Lockwood P., Apps M. A., Muhammed K., & Husain M. (2017). Distinct subtypes of apathy revealed by the apathy motivation index. *PloS one*, 12(1), e0169938. <https://doi.org/10.1371/journal.pone.0169938> PMID: 28076387
62. Spitzer R. L., Kroenke K., Williams J. B., & Löwe B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of internal medicine*, 166(10), 1092–1097. <https://doi.org/10.1001/archinte.166.10.1092> PMID: 16717171
63. Üstün, T. B., Kostanjsek, N., Chatterji, S., & Rehm, J. (Eds.). (2010). Measuring health and disability: Manual for WHO disability assessment schedule WHODAS 2.0. World Health Organization.
64. Rozental A., Kottorp A., Forsström D., Månsson K., Boettcher J., Andersson G., et al. (2019). The Negative Effects Questionnaire: Psychometric properties of an instrument for assessing adverse and unwanted events in psychological treatments. *Behavioural and Cognitive Psychotherapy*, 47(5), 559–572.
65. Dietrichkeit M., Hagemann-Goebel M., Nestoriuc Y., Moritz S., & Jelinek L. (2021). Side effects of the metacognitive training for depression compared to a cognitive remediation training in patients with

- depression. *Scientific Reports*, 11(1), 1–10. <https://doi.org/10.1038/s41598-021-87198-8> PMID: [33846503](https://pubmed.ncbi.nlm.nih.gov/33846503/)
66. Roberts R., Callow N., Hardy L., Markland D., & Bringer J. (2008). Movement imagery ability: development and assessment of a revised version of the vividness of movement imagery questionnaire. *Journal of Sport and Exercise Psychology*, 30(2), 200–221. <https://doi.org/10.1123/jsep.30.2.200> PMID: [18490791](https://pubmed.ncbi.nlm.nih.gov/18490791/)
 67. Ziv G., Lidor R., Arnon M., & Zeev A. (2017). The Vividness of Movement Imagery Questionnaire (VMIQ-2)—translation and reliability of a Hebrew version. *Israeli Journal of Psychiatry*, 54, 48–53.
 68. Armento M. E., & Hopko D. R. (2007). The environmental reward observation scale (EROS): development, validity, and reliability. *Behavior therapy*, 38(2), 107–119. <https://doi.org/10.1016/j.beth.2006.05.003> PMID: [17499078](https://pubmed.ncbi.nlm.nih.gov/17499078/)
 69. Treynor W., Gonzalez R., & Nolen-Hoeksema S. (2003). Rumination reconsidered: A psychometric analysis. *Cognitive therapy and research*, 27(3), 247–259. <https://doi.org/10.1023/A:1023910315561>
 70. Manos R. C., Kanter J. W., & Luo W. (2011). The behavioral activation for depression scale—short form: development and validation. *Behavior Therapy*, 42(4), 726–739. <http://dx.doi.org/10.1016/j.beth.2011.04.004> PMID: [22036000](https://pubmed.ncbi.nlm.nih.gov/22036000/)
 71. Iverson G. L., & Tulskey D. S. (2003). Detecting malingering on the WAIS-III: Unusual Digit Span performance patterns in the normal population and in clinical groups. *Archives of Clinical Neuropsychology*, 18(1), 1–9.
 72. Kazdin A. E. (2007). Mediators and mechanisms of change in psychotherapy research. *Annual Review Clinical Psychology*, 3, 1–27. <https://doi.org/10.1146/annurev.clinpsy.3.022806.091432> PMID: [17716046](https://pubmed.ncbi.nlm.nih.gov/17716046/)