

Analyzing the Ringelmann Effect with the Repeated Measures ANOVA

Nikolas Höft

Constantin Meyer-Grant

Joachim Munch

Quang Nguyen Duc

Frederik Schreck

Statistical Programming Languages

Humboldt-Universität zu Berlin




The Ringelmann Effect

- Maximilian Ringelmann (1861-1931):
 - ▶ French professor of agricultural engineering
- Work performance depends of number of group size
- Decreasing individual performance with increasing group size
- Example: Pulling weights with different sized groups




Overview

- The Ringelmann Effect can be investigated with an experimental design
 - ▶ Dependent Variable: Individual performance
 - ▶ Independent Variable / Factor: Group size
 - ▶ Realization of different factor levels
- For our purpose: Data simulation  Quantlet Data Simulation



The Ringelmann Effect

- The Ringelmann Effect can be investigated with an experimental design
 - ▶ Dependent Variable: Individual performance
 - ▶ Independent Variable / Factor: Group size
 - ▶ Realization of different factor levels
- For our purpose: Data simulation  Quantlet Data Simulation



Code

Analyzing the Ringelmann Effect with the Repeated Measures ANOVA



The Ringelmann Effect




Figure 1: Include a short, but meaningful caption.



The Repeated Measures ANOVA: The ANOVA model

- ANOVA: Analysis of Variance
- Comparison of the k factor level means

$$H_0 : \mu_1 = \mu_2 = \dots = \mu_k$$

$$H_1 : \exists i \neq j : \mu_i \neq \mu_j$$

- Test is accomplished by decomposition of variance components



Code

Analyzing the Ringelmann Effect with the Repeated Measures ANOVA



The Repeated Measures ANOVA: An Advantageous Model

- Problem: In case of large variance between different subjects
⇒ High error variance ⇒ Loss of power in F-Test
 - Repeated Measures ANOVA considers the between subject variance separately
⇒ Relatively low error variance ⇒ Gain of power in F-Test
- 🔍 Reduction of error variance



The Repeated Measures ANOVA: An Advantageous Model




Figure 2: Include a short, but meaningful caption.



The Repeated Measures ANOVA: An Advantageous Model

- Design Requirement: Each subject has to be measured under all factor levels

Title	Title
2.13	1.45
3.14	6.85

Table 1: Example Data Matrix



Code

Analyzing the Ringelmann Effect with the Repeated Measures ANOVA



Table



The Repeated Measures ANOVA: Confidence Intervals

- The computation of the confidence intervals has to be adjusted in the Repeated Measures ANOVA



Confidence Intervals



Code

Analyzing the Ringelmann Effect with the Repeated Measures ANOVA



The Repeated Measures ANOVA: Confidence Intervals




Figure 3: Include a short, but meaningful caption.



The Repeated Measures ANOVA: Effect Size Measures

□ Two measures of effect size:

▶ η^2

▶ η_p^2



Code

Analyzing the Ringelmann Effect with the Repeated Measures ANOVA



Table



An Important Requirement

- Sphericity: The variance of differences are equal for each pair of factor levels
- Test for sphericity: Mauchly test
- Measurement of sphericity ($\epsilon \in [0, 1]$):
 - ▶ Greenhouse & Geisser: ϵ_{GG}
 - ▶ Box: ϵ_B
 - ▶ Huynh & Feldt: ϵ_{HF}
- These can be used to correct the degrees of freedom and therefore adjust the p-values if sphericity is violated

