

# Totally geodesic submanifolds of the homogeneous nearly Kähler six-manifolds

Juan Manuel Lorenzo Naveiro

CITMAGA – Universidade de Santiago de Compostela

Geometry day, 11 April 2024

Joint work with Alberto Rodríguez-Vázquez




CENTRO DE INVESTIGACIÓN  
E TECNOLOXÍA MATEMÁTICA  
DE GALICIA




# Totally geodesic submanifolds


- $\Sigma, M$  complete Riemannian manifolds,  $f: \Sigma \rightarrow M$  isometric immersion.
- $f$  is *totally geodesic* if:
  - $\gamma(t)$  geodesic in  $\Sigma \Rightarrow f(\gamma(t))$  geodesic in  $M$ .
  - $\text{II} = 0$ .



$$\mathbb{R}^k \subseteq \mathbb{R}^n$$



$$S^k \subseteq S^n$$



$$\mathbb{R}H^k \subseteq \mathbb{R}H^n$$

# Totally geodesic submanifolds

- $\Sigma, M$  complete Riemannian manifolds,  $f: \Sigma \rightarrow M$  isometric immersion.
- $f$  is *totally geodesic* if:
  - $\gamma(t)$  geodesic in  $\Sigma \Rightarrow f(\gamma(t))$  geodesic in  $M$ .
  - $\text{II} = 0$ .

## General problem

Classify totally geodesic submanifolds of  $M$  up to congruence.

# Nearly Kähler geometry

- $(M, J)$  is nearly Kähler if  $(\nabla_X J)X = 0$  for all  $X \in \mathfrak{X}(M)$ .
- Butruille ('06): classification of homogeneous NK 6-manifolds:

$$\mathbb{S}^6 = \frac{G_2}{SU(3)}$$

$$\mathbb{CP}^3 = \frac{Sp(2)}{U(1) \times Sp(1)}$$

$$F(\mathbb{C}^3) = \frac{SU(3)}{T^2}$$

$$\mathbb{S}^3 \times \mathbb{S}^3 = \frac{SU(2) \times SU(2) \times SU(2)}{\Delta SU(2)}$$

$$\begin{array}{c} \sigma \in \text{Aut}(G) \\ \sigma^3 = \text{Id} \end{array} \longrightarrow \sigma_* = \frac{1}{2}\text{Id} + \frac{\sqrt{3}}{2}J$$

# Previously known results


- $F(\mathbb{C}^3)$ :
  - Totally geodesic + Lagrangian (Storm '20).
  - Totally geodesic + almost complex + 2 – dim (Vrancken, Cwiklinski '22).
- $\mathbb{CP}^3$ :
  - Totally geodesic + Lagrangian (Aslan '23, Liefsoens '22).
- $\mathbb{S}^3 \times \mathbb{S}^3$ :
  - Totally geodesic + Lagrangian (Zhang, Dioos, Hu, Vrancken, Wang '16).
  - Totally geodesic + almost complex + 2 – dim (Bolton, Dillen, Dioos, Vrancken '22).

# Reductive homogeneous spaces

- $M = G/K$  homogeneous space,  $o = eK$ .
- $G \curvearrowright M$  gives a homomorphism  $G \rightarrow I(M)$ .
- $X \in \mathfrak{g} \mapsto X^* \in \mathfrak{X}(M)$ .

$$X_p^* = \left. \frac{d}{dt} \right|_{t=0} \text{Exp}(tX) \cdot p$$

- $T_o M = \mathfrak{g}/\mathfrak{k}$  via  $X + \mathfrak{k} \mapsto X_o^*$ .



$M = G/K$  is *reductive* if there is a  $\mathfrak{p} \subseteq \mathfrak{g}$  such that  
 $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ ,  $\text{Ad}(K)\mathfrak{p} = \mathfrak{p}$ .

- $\mathfrak{p} = \mathfrak{g}/\mathfrak{k} = T_o M$ .
- Isotropy representation  $K \curvearrowright T_o M \Leftrightarrow$  Adjoint representation  $K \curvearrowright \mathfrak{p}$ .
- $G$ -invariant tensor fields on  $M \Leftrightarrow \text{Ad}(K)$ -invariant tensors on  $\mathfrak{p}$ .

$$\langle \cdot, \cdot \rangle \text{ } G\text{-invariant metric} = \langle \cdot, \cdot \rangle \text{ } \text{Ad}(K)\text{-invariant metric on } \mathfrak{p}$$

- $\mathfrak{p} = \mathfrak{g}/\mathfrak{k} = T_o M$ .
- Isotropy representation  $K \curvearrowright T_o M \rightleftarrows$  Adjoint representation  $K \curvearrowright \mathfrak{p}$ .
- $G$ -invariant tensor fields on  $M \rightleftarrows \text{Ad}(K)$ -invariant tensors on  $\mathfrak{p}$ .

### Theorem (Nomizu '54)

$$\begin{array}{ccc} \text{Invariant connections} & & \text{Ad-equivariant maps} \\ \nabla: \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M) & = & \alpha: \mathfrak{p} \otimes \mathfrak{p} \rightarrow \mathfrak{p} \end{array}$$

$$\alpha(X, Y) = \Lambda(X)Y = \nabla_{X^*} Y^* + [X, Y]_{\mathfrak{p}},$$

$$R(X, Y) = [\Lambda(X), \Lambda(Y)] - \Lambda([X, Y]_{\mathfrak{p}}) - \text{ad}([X, Y]_{\mathfrak{p}}),$$

$$T(X, Y) = \alpha(X, Y) - \alpha(Y, X) - [X, Y]_{\mathfrak{p}}.$$



# Natural reductivity

$$\alpha(X, Y) = \frac{1}{2} [X, Y]_{\mathfrak{p}} + U(X, Y),$$

$$2\langle U(X, Y), Z \rangle = \langle [Z, X]_{\mathfrak{p}}, Y \rangle + \langle X, [Z, Y]_{\mathfrak{p}} \rangle.$$

- $M = G/K$  is *naturally reductive* if  $U = 0$ .  
 $\Leftrightarrow \exp_o(tX) = \text{Exp}(tX) \cdot o$  for all  $X \in \mathfrak{p}$ .
- $M = G/K$  is *normal homogeneous* if:
  1.  $G$  has a bi-invariant metric.
  2. The complement  $\mathfrak{p} = \mathfrak{g} \ominus \mathfrak{k}$ .
  3. The metric on  $M$  is induced from the metric on  $G$ .

# Totally geodesic submanifolds of $G/K$

- $\Sigma \subseteq M$  totally geodesic,  $p \in \Sigma$ . Then,

$$\Sigma = \exp_p(T_p\Sigma).$$

- If  $V \subseteq T_pM$ , when is  $\exp_p(V)$  totally geodesic?

## Theorem (Cartan '51, Hermann '59)

$M$  analytic Riemannian manifold,  $V \subseteq T_pM$ . The following are equivalent:

- $V$  generates a totally geodesic submanifold.
- $V$  is  $\nabla^k R$ -invariant for all  $k = 0, 1, 2, \dots$

# Totally geodesic submanifolds of $G/K$

- $\Sigma \subseteq M$  totally geodesic,  $p \in \Sigma$ . Then,

$$\Sigma = \exp_p(T_p\Sigma).$$

- If  $V \subseteq T_pM$ , when is  $\exp_p(V)$  totally geodesic?

## Theorem (Cartan) for naturally reductive spaces

$M$  naturally reductive,  $\mathfrak{v} \subseteq \mathfrak{p}$ . The following are equivalent:

- $\mathfrak{v}$  generates a totally geodesic submanifold.
- $\mathfrak{v}$  is  $\nabla^k R$ -invariant for all  $k = 0, 1, 2, \dots, d$ .

## Tojo's criterion ('96)

$M$  naturally reductive,  $\mathfrak{v} \subseteq \mathfrak{p}$ . The following are equivalent:

- $\mathfrak{v}$  generates a totally geodesic submanifold.
- $R(e^{-\Lambda(X)}\mathfrak{v}, e^{-\Lambda(X)}\mathfrak{v})e^{-\Lambda(X)}\mathfrak{v} \subseteq e^{-\Lambda(X)}\mathfrak{v}$  for all  $X \in \mathfrak{v}$ .

## Sufficient criterion (LN—Rodríguez-Vázquez)

$M$  naturally reductive,  $\mathfrak{v} \subseteq \mathfrak{p}$  invariant under  $R$  and  $\alpha$ :

1. The subalgebra  $\mathfrak{s} = [\mathfrak{v}, \mathfrak{v}] + \mathfrak{v} = [\mathfrak{v}, \mathfrak{v}]_{\mathfrak{k}} \oplus \mathfrak{v}$  satisfies

$$\mathfrak{s} = (\mathfrak{s} \cap \mathfrak{k}) \oplus (\mathfrak{s} \cap \mathfrak{p}) = \mathfrak{s}_{\mathfrak{k}} \oplus \mathfrak{s}_{\mathfrak{p}}.$$

2. The set  $\exp_o(\mathfrak{v}) = S \cdot o$  is totally geodesic.


## Sufficient criterion (LN—Rodríguez-Vázquez)

$M$  naturally reductive,  $\mathfrak{v} \subseteq \mathfrak{p}$  invariant under  $R$  and  $\alpha$ :

1. The subalgebra  $\mathfrak{s} = [\mathfrak{v}, \mathfrak{v}] + \mathfrak{v} = [\mathfrak{v}, \mathfrak{v}]_{\mathfrak{k}} \oplus \mathfrak{v}$  satisfies

$$\mathfrak{s} = (\mathfrak{s} \cap \mathfrak{k}) \oplus (\mathfrak{s} \cap \mathfrak{p}) = \mathfrak{s}_{\mathfrak{k}} \oplus \mathfrak{s}_{\mathfrak{p}}.$$

2. The set  $\exp_o(\mathfrak{v}) = S \cdot o$  is totally geodesic.



# The flag manifold $F(\mathbb{C}^3)$

- $F(\mathbb{C}^3) = \frac{\mathrm{SU}(3)}{\mathrm{T}^2}; I^0(F(\mathbb{C}^3)) = \mathrm{PSU}(3).$

- $\mathfrak{p} = \left\{ \begin{pmatrix} 0 & x & z \\ -\bar{x} & 0 & y \\ -\bar{z} & -\bar{y} & 0 \end{pmatrix} : x, y, z \in \mathbb{C} \right\} = \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3.$

$$\mathrm{diag}(e^{i\theta_1}, e^{i\theta_2}, e^{-i(\theta_1+\theta_2)}) \bullet \begin{matrix} \nearrow e^{i(\theta_1-\theta_2)} \\ \rightarrow e^{i(\theta_1+2\theta_2)} \\ \searrow e^{i(2\theta_1+\theta_2)} \end{matrix}$$

# The flag manifold $F(\mathbb{C}^3)$

- $F(\mathbb{C}^3) = \frac{\mathrm{SU}(3)}{\mathrm{T}^2}; I^0(F(\mathbb{C}^3)) = \mathrm{PSU}(3).$
- $\mathfrak{p} = \left\{ \begin{pmatrix} 0 & x & z \\ -\bar{x} & 0 & y \\ -\bar{z} & -\bar{y} & 0 \end{pmatrix} : x, y, z \in \mathbb{C} \right\} = \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3.$
- $\langle X, Y \rangle = -\mathrm{tr}(XY)$

# The flag manifold $F(\mathbb{C}^3)$

- $F(\mathbb{C}^3) = \frac{SU(3)}{T^2}; I^0(F(\mathbb{C}^3)) = PSU(3).$

General fact:

$K \subseteq H \subseteq G$  gives a fibration  
 $H/K \rightarrow G/K \rightarrow G/H.$

- $\mathfrak{p} = \left\{ \begin{pmatrix} 0 & x & z \\ -\bar{x} & 0 & y \\ -\bar{z} & -\bar{y} & 0 \end{pmatrix} : x, y, z \in \mathbb{C} \right\} = \mathfrak{p}_1 \oplus \mathfrak{p}_2 \oplus \mathfrak{p}_3.$

- $T^2 \subseteq U(2) \subseteq SU(3)$  gives the fibration  $\mathbb{CP}^1 \rightarrow F(\mathbb{C}^3) \rightarrow \mathbb{CP}^2.$

- $\mathcal{V}_o = \mathfrak{p}_1, \mathcal{H}_o = \mathfrak{p}_2 \oplus \mathfrak{p}_3.$



# The complex projective space $\mathbb{CP}^3$

- $\mathbb{CP}^3 = \frac{\mathrm{Sp}(2)}{\mathrm{U}(1) \times \mathrm{Sp}(1)}; I^0(\mathbb{CP}^3) = \frac{\mathrm{Sp}(2)}{\mathbb{Z}^2}.$
- $\mathfrak{p} = \left\{ \begin{pmatrix} \textcolor{brown}{z}^j & -\textcolor{violet}{\bar{q}} \\ \textcolor{violet}{q} & 0 \end{pmatrix} : z \in \mathbb{C}, q \in \mathbb{H} \right\} = \textcolor{brown}{\mathfrak{p}}_1 \oplus \textcolor{violet}{\mathfrak{p}}_2.$

$$\begin{aligned} (\lambda, \mu) \cdot \textcolor{brown}{z} &= \textcolor{brown}{\lambda}^2 \textcolor{brown}{z} \\ (\lambda, \mu) \cdot \textcolor{violet}{q} &= \textcolor{violet}{\mu} \textcolor{violet}{q} \bar{\lambda} \end{aligned}$$

# The complex projective space $\mathbb{CP}^3$

- $\mathbb{CP}^3 = \frac{\mathrm{Sp}(2)}{\mathrm{U}(1) \times \mathrm{Sp}(1)}; I^0(\mathbb{CP}^3) = \frac{\mathrm{Sp}(2)}{\mathbb{Z}^2}.$
- $\mathfrak{p} = \left\{ \begin{pmatrix} z^j & -\bar{q} \\ q & 0 \end{pmatrix} : z \in \mathbb{C}, q \in \mathbb{H} \right\} = \mathfrak{p}_1 \oplus \mathfrak{p}_2.$
- $\langle X, Y \rangle = -2\mathrm{Re} \, \mathrm{tr}_{\mathbb{H}}(XY).$

# The complex projective space $\mathbb{CP}^3$

- $\mathbb{CP}^3 = \frac{\mathrm{Sp}(2)}{\mathrm{U}(1) \times \mathrm{Sp}(1)}; I^0(\mathbb{CP}^3) = \frac{\mathrm{Sp}(2)}{\mathbb{Z}^2}.$
- $\mathfrak{p} = \left\{ \begin{pmatrix} \textcolor{brown}{z}j & -\textcolor{violet}{\bar{q}} \\ \textcolor{violet}{q} & 0 \end{pmatrix} : z \in \mathbb{C}, q \in \mathbb{H} \right\} = \textcolor{brown}{\mathfrak{p}}_1 \oplus \textcolor{violet}{\mathfrak{p}}_2.$
- $\mathrm{U}(1) \times \mathrm{Sp}(1) \subseteq \mathrm{Sp}(1) \times \mathrm{Sp}(1) \subseteq \mathrm{Sp}(2)$  gives the fibration  $\mathbb{CP}^1 \rightarrow \mathbb{CP}^3 \rightarrow \mathbb{HP}^1 = \mathbb{S}^4.$
- $\mathcal{V}_o = \mathfrak{p}_1, \mathcal{H}_o = \mathfrak{p}_2.$

# The almost product $\mathbb{S}^3 \times \mathbb{S}^3$

- $\mathbb{S}^3 \times \mathbb{S}^3 = \frac{\mathrm{SU}(2)^3}{\Delta \mathrm{SU}(2)}; I^0(\mathbb{S}^3 \times \mathbb{S}^3) = \frac{\mathrm{SU}(2)^3}{\Delta \mathbb{Z}_2}.$
- $\mathfrak{p} = \{(X, Y, Z): X + Y + Z = 0\} = \mathfrak{p}_1 \oplus \mathfrak{p}_2.$ 
  - $\mathfrak{p}_1 = \{(X, -2X, X): X \in \mathfrak{su}(2)\} \cong \mathrm{Ad}.$
  - $\mathfrak{p}_2 = \{(X, 0, -X): X \in \mathfrak{su}(2)\} \cong \mathrm{Ad}.$
- $\langle (X, Y, Z), (X', Y', Z') \rangle = -\mathrm{tr}(XX') - \mathrm{tr}(YY') - \mathrm{tr}(ZZ').$

# The almost product $\mathbb{S}^3 \times \mathbb{S}^3$

- $\mathbb{S}^3 \times \mathbb{S}^3 = \frac{SU(2)^3}{\Delta SU(2)}$ ;  $I^0(\mathbb{S}^3 \times \mathbb{S}^3) = \frac{SU(2)^3}{\Delta \mathbb{Z}_2}$ .
- $\mathfrak{p} = \{(X, Y, Z): X + Y + Z = 0\} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$ .
- $\Delta SU(2) \subseteq \Delta_{13} SU(2) \times SU(2)_2 \subseteq SU(2)^3$  gives the fibration  $\mathbb{S}^3 \rightarrow \mathbb{S}^3 \times \mathbb{S}^3 \rightarrow \mathbb{S}^3$ .
- $\mathcal{V}_o = \mathfrak{p}_1$ ,  $\mathcal{H}_o = \mathfrak{p}_2$ .

# Low codimension

## Theorem (Nikolayevsky '15)

$M$  simply connected, compact and homogeneous.  
If there exists  $\Sigma \subseteq M$  totally geodesic hypersurface, then  $M = \mathbb{S}^n$ .

## Corollary

$M \in \{F(\mathbb{C}^3), \mathbb{CP}^3, \mathbb{S}^3 \times \mathbb{S}^3\}$  has no totally geodesic hypersurfaces.

# Low codimension

## Theorem (Nikolayevsky '15)

$M$  simply connected, compact and homogeneous.  
If there exists  $\Sigma \subseteq M$  totally geodesic hypersurface, then  $M = \mathbb{S}^n$ .

## Theorem

$M \in \{F(\mathbb{C}^3), \mathbb{CP}^3, \mathbb{S}^3 \times \mathbb{S}^3\}$  has no totally geodesic submanifolds of codimension  $\leq 2$ .

# Dimension three

Ambient	Submanifold	Orbit of
$F(\mathbb{C}^3)$	$F(\mathbb{R}^3) = \mathbb{S}^3/Q_8$	$SO(3)$
	$\mathbb{S}_{\mathbb{C}}^3$	$SU(2)$
$\mathbb{CP}^3$	$\mathbb{RP}_{\mathbb{C}}^3$	$SU(2)^j$
$\mathbb{S}^3 \times \mathbb{S}^3$	$\mathbb{S}^3(2/\sqrt{3})$	$SU(2)_2$
	$\mathbb{S}_{\mathbb{C}}^3$	$\Delta_{13}SU(2) \times SU(2)_2$

All of these are Lagrangian.




# Dimension two in $F(\mathbb{C}^3)$

Submanifold	Orbit of	Relationship with $J$
$T_A = \mathbb{R}^2/A$	$T^2$	Almost complex
$\mathbb{CP}^1 = \mathbb{S}^2(1/\sqrt{2})$	$U(2)$	Almost complex
$\mathbb{S}^2(\sqrt{2})$	$SO(3)$	Almost complex
$\mathbb{RP}^2 \rightarrow F(\mathbb{R}^3)$	Inhomogeneous	Totally real

# Dimension two in $F(\mathbb{C}^3)$



Submanifold	Orbit of	Relationship with $J$
$T_A = \mathbb{R}^2/A$	$T^2$	Almost complex



$$A = \mathbb{Z} - \text{span} \left\{ \left( 0, \frac{2\sqrt{2}\pi}{\sqrt{3}} \right), \left( \sqrt{2}\pi, \frac{\sqrt{2}}{\sqrt{3}}\pi \right) \right\}$$

# Dimension two in $F(\mathbb{C}^3)$

Submanifold	Orbit of	Relationship with $J$
$T_A = \mathbb{R}^2/A$	$T^2$	Almost complex



# Dimension two in $\mathbb{CP}^3$

Submanifold	Orbit of	Relationship with $J$
$\mathbb{S}^2(1/\sqrt{2})$	$\mathrm{Sp}(1)_f$	Almost complex
$\mathbb{S}^2(1)$	$\mathrm{SU}(2)$	Almost complex
$\mathbb{S}^2(\sqrt{5})$	$\mathrm{SU}(2)_{\Lambda_3}$	Almost complex

$\Lambda_3 = \mathrm{Sym}^3(\mathbb{C}^2)$ : three-dimensional irrep. of  $\mathrm{SU}(2)$ .


$$\begin{pmatrix} a & -\bar{b} \\ b & \bar{a} \end{pmatrix} \mapsto \begin{pmatrix} a(|a|^2 - 2|b|^2) & -\sqrt{3}a^2\bar{b} \\ \sqrt{3}a^2\bar{b} & a^3 \end{pmatrix} + j \begin{pmatrix} b(2|a|^2 - |b|^2) & -\sqrt{3}ab^2 \\ \sqrt{3}\bar{a}b^2 & -b^3 \end{pmatrix}$$

# Dimension two in $\mathbb{S}^3 \times \mathbb{S}^3$

Submanifold	Orbit of	Relationship with $J$
$T_B = \mathbb{R}^2/B$	$T \subseteq U(1)^3$	Almost complex
$\mathbb{S}^2(\sqrt{3}/\sqrt{2})$	$\Delta SU(2)$	Almost complex
$\mathbb{S}^2 \subseteq \mathbb{S}^3(2/\sqrt{3})$	$H \subseteq \Delta_{13} SU(2) \times SU(2)_2$	Totally real

# Dimension two in $\mathbb{S}^3 \times \mathbb{S}^3$



Submanifold	Orbit of	Relationship with $J$
$T_B = \mathbb{R}^2/B$	$T \subseteq U(1)^3$	Almost complex



$$B = \mathbb{Z} - \text{span} \left\{ \left( \frac{4\pi}{\sqrt{3}}, 0 \right), \left( \frac{2\pi}{\sqrt{3}}, 2\pi \right) \right\}$$

# Dimension two in $\mathbb{S}^3 \times \mathbb{S}^3$

Submanifold	Orbit of	Relationship with $J$
$T_B = \mathbb{R}^2/B$	$T \subseteq U(1)^3$	Almost complex



Ambient	Submanifold	Orbit of	Relationship with $J$
$F(\mathbb{C}^3)$	$F(\mathbb{R}^3)$	$SO(3)$	Lagrangian
	$\mathbb{S}^3_{\mathbb{C}}$	$SU(2)$	Lagrangian
	$T_A$	$T^2$	Almost complex
	$\mathbb{CP}^1 = \mathbb{S}^2(1/\sqrt{2})$	$U(2)$	Almost complex
	$\mathbb{S}^2(\sqrt{2})$	$SO(3)$	Almost complex
	$\mathbb{RP}^2 \rightarrow F(\mathbb{R}^3)$	Inhomogeneous	Totally real
$\mathbb{CP}^3$	$\mathbb{RP}^3_{\mathbb{C}}$	$SU(2)^j$	Lagrangian
	$\mathbb{S}^2(1/\sqrt{2})$	$Sp(1)_f$	Almost complex
	$\mathbb{S}^2(1)$	$SU(2)$	Almost complex
	$\mathbb{S}^2(\sqrt{5})$	$SU(2)_{\Lambda_3}$	Almost complex
$\mathbb{S}^3 \times \mathbb{S}^3$	$\mathbb{S}^3(2/\sqrt{3})$	$SU(2)_2$	Lagrangian
	$\mathbb{S}^3_{\mathbb{C}}$	$\Delta_{13}SU(2) \times SU(2)_2$	Lagrangian
	$T_B$	$T \subseteq U(1)^3$	Almost complex
	$\mathbb{S}^2(\sqrt{3}/\sqrt{2})$	$\Delta SU(2)$	Almost complex
	$\mathbb{S}^2 \subseteq \mathbb{S}^3(2/\sqrt{3})$	$H \subseteq \Delta_{13}SU(2) \times SU(2)_2$	Totally real





*That's all Folks!*

Ambient	Submanifold	Orbit of	Relationship with $J$
$F(\mathbb{C}^3)$	$F(\mathbb{R}^3)$	$SO(3)$	Lagrangian
	$\mathbb{S}^3_{\mathbb{C}}$	$SU(2)$	Lagrangian
	$T_A$	$T^2$	Almost complex
	$\mathbb{CP}^1 = \mathbb{S}^2(1/\sqrt{2})$	$U(2)$	Almost complex
	$\mathbb{S}^2(\sqrt{2})$	$SO(3)$	Almost complex
	$\mathbb{RP}^2 \rightarrow F(\mathbb{R}^3)$	Inhomogeneous	Totally real
$\mathbb{CP}^3$	$\mathbb{RP}^3_{\mathbb{C}}$	$SU(2)^j$	Lagrangian
	$\mathbb{S}^2(1/\sqrt{2})$	$Sp(1)_f$	Almost complex
	$\mathbb{S}^2(1)$	$SU(2)$	Almost complex
	$\mathbb{S}^2(\sqrt{5})$	$SU(2)_{\Lambda_3}$	Almost complex
$\mathbb{S}^3 \times \mathbb{S}^3$	$\mathbb{S}^3(2/\sqrt{3})$	$SU(2)_2$	Lagrangian
	$\mathbb{S}^3_{\mathbb{C}}$	$\Delta_{13}SU(2) \times SU(2)_2$	Lagrangian
	$T_B$	$T \subseteq U(1)^3$	Almost complex
	$\mathbb{S}^2(\sqrt{3}/\sqrt{2})$	$\Delta SU(2)$	Almost complex
	$\mathbb{S}^2 \subseteq \mathbb{S}^3(2/\sqrt{3})$	$H \subseteq \Delta_{13}SU(2) \times SU(2)_2$	Totally real