

Polar homogeneous foliations on hyperbolic spaces

Juan Manuel Lorenzo Naveiro
University of Oklahoma
Symmetry and Shape, 05/11/2025

Joint work with J. C. Díaz-Ramos
(Universidade de Santiago de Compostela)

Polar actions

M complete Riemannian manifold, $H \curvearrowright M$ proper isometric action.

Polar actions

M complete Riemannian manifold, $H \curvearrowright M$ proper isometric action.


$H \curvearrowright M$ is *polar* if $\exists \Sigma \subseteq M$ with:

- $\Sigma \cap (H \cdot p) \neq \emptyset$ for all $p \in M$,
- $T_p \Sigma \perp T_p(H \cdot p)$ for all $p \in \Sigma$.


Σ is a *section*.




Examples on $\mathbb{RH}^2 = \mathrm{SL}(2, \mathbb{R})/\mathrm{SO}(2)$



K



A



N

Two actions $H_1 \curvearrowright M$, $H_2 \curvearrowright M$ are *orbit equivalent* if $\exists f \in I(M)$ with $f(H_1 \cdot p) = H_2 \cdot f(p)$ for all $p \in M$.

Main problem. Given M , classify polar actions on M up to orbit equivalence.

Known results

- Polar representations: Dadok ('85).

Every polar representation is orbit equivalent to the *isotropy representation* of a symmetric space.

Known results

- Polar representations: Dadok ('85).

Every polar representation is orbit equivalent to the *isotropy representation* of a symmetric space.

- $\mathbb{R}H^n$: Wu ('92).

Known results

- Polar representations: Dadok ('85).

Every polar representation is orbit equivalent to the *isotropy representation* of a symmetric space.

- $\mathbb{R}H^n$: Wu ('92).
- Compact rank one symmetric spaces: Podestà, Thorbergsson ('99).

Known results

- Polar representations: Dadok ('85).

Every polar representation is orbit equivalent to the *isotropy representation* of a symmetric space.

- $\mathbb{R}H^n$: Wu ('92).
- Compact rank one symmetric spaces: Podestà, Thorbergsson ('99).
- Compact type, higher rank: Kollross, Lytchak.

The noncompact setting

- $\mathbb{C}H^n$: Díaz-Ramos, Domínguez-Vázquez, Kollross ('17).

The noncompact setting

- $\mathbb{C}H^n$: Díaz-Ramos, Domínguez-Vázquez, Kollross ('17).
- Cohomogeneity one actions: Berndt, Brück, Díaz-Ramos, Domínguez-Vázquez, Otero, Rodríguez-Vázquez, Sanmartín-López, Solonenko, Tamaru ('01-'25).

The noncompact setting

- $\mathbb{C}H^n$: Díaz-Ramos, Domínguez-Vázquez, Kollross ('17).
- Cohomogeneity one actions: Berndt, Brück, Díaz-Ramos, Domínguez-Vázquez, Otero, Rodríguez-Vázquez, Sanmartín-López, Solonenko, Tamaru ('01-'25).
- Hyperpolar homogeneous foliations: Berndt, Díaz-Ramos, Tamaru ('10).

The noncompact setting

- $\mathbb{C}H^n$: Díaz-Ramos, Domínguez-Vázquez, Kollross ('17).
- Cohomogeneity one actions: Berndt, Brück, Díaz-Ramos, Domínguez-Vázquez, Otero, Rodríguez-Vázquez, Sanmartín-López, Solonenko, Tamaru ('01-'25).
- Hyperpolar homogeneous foliations: Berndt, Díaz-Ramos, Tamaru ('10).
- Codimension two polar homogeneous foliations: Díaz-Ramos, LN ('25).

Symmetric spaces of noncompact type

- $M = G/K$ irreducible symmetric space of noncompact type.

Symmetric spaces of noncompact type

- $M = G/K$ irreducible symmetric space of noncompact type.
- $\theta \in \text{Aut}(\mathfrak{g})$ *Cartan involution* $\leadsto \mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ *Cartan decomposition*.

Symmetric spaces of noncompact type

- $M = G/K$ irreducible symmetric space of noncompact type.
- $\theta \in \text{Aut}(\mathfrak{g})$ *Cartan involution* $\leadsto \mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ *Cartan decomposition*.
- $\langle X, Y \rangle = -\mathcal{B}(X, \theta Y)$ inner product on \mathfrak{g} .

Symmetric spaces of noncompact type

- $M = G/K$ irreducible symmetric space of noncompact type.
- $\theta \in \text{Aut}(\mathfrak{g})$ *Cartan involution* $\leadsto \mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ *Cartan decomposition*.
- $\langle X, Y \rangle = -\mathcal{B}(X, \theta Y)$ inner product on \mathfrak{g} .

Iwasawa decomposition:

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}, \quad G = KAN.$$



Noncompact rank one symmetric spaces

- $M = G/K = \mathbb{F}H^n$ symmetric space of noncompact type and rank one.
- $\mathfrak{a} = \mathbb{R}$ and $\mathfrak{n} = \mathfrak{g}_\alpha \oplus \mathfrak{g}_{2\alpha}$. Moreover, $-1 \leq \sec \leq -\frac{1}{4}$.

M	G	K	\mathfrak{g}_α	$\mathfrak{g}_{2\alpha}$
$\mathbb{R}H^n$	$SO^0(1, n)$	$SO(n)$	\mathbb{R}^{n-1}	0
$\mathbb{C}H^n$	$SU(1, n)$	$S(U(1) \times U(n))$	\mathbb{C}^{n-1}	\mathbb{R}
$\mathbb{H}H^n$	$Sp(1, n)$	$Sp(1) \times Sp(n-1)$	\mathbb{H}^{n-1}	\mathbb{R}^3
$\mathbb{O}H^2$	F_4^{-20}	$Spin(9)$	\mathbb{O}	\mathbb{R}^7



Nilpotent foliation $\mathcal{F}_{\alpha, \mathfrak{v}}$

$$\mathfrak{s}_{\alpha, \mathfrak{v}} = \mathfrak{n} \ominus \mathfrak{v}, \quad \mathfrak{v} \subseteq \mathfrak{g}_{\alpha} \text{ abelian subspace.}$$



Solvable foliation $\mathcal{F}_{0,\mathfrak{v}}$

$$\mathfrak{s}_{0,\mathfrak{v}} = \mathfrak{a} \oplus (\mathfrak{n} \ominus \mathfrak{v}), \quad \mathfrak{v} \subseteq \mathfrak{g}_\alpha \text{ abelian subspace.}$$



Theorem (Díaz-Ramos, LN). The hyperbolic planes $M = \mathbb{H}^2$ and $\mathbb{O}H^2$ admit exactly four (nontrivial) polar homogeneous foliations up to orbit equivalence:

$$AN \curvearrowright M, \quad N = S_{a,0} \curvearrowright M, \quad S_{0,\ell} \curvearrowright M, \quad S_{a,\ell} \curvearrowright M.$$

Remark. $\dim \ell = 1$.

Sections of polar actions $S \curvearrowright \mathbb{H}^n$

- $S \curvearrowright \mathbb{O}H^2$ polar action with singular orbits and cohomogeneity ≥ 2
 \leadsto The section is $\mathbb{R}H^k(c)$, where $c \in \{-\frac{1}{4}, -1\}$ (Kollross '20).

Sections of polar actions $S \curvearrowright \mathbb{H}^n$

- $S \curvearrowright \mathbb{O}H^2$ polar action with singular orbits and cohomogeneity ≥ 2
 \leadsto The section is $\mathbb{R}H^k(c)$, where $c \in \{-\frac{1}{4}, -1\}$ (Kollross '20).
- Kollross' argument also holds for \mathbb{H}^n .

Sections of polar actions $S \curvearrowright \mathbb{H}^n$

- $S \curvearrowright \mathbb{O}H^2$ polar action with singular orbits and cohomogeneity ≥ 2
 \leadsto The section is $\mathbb{R}H^k(c)$, where $c \in \{-\frac{1}{4}, -1\}$ (Kollross '20).
- Kollross' argument also holds for $\mathbb{H}H^n$.

Theorem (Díaz-Ramos, LN). If $S \curvearrowright \mathbb{H}H^n$ is a nontrivial polar action with cohomogeneity ≥ 2 , then its section is a totally geodesic $\mathbb{R}H^k$ with constant curvature $c \in \{-1, -\frac{1}{4}\}$.

Standard polar foliations on $\mathbb{H}H^n$

\mathcal{F} homogeneous foliation on M .

\mathcal{F} is *standard* if it is induced by some $S \subseteq AN$ up to orbit equivalence.

Note. $S_{\mathfrak{b}, \mathfrak{v}} \curvearrowright \mathbb{H}H^n$ is standard, where $\mathfrak{s}_{\mathfrak{b}, \mathfrak{v}} = (\mathfrak{a} \ominus \mathfrak{b}) \oplus (\mathfrak{n} \ominus \mathfrak{v})$.

Standard polar foliations on $\mathbb{H}H^n$

\mathcal{F} homogeneous foliation on M .

\mathcal{F} is *standard* if it is induced by some $S \subseteq AN$ up to orbit equivalence.

Note. $S_{\mathfrak{b}, \mathfrak{v}} \curvearrowright \mathbb{H}H^n$ is standard, where $\mathfrak{s}_{\mathfrak{b}, \mathfrak{v}} = (\mathfrak{a} \ominus \mathfrak{b}) \oplus (\mathfrak{n} \ominus \mathfrak{v})$.

Theorem (Díaz-Ramos, LN). If \mathcal{F} is a (nontrivial) polar standard homogeneous foliation on $\mathbb{H}H^n$, then \mathcal{F} is isometrically congruent to the orbit foliation induced by some $S_{\mathfrak{b}, \mathfrak{v}}$.

Polar homogeneous foliations on hyperbolic spaces

Juan Manuel Lorenzo Naveiro
University of Oklahoma
Symmetry and Shape, 05/11/2025

Joint work with J. C. Díaz-Ramos
(Universidade de Santiago de Compostela)