

---

# MIDTERM EXAM 1 (FORM B)

---

NAME: \_\_\_\_\_

ID N°: \_\_\_\_\_

## INSTRUCTIONS

- This is a closed-book exam. You may not use your notes, textbook or any outside resources to assist you.
- Write your answer to each question in the space assigned to each problem. If you want the instructor to read the scratch working page for the answer to a specific problem, explicitly write that on the space assigned to said problem.
- Always remember to show your work. An answer without a proper justification will earn little to no credit, even if it is correct.
- Circle or box your final answer to each question.
- Make sure that your answers are organized and legible. Use a dark pencil or a pen. Any work that is hard or impossible to read will not be graded.
- In this exam, the use of determinants is not allowed. We will allow determinants once we cover them in this course.
- The use of calculators of any type is not allowed.

This exam consists of four problems, and your total score will be out of  $10 + 10 + 10 + 10 = 40$  points.

**Good Luck!**

This page is intentionally left blank

**Problem 1** (10 points). Suppose  $A$  and  $B$  are  $5 \times 3$  matrices.

- (a) (6 points) Find, in terms of  $A$  and  $B$ , the matrix  $X$  that satisfies

$$2(X - 3A^T)^T = 6X^T - B.$$

What is the size of  $X$ ?

- (b) (4 points) Calculate the matrix  $C^2 + C - 17I_2$ , where

$$C = \begin{bmatrix} -2 & 3 \\ 5 & 1 \end{bmatrix}.$$

**Problem 2** (2 points per part). On each of the following questions, **completely fill in** the circle next to the correct answer. Each question only has one correct answer. You do not need to justify your choices.

- (1) If  $A$  is an  $m \times n$  matrix, the rank  $r$  of  $A$  always satisfies:
- ☐  $r \leq m$  (but not necessarily  $r \leq n$ ).
  - ☐  $r \leq n$  (but not necessarily  $r \leq m$ ).
  - ☐  $r \leq n$  and  $r \leq m$ .
  - ☐ None of the above.
- (2) A system of linear equations is called *consistent* if:
- ☐ It has exactly one solution.
  - ☐ It has infinitely many solutions.
  - ☐ It has no solutions.
  - ☐ It has at least one solution.
- (3) If  $A\mathbf{x} = \mathbf{b}$  is a *non-homogeneous* linear system and  $\mathbf{x}_1, \mathbf{x}_2$  are solutions to it, then:
- ☐  $\mathbf{x}_1 + \mathbf{x}_2$  is a solution to  $A\mathbf{x} = \mathbf{b}$ .
  - ☐  $\mathbf{x}_1 - \mathbf{x}_2$  is a solution to  $A\mathbf{x} = \mathbf{b}$ .
  - ☐  $\mathbf{x}_1 + \mathbf{x}_2$  is a solution to  $A\mathbf{x} = \mathbf{0}$ .
  - ☐  $\mathbf{x}_1 - \mathbf{x}_2$  is a solution to  $A\mathbf{x} = \mathbf{0}$ .
- (4) Given the matrices  $X = \begin{bmatrix} 1 & 0 & 2 \end{bmatrix}$  and  $Y = \begin{bmatrix} 2 & 1 & 3 \end{bmatrix}^T$ , we have:
- ☐ The matrices  $5X + Y$ ,  $X(2Y)$  and  $4Y^T$  exist.
  - ☐ The matrices  $XY$ ,  $YX$  and  $2X + 5Y^T$  exist.
  - ☐ The matrices  $Y(3X^T)$ ,  $(X + Y^T)^T$  and  $Y^2$  exist.
  - ☐ The matrices  $X^T X$ ,  $-2X + Y$  and  $YX^2$  exist.
- (5) A homogeneous system of equations  $A\mathbf{x} = \mathbf{0}$  has 20 variables and 8 equations. If the augmented matrix  $[A | \mathbf{0}]$  has rank 5, then:
- ☐ The system is consistent regardless of  $A$ , and its general solution involves 12 parameters.
  - ☐ The system is consistent regardless of  $A$ , and its general solution involves 15 parameters.
  - ☐ There are matrices  $A$  for which the system is inconsistent. However, when the system is consistent, its general solution involves 12 parameters.
  - ☐ There are matrices  $A$  for which the system is inconsistent. However, when the system is consistent, its general solution involves 15 parameters.

**Problem 3** (10 points). Consider the system of linear equations  $A\mathbf{x} = \mathbf{b}$  whose augmented matrix is:

$$[A | \mathbf{b}] = \left[ \begin{array}{ccc|c} 1 & 0 & 6 & a \\ 3 & 1 & 5 & b \\ -4 & -2 & 2 & c \end{array} \right].$$

- (a) (2 points) What conditions do  $a$ ,  $b$  and  $c$  have to satisfy in order for the system  $A\mathbf{x} = \mathbf{b}$  to be homogeneous?

- (b) (4 points) Using the Gaussian algorithm, show that the system  $A\mathbf{x} = \mathbf{b}$  is consistent if and only if  $c = 2a - 2b$ .

- (c) (4 points) Calculate a set of basic solutions to the system  $A\mathbf{x} = \mathbf{0}$ .  
(**Hint:** Recycle the calculations from the previous part)

**Problem 4** (10 points). Consider the matrix

$$A = \begin{bmatrix} 1 & -1 & 1 \\ -2 & 2 & -1 \\ 3 & -2 & 1 \end{bmatrix}.$$

(a) (6 points) Using elementary row operations, find the inverse  $A^{-1}$  of  $A$ .

(b) (4 points) Use the previous part to find the solutions to the system

$$\begin{cases} x - y + z = 2 \\ -2x + 2y - z = -1 \\ 3x - 2y + z = 3 \end{cases}$$

## Scratch working page