

Exam Summary

Module Name

Student

Last updated: 14. March 2026

Table of Contents

1. Week 1: Linear Algebra for Deep Learning	3
1.1. Chapter 1: Linear Algebra for Deep Learning	3

Week 1: Linear Algebra for Deep Learning

Chapter 1: Linear Algebra for Deep Learning

Part 1 — The Building Blocks

Type	Dims	Notation	Example
Scalar	0D	a	single loss value
Vector	1D	$\mathbf{x}$ — always column	$[1, 2, 3]^\top$
Matrix	2D	$\mathbf{A}$, element $A_{i,j}$	weight layer
Tensor	3D+	$\mathbf{A}$, element $A_{i,j,k}$	image batch

Column vector and matrix notation:

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \quad \mathbf{A} = \begin{pmatrix} A_{1,1} & A_{1,2} \\ A_{2,1} & A_{2,2} \\ A_{3,1} & A_{3,2} \end{pmatrix}$$

Part 2 — Operations

Matrix addition (same size only):

$$C_{i,j} = A_{i,j} + B_{i,j}$$

Scalar on matrix:

$$D_{i,j} = a \cdot B_{i,j} + c$$

Broadcasting (vector added to every column of matrix):

$$C_{i,j} = A_{i,j} + b_j$$

Matrix multiplication:

$$\mathbf{C} = \mathbf{A}\mathbf{B} \quad C_{i,j} = \sum_k A_{i,k} \cdot B_{k,j}$$

Shape: $(m \times k) \cdot (k \times n) = (m \times n)$ — inner dims must match.

Inner / dot product (vectors only):

$$\mathbf{x}^\top \mathbf{y} = \sum_i x_i y_i = \|\mathbf{x}\|_2 \|\mathbf{y}\|_2 \cos \theta$$

Hadamard (element-wise) product:

$$\mathbf{C} = \mathbf{A} \odot \mathbf{B} \quad C_{i,j} = A_{i,j} \cdot B_{i,j}$$

Properties of matrix multiplication:

- Distributive: $A(B + C) = AB + AC$
- Associative: $A(BC) = (AB)C$

Exam Trap: $AB \neq BA$ — matrix multiplication is **NOT commutative**. This is on every exam.

Transpose:

$$(A^\top)_{i,j} = A_{j,i} \quad (AB)^\top = B^\top A^\top$$

Part 3 — Norms

General L^p norm:

$$\|x\|_p = \left(\sum_i |x_i|^p \right)^{\frac{1}{p}}, \quad p \geq 1$$

Norm	Formula	Use
L^2 (Euclidean)	$\ x\ _2 = \sqrt{\sum_i x_i^2}$	most common
Squared L^2	$\ x\ _2^2 = x^\top x$	easier to compute
L^1	$\ x\ _1 = \sum_i x_i $	sparsity
L^0	count of non-zeros	not a true norm
L^∞	$\ x\ _\infty = \max_i x_i $	max element
Frobenius	$\ A\ _F = \sqrt{\sum_{i,j} A_{i,j}^2} = \sqrt{\text{tr}(AA^\top)}$	matrix size

Part 4 — Identity, Inverse, Solving Systems

$$A^{-1}A = I \quad Ax = b \Rightarrow x = A^{-1}b$$

A^{-1} exists only if: A is square AND columns are linearly independent (non-singular).

Situation	Solutions
$m > n$ (overdetermined)	None (generically)
$m < n$ (underdetermined)	Infinitely many
$m = n$, independent cols	Exactly one

Part 5 — Special Matrices**Diagonal:**

$$\text{diag}(v)x = v \odot x \quad \text{diag}(v)^{-1} = \text{diag}([1/v_1, \dots, 1/v_n]^\top)$$

Symmetric:

$$A = A^\top$$

Orthogonal (rows AND cols are orthonormal):

$$A^\top A = AA^\top = I \Rightarrow A^{-1} = A^\top$$

Inverting an orthogonal matrix = just transposing it.

Part 6 — Eigendecomposition

$$A\mathbf{v} = \lambda\mathbf{v}$$

- $\mathbf{v}$ = eigenvector (direction unchanged by A , unit length $\|\mathbf{v}\| = 1$)
- λ = eigenvalue (scaling factor)

Full decomposition:

$$A = V \text{diag}(\boldsymbol{\lambda}) V^{-1} \quad V = [\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(n)}]$$

Intuition: **rotate** (by V) $\rightarrow$ **scale** (by λ) $\rightarrow$ **rotate back** (V^{-1}).

For real symmetric matrices: n real eigenvalues, orthogonal eigenvectors $\rightarrow V^{-1} = V^\top$.

Optimization (exam!):

$$\max_{\|\mathbf{x}\|=1} \mathbf{x}^\top A \mathbf{x} \rightarrow \mathbf{x} = \text{eigenvector of max eigenvalue}$$

$$\min_{\|\mathbf{x}\|=1} \mathbf{x}^\top A \mathbf{x} \rightarrow \mathbf{x} = \text{eigenvector of min eigenvalue}$$

Part 7 — Singular Value Decomposition (SVD)

Works for **any** matrix (generalization of eigendecomposition):

$$A = U D V^\top$$

- U : orthogonal — left singular vectors
- D : diagonal — singular values
- V : orthogonal — right singular vectors

Part 8 — Moore-Penrose Pseudoinverse

When A^{-1} doesn't exist:

$$A^+ = V D^+ U^\top \quad (D^+ : \text{reciprocal of each non-zero diagonal})$$

Case	Result
More rows than cols	Minimises $\ A\mathbf{x} - \mathbf{b}\ _2$ (least squares)
More cols than rows	Minimises $\ \mathbf{x}\ _2$ among all solutions

Part 9 — Trace

$$\text{tr}(A) = \sum_i A_{i,i} = \sum_i \lambda_i$$

- $\text{tr}(A^\top) = \text{tr}(A)$
- $\text{tr}(ABC) = \text{tr}(CAB) = \text{tr}(BCA) \leftarrow$ cyclic invariant
- $\|A\|_F = \sqrt{\text{tr}(AA^\top)}$

Part 10 — Determinant

$$\det(A) = \prod_i \lambda_i$$

- $\det(A) = 0 \rightarrow$ singular, not invertible
- $\det(A) \neq 0 \rightarrow$ invertible
- Geometric meaning: factor by which A scales volume

Cheat Sheet**WEEK 1 — LINEAR ALGEBRA CHEAT SHEET**

Structures: scalar=0D · vector=1D(col) · matrix=2D · tensor=3D+

Multiply:

- Standard: $C_{i,j} = \sum_k A_{i,k} B_{k,j}$
- Dot product: $\mathbf{x}^\top \mathbf{y} = \|\mathbf{x}\| \|\mathbf{y}\| \cos \theta$
- Hadamard $\odot$: $C_{i,j} = A_{i,j} B_{i,j}$
- $AB \neq BA$ — NEVER commutative

Norms: $L^2 = \sqrt{\sum x^2}$ · $L^1 = \sum |x|$ · $L^\infty = \max |x|$ · $\|A\|_F = \sqrt{\text{tr}(AA^\top)}$

Special: Symmetric $A = A^\top$ · Orthogonal $A^{-1} = A^\top$ · Diagonal: invert = reciprocal diagonal

Eigen: $A\mathbf{v} = \lambda\mathbf{v} \rightarrow A = V \text{diag}(\lambda) V^{-1}$

SVD: $A = UDV^\top$ (always works)

Pseudoinverse: $A^+ = VD^+U^\top$

Trace: $\text{tr}(A) = \sum A_{i,i} = \sum \lambda_i$ (cyclic invariant)

Det: $\det(A) = \prod \lambda_i$ · zero = singular = not invertible