

1 Elliptic Equation

The linearised elliptic equation is

$$G = Hv - \frac{H^3}{3} \left(\frac{\partial^2 v}{\partial x^2} \right)$$

Taking the weak version of this we get that

$$\int_{\Omega} Gv \, dx = H \int_{\Omega} vv \, dx - \frac{H^3}{3} \int_{\Omega} \frac{\partial^2 v}{\partial x^2} v \, dx$$

$$\int_{\Omega} Gv \, dx = H \int_{\Omega} vv \, dx + \frac{H^3}{3} \int_{\Omega} \frac{\partial v}{\partial x} \frac{\partial v}{\partial x} \, dx$$

In particular for the basis function ϕ_j we must have

$$\int_{\Omega} G\phi_j \, dx = H \int_{\Omega} v\phi_j \, dx + \frac{H^3}{3} \int_{\Omega} \frac{\partial v}{\partial x} \frac{\partial (\phi_j)}{\partial x} \, dx$$

We use the FEM discretisation from []

$$G = \sum_j G_{j-1/2}^+ \psi_{j-1/2}^+ + G_{j+1/2}^- \psi_{j+1/2}^-$$

and

$$v = \sum_j v_{j-1/2} \phi_{j-1/2} + v_j \phi_j + v_{j+1/2} \phi_{j+1/2} \quad (1)$$