

$$\mathcal{C}_2 = \frac{2 \cos(k\Delta x) - 2}{\Delta x^2}$$

$$\mathcal{C}_4 = \frac{-2 \cos(2k\Delta x) + 32 \cos(k\Delta x) - 30}{12\Delta x^2}$$

$$\mathcal{G} = \left[ H - \frac{H^3}{3} \mathcal{C} \right]$$

$$\mathcal{M}_3 = \frac{26 - 2 \cos(k\Delta x)}{24}$$

$$\mathcal{M}_1 = \mathcal{M}_2 = 1. \quad \mathcal{R}_1^+ = e^{ik\Delta x} \text{ and } \mathcal{R}_1^- = 1.$$

$$\mathcal{R}_2^- = 1 + \frac{i \sin(k\Delta x)}{2}$$

$$\mathcal{R}_2^+ = e^{ik\Delta x} \left( 1 - \frac{i \sin(k\Delta x)}{2} \right)$$

$$R_3^- = \frac{\mathcal{M}_3}{6} [5 + -e^{-ik\Delta x} + 2e^{ik\Delta x}]$$

$$R_3^+ = \frac{\mathcal{M}_3 e^{ik\Delta x}}{6} [5 + 2e^{-ik\Delta x} - e^{ik\Delta x}]$$

$$R_2^u = \frac{e^{ik\Delta x} + 1}{2}$$

$$R_3^u = \frac{-3e^{2ik\Delta x} + 27e^{ik\Delta x} + 27 - 3e^{-ik\Delta x}}{48}$$

$$\mathcal{F}^{h,u} = H\mathcal{R}^u$$

$$\mathcal{F}^{h,h} = -\frac{\sqrt{gH}}{2} [\mathcal{R}^+ - \mathcal{R}^-]$$

$$\mathcal{F}^{u,u} = -\frac{\sqrt{gH}}{2} \mathcal{G} [\mathcal{R}^+ - \mathcal{R}^-]$$

$$\mathcal{F}^{u,h} = \frac{gH\mathcal{R}^- + gH\mathcal{R}^+}{2}$$

$$\mathcal{D} = 1 - e^{-ik\Delta x}$$

# 1 Taylor Expansions

Around 0 we have

$$\cos(x) = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \frac{1}{8!}x^8 - \frac{1}{10!}x^{10} + O(x^{12})$$

$$\sin(x) = \frac{1}{1!}x^1 - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \frac{1}{9!}x^9 + O(x^{11})$$

$$e^{ix} = \cos(x) + i \sin(x)$$

$$e^{ix} = 1 + ix - \frac{1}{2}x^2 - \frac{i}{3!}x^3 + \frac{1}{4!}x^4 + \frac{i}{5!}x^5 - \frac{1}{6!}x^6 - \frac{i}{7!}x^7 + \frac{1}{8!}x^8 + \frac{i}{9!}x^9 - \frac{1}{10!}x^{10} + O(x^{11})$$

# 2 Going Down List

So

$$\mathcal{G}_a = \left[ H + \frac{H^3}{3}k^2 \right]$$

$$\mathcal{C}_2 = \frac{2 \left( -\frac{1}{2!}(k\Delta x)^2 + \frac{1}{4!}(k\Delta x)^4 - \frac{1}{6!}(k\Delta x)^6 + \frac{1}{8!}(k\Delta x)^8 - \frac{1}{10!}(k\Delta x)^{10} + O(x^{12}) \right)}{\Delta x^2}$$

$$\mathcal{C}_2 = \frac{-(k\Delta x)^2 + \frac{2}{4!}(k\Delta x)^4 - \frac{2}{6!}(k\Delta x)^6 + \frac{2}{8!}(k\Delta x)^8 - \frac{2}{10!}(k\Delta x)^{10} + O(x^{12})}{\Delta x^2}$$

$$\mathcal{C}_2 = -k^2 + \frac{2}{4!}k^4(\Delta x)^2 - \frac{2}{6!}k^6(\Delta x)^4 + \frac{2}{8!}k^8(\Delta x)^6 - \frac{2}{10!}k^{10}(\Delta x)^8 + O(x^{10})$$

$$\mathcal{C}_2 = -k^2 + \frac{1}{12}k^4(\Delta x)^2 + O(x^4)$$

$$\mathcal{C}_4 = \frac{-2 \cos(2k\Delta x) + 32 \cos(k\Delta x) - 30}{12\Delta x^2}$$

$$-2 \cos(2k\Delta x) = -2 \left[ 1 - \frac{1}{2}2^2 k^2 \Delta x^2 + \frac{1}{24}2^4 k^4 \Delta x^4 - \frac{1}{720}2^6 k^6 \Delta x^6 + \frac{1}{40320}2^8 k^8 \Delta x^8 + O(\Delta x^{10}) \right]$$

$$= -2 + \frac{1}{2}2^3 k^2 \Delta x^2 - \frac{1}{24}2^5 k^4 \Delta x^4 + \frac{1}{720}2^7 k^6 \Delta x^6 - \frac{1}{40320}2^9 k^8 \Delta x^8 + O(\Delta x^{10})$$

$$= -2 + \frac{8}{2}k^2 \Delta x^2 - \frac{32}{24}k^4 \Delta x^4 + \frac{128}{720}k^6 \Delta x^6 - \frac{512}{40320}k^8 \Delta x^8 + O(\Delta x^{10})$$

$$32 \cos(k\Delta x) = 32 - 16(k\Delta x)^2 + \frac{32}{24}(k\Delta x)^4 - \frac{32}{720}(k\Delta x)^6 + \frac{32}{40320}(k\Delta x)^8 + O((k\Delta x)^{10})$$

$$32 \cos(k\Delta x) - 2 \cos(2k\Delta x) = 30 - 12(k\Delta x)^2 + \frac{96}{720}(k\Delta x)^6 - \frac{480}{40320}(k\Delta x)^8 + O((k\Delta x)^{10})$$

$$\mathcal{C}_4 = \frac{1}{12\Delta x^2} \left[ -12(k\Delta x)^2 + \frac{96}{720}(k\Delta x)^6 - \frac{480}{40320}(k\Delta x)^8 + O((k\Delta x)^{10}) \right]$$

$$\mathcal{C}_4 = \left[ -(k)^2 + \frac{96}{8640}k^6(\Delta x)^4 - \frac{480}{483840}k^8(\Delta x)^6 + O((k\Delta x)^8) \right]$$

$$\mathcal{C}_4 = \left[ -(k)^2 + \frac{1}{90}k^6(\Delta x)^4 + O(\Delta x^6) \right]$$

Analytically

$$G = uh - h^3/3(u_{xx})$$

For the fourier nodes we have

$$G = uH - (-1)(k^2)H^3/3(u)$$

$$G = uH + (k^2)H^3/3u$$

Using above derivative approximations

$$\mathcal{G}_2 = H - \frac{H^3}{3} \left[ -k^2 + \frac{1}{12}k^4(\Delta x)^2 + O(x^4) \right]$$

$$\mathcal{G}_2 = H + \frac{H^3}{3}k^2 - \frac{H^3k^4(\Delta x)^2}{36} + O(x^4)$$

$$\mathcal{G}_4 = H - \frac{H^3}{3} \left[ -(k)^2 + \frac{1}{90}k^6(\Delta x)^4 + O(\Delta x^6) \right]$$

$$\mathcal{G}_4 = H + \frac{H^3k^2}{3} - \frac{H^3k^6(\Delta x)^4}{270} + O(\Delta x^6)$$

Analytically we have

$$\bar{u}_j = \frac{1}{\Delta x} \int_{x_{j-1/2}}^{x_{j+1/2}} u_j dx$$

$$\bar{u}_j = \frac{1}{\Delta x} \frac{-i}{k} u_j \left[ e^{ik\Delta x/2} - e^{-ik\Delta x/2} \right] dx$$

$$\bar{u}_j = \frac{1}{\Delta x} \frac{-i}{k} u_j [2i \sin(ik\Delta x/2)]$$

$$\bar{u}_j = \frac{1}{\Delta x} \frac{2}{k} [\sin(ik\Delta x/2)] u_j$$

So

$$u_j = k \frac{\Delta x}{2} [\csc(ik\Delta x/2)] \bar{u}_j$$

$$\mathcal{M}_a = k \frac{\Delta x}{2} [\csc(ik\Delta x/2)]$$

The Taylor expansion of this is

$$\mathcal{M}_a = 1 + \frac{1}{24}(k\Delta x)^2 + \frac{7}{5760}(k\Delta x)^4 + O(x^6)$$

$$\mathcal{M}_3 = \frac{26 - 2 \left( 1 - \frac{1}{2!}(k\Delta x)^2 + \frac{1}{4!}(k\Delta x)^4 - \frac{1}{6!}(k\Delta x)^6 + \frac{1}{8!}(k\Delta x)^8 - \frac{1}{10!}(k\Delta x)^{10} + O(x^{12}) \right)}{24}$$

$$\mathcal{M}_3 = \frac{12 - \left(-\frac{1}{2!}(k\Delta x)^2 + \frac{1}{4!}(k\Delta x)^4 - \frac{1}{6!}(k\Delta x)^6 + \frac{1}{8!}(k\Delta x)^8 - \frac{1}{10!}(k\Delta x)^{10} + O(x^{12})\right)}{12}$$

$$\mathcal{M}_3 = \frac{12 + \frac{1}{2!}(k\Delta x)^2 - \frac{1}{4!}(k\Delta x)^4 + \frac{1}{6!}(k\Delta x)^6 - \frac{1}{8!}(k\Delta x)^8 + \frac{1}{10!}(k\Delta x)^{10} + O(x^{12})}{12}$$

$$\mathcal{M}_3 = 1 + \frac{1}{24}(k\Delta x)^2 - \frac{1}{288}(k\Delta x)^4 + \frac{1}{8640}(k\Delta x)^6 + O(x^8)$$

$$\mathcal{M}_3 = 1 + \frac{1}{24}(k\Delta x)^2 - \frac{1}{288}(k\Delta x)^4 + O(x^6)$$

So we see that  $\mathcal{M}_3$  has a fourth order error compared to the analytic solution.

Reconstruction, analytically we have

$$u_{j+1/2} = u_j e^{ik\Delta x/2}$$

So

$$\mathcal{R}_a = \mathcal{R}_a^+ = \mathcal{R}_a^- = e^{ik\Delta x/2}$$

So we have the Taylor expansion

$$\mathcal{R}_a = 1 + i(k\Delta x/2) - \frac{1}{2}(k\Delta x/2)^2 - \frac{i}{3!}(k\Delta x/2)^3 + \frac{1}{4!}(k\Delta x/2)^4 + \frac{i}{5!}(k\Delta x/2)^5 - \frac{1}{6!}(k\Delta x/2)^6 + O(x^7)$$

$$\mathcal{R}_a = 1 + i\frac{1}{2}(k\Delta x) - \frac{1}{2}\frac{1}{4}(k\Delta x)^2 - \frac{i}{3!}\frac{1}{8}(k\Delta x)^3 + \frac{1}{4!}\frac{1}{16}(k\Delta x)^4 + \frac{i}{5!}\frac{1}{32}(k\Delta x)^5 - \frac{1}{6!}\frac{1}{64}(k\Delta x)^6 + O(x^7)$$

$$\mathcal{R}_a = 1 + \frac{i}{2}k\Delta x - \frac{1}{8}k^2\Delta x^2 - \frac{i}{48}k^3\Delta x^3 + O(x^4)$$

$$\mathcal{R}_1^+ = e^{ik\Delta x} \text{ and } \mathcal{R}_1^- = 1.$$

$$\mathcal{R}_1^+ = 1 + ik\Delta x - \frac{1}{2}(k\Delta x)^2 - \frac{i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 + O((k\Delta x)^5)$$

So both have first order errors.

$$\mathcal{R}_2^- = 1 + \frac{i \sin(k\Delta x)}{2}$$

$$\mathcal{R}_2^- = 1 + \frac{i}{2} \left[ \frac{1}{1!}(k\Delta x)^1 - \frac{1}{3!}(k\Delta x)^3 + \frac{1}{5!}(k\Delta x)^5 - \frac{1}{7!}(k\Delta x)^7 + \frac{1}{9!}(k\Delta x)^9 + O(x^{11}) \right]$$

$$\mathcal{R}_2^- = 1 + \frac{i}{2}(k\Delta x) - \frac{i}{2} \frac{1}{3!}(k\Delta x)^3 + \frac{i}{2} \frac{1}{5!}(k\Delta x)^5 - \frac{i}{2} \frac{1}{7!}(k\Delta x)^7 + \frac{i}{2} \frac{1}{9!}(k\Delta x)^9 + O(x^{11})$$

$$\mathcal{R}_2^+ = e^{ik\Delta x} \left( 1 - \frac{i \sin(k\Delta x)}{2} \right)$$

$$\mathcal{R}_2^+ = e^{ik\Delta x} \left( 1 - \frac{i}{2}(k\Delta x) + \frac{i}{2} \frac{1}{3!}(k\Delta x)^3 - \frac{i}{2} \frac{1}{5!}(k\Delta x)^5 + \frac{i}{2} \frac{1}{7!}(k\Delta x)^7 - \frac{i}{2} \frac{1}{9!}(k\Delta x)^9 + O(x^{11}) \right)$$

$$\begin{aligned} \mathcal{R}_2^+ &= \left( 1 + i(k\Delta x) - \frac{1}{2}(k\Delta x)^2 - \frac{i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 + \frac{i}{5!}(k\Delta x)^5 + O((\Delta x^2)) \right) \\ &\quad \times \left( 1 - \frac{i}{2}(k\Delta x) + \frac{i}{2} \frac{1}{3!}(k\Delta x)^3 - \frac{i}{2} \frac{1}{5!}(k\Delta x)^5 + O(x^7) \right) \quad (1) \end{aligned}$$

$$\mathcal{R}_2^+ = 1 + \frac{ik\Delta x}{2} + \frac{i}{6}k^3\Delta x^3 + O(\Delta x^4)$$

So for both second order reconstructions we have a second order error.

$$R_3^- = \frac{\mathcal{M}_3}{6} [5 + -e^{-ik\Delta x} + 2e^{ik\Delta x}]$$

$$\begin{aligned} R_3^- &= \frac{1 + \frac{1}{24}(k\Delta x)^2 - \frac{1}{288}(k\Delta x)^4 + O(x^6)}{6} \\ &\quad \left[ 5 - \left[ 1 - ik\Delta x - \frac{1}{2}(k\Delta x)^2 + \frac{i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 - \frac{i}{5!}(k\Delta x)^5 - O(\Delta x^6) \right] \right. \\ &\quad \left. + 2 \left[ 1 + i(k\Delta x) - \frac{1}{2}(k\Delta x)^2 - \frac{i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 + \frac{i}{5!}(k\Delta x)^5 + O(\Delta x^6) \right] \right] \quad (2) \end{aligned}$$

$$R_3^- = \frac{1 + \frac{1}{24}(k\Delta x)^2 - \frac{1}{288}(k\Delta x)^4 + O(x^6)}{6} \left[ 6 + 3ik\Delta x - \frac{1}{2}(k\Delta x)^2 - \frac{3i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 O(\Delta x^5) \right] \quad (3)$$

$$\begin{aligned} R_3^- &= \frac{1}{6} \left[ 6 + 3ik\Delta x - \frac{1}{2}(k\Delta x)^2 - \frac{3i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 O(\Delta x^5) \right] \\ &+ \frac{\frac{1}{24}(k\Delta x)^2}{6} \left[ 6 + 3ik\Delta x - \frac{1}{2}(k\Delta x)^2 - \frac{3i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 O(\Delta x^5) \right] \\ &+ \frac{-\frac{1}{288}(k\Delta x)^4}{6} \left[ 6 + 3ik\Delta x - \frac{1}{2}(k\Delta x)^2 - \frac{3i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 O(\Delta x^5) \right] \\ &+ O(x^6) \quad (4) \end{aligned}$$

$$\begin{aligned} R_3^- &= 1 + \frac{i}{2}k\Delta x \\ &+ \frac{1}{6} \left[ -\frac{1}{2}(k\Delta x)^2 - \frac{3i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 O(\Delta x^5) \right] \\ &+ \frac{\frac{1}{24}(k\Delta x)^2}{6} \left[ 6 + 3ik\Delta x - \frac{1}{2}(k\Delta x)^2 - \frac{3i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 O(\Delta x^5) \right] \\ &+ \frac{-\frac{1}{288}(k\Delta x)^4}{6} \left[ 6 + 3ik\Delta x - \frac{1}{2}(k\Delta x)^2 - \frac{3i}{3!}(k\Delta x)^3 + \frac{1}{4!}(k\Delta x)^4 O(\Delta x^5) \right] \\ &+ O(x^6) \quad (5) \end{aligned}$$

$$R_3^- = 1 + \frac{i}{2}k\Delta x - \frac{1}{24}(k\Delta x)^2 - \frac{1}{16}(k\Delta x)^3 + O(\Delta x^4) \quad (6)$$

So our reconstruction is under order...

$$R_3^+ = \frac{\mathcal{M}_3 e^{ik\Delta x}}{6} [5 + 2e^{-ik\Delta x} - e^{ik\Delta x}]$$

from wolfram

$$R_3^+ = 1 + \frac{i}{2}k\Delta x - \frac{1}{24}(k\Delta x)^2 - \frac{5}{48}(k\Delta x)^3 + O(\Delta x^4)$$

Again we have second order errors. But they are the same... Hopefully some cancellation works out.

$$R_2^u = \frac{e^{ik\Delta x} + 1}{2}$$

$$R_2^u = 1 + \frac{i}{2}k\Delta x - \frac{1}{4}(k\Delta x)^2 - \frac{i}{12}(k\Delta x)^3 + O(\Delta x^4)$$

$$R_3^u = \frac{-3e^{2ik\Delta x} + 27e^{ik\Delta x} + 27 - 3e^{-ik\Delta x}}{48}$$

$$R_3^u = 1 + \frac{i}{2}k\Delta x - \frac{1}{8}(k\Delta x)^2 - \frac{i}{48}(k\Delta x)^3 + O(\Delta x^4)$$

So here we get correct order of errors. In fact the reconstruction is higher than 3rd.

Ok so for flux we have to get the analytic result. Is

$$F_1^h(u_j, h_j) = kiHu_j$$

$$F_a^h(u_j, h_j) = kigHh_j$$

$$\mathcal{D} = 1 - e^{-ik\Delta x}$$

$$\mathcal{D} = ik\Delta x + \frac{(k\Delta x)^2}{2} - \frac{i(k\Delta x)^3}{6} + O(\Delta x^4)$$

$$\mathcal{F}_1^{h,u} = H\mathcal{R}_2^u$$

$$\mathcal{F}_1^{h,u} = H \left[ 1 + \frac{i}{2}k\Delta x - \frac{1}{4}(k\Delta x)^2 - \frac{i}{12}(k\Delta x)^3 + O(\Delta x^4) \right]$$

$$\mathcal{D}\mathcal{F}_1^{h,u} = H \left[ ik\Delta x - \frac{i}{6}(k\Delta x)^3 + O(\Delta x^4) \right]$$



$$\begin{aligned}
\mathcal{F}_1^{h,h} &= -\frac{\sqrt{gH}}{2} [\mathcal{R}_1^+ - \mathcal{R}_1^-] \\
\mathcal{F}_1^{h,h} &= -\frac{\sqrt{gH}}{2} \left[ ik\Delta x - \frac{(k\Delta x)^2}{2} - \frac{i(k\Delta x)^3}{6} + O(\Delta x^4) \right] \\
\mathcal{D}\mathcal{F}_1^{h,h} &= -\frac{\sqrt{gH}}{2} [(k\Delta x)^2 + O(\Delta x^4)] \\
F_1^h &= \frac{1}{\Delta x} [\mathcal{D}\mathcal{F}_1^{h,u} u_j + \mathcal{D}\mathcal{F}_1^{h,h} h_j] \\
F_1^h &= H \left[ ik - \frac{i}{6} k^3 (\Delta x)^2 + O(\Delta x^3) \right] u_j \\
&\quad + -\frac{\sqrt{gH}}{2} [k^2 (\Delta x) + O(\Delta x^3)] h_j \quad (7)
\end{aligned}$$

So first order error is completely on  $h$  interesting.

$$\begin{aligned}
\mathcal{F}_1^{u,u} &= -\frac{\sqrt{gH}}{2} \mathcal{G}_1 [\mathcal{R}_1^+ - \mathcal{R}_1^-] \\
\mathcal{F}_1^{u,u} &= -\frac{\sqrt{gH}}{2} \left[ \frac{1}{3} iH(H^2 + 3)(k\Delta x) - \frac{1}{6} iH(H^2 + 3)(k\Delta x)^2 - \frac{1}{12} iH(H^2 + 3)(k\Delta x)^3 + O(\Delta x^4) \right] \\
\mathcal{D}\mathcal{F}_1^{u,u} &= -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{3} iH(H^2 + 3)(k\Delta x)^2 + \frac{1}{36} H(2H^2 + 3)(k\Delta x)^4 + O(\Delta x^6) \right] \\
\mathcal{F}_1^{u,h} &= \frac{gH}{2} [\mathcal{R}_1^- + \mathcal{R}_1^+] \\
\mathcal{F}_1^{u,h} &= \frac{gH}{2} \left[ 2 + ik\Delta x - \frac{(k\Delta x)^2}{2} - \frac{i(k\Delta x)^3}{6} + O(\Delta x^4) \right] \\
\mathcal{D}\mathcal{F}_1^{u,h} &= \frac{gH}{2} \left[ 2ik\Delta x - \frac{i(k\Delta x)^3}{3} + O(\Delta x^4) \right]
\end{aligned}$$

So

$$F_1^u = -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{3}iH(H^2 + 3)k^2(\Delta x) + \frac{1}{36}H(2H^2 + 3)k^4(\Delta x)^3 + O(\Delta x^5) \right] u_j \\ + \frac{gH}{2} \left[ 2ik - \frac{ik^3(\Delta x)^2}{3} + O(\Delta x^3) \right] h_j \quad (8)$$

$$F_1^u = -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{3}iH(H^2 + 3)k^2(\Delta x) + \frac{1}{36}H(2H^2 + 3)k^4(\Delta x)^3 + O(\Delta x^5) \right] u_j \\ + gH \left[ ik - \frac{ik^3(\Delta x)^2}{6} + O(\Delta x^3) \right] h_j \quad (9)$$

OK Seeing a pattern, but we have to make sure this holds as we go, anyway we now calculate just the D F together always to make it simpler.

$$\mathcal{DF}_2^{h,u} = H\mathcal{DR}_2^u$$

$$\mathcal{DF}_2^{h,u} = H \left[ i(k\Delta x) - \frac{i}{6}(k\Delta x)^3 + \frac{i}{120}(k\Delta x)^5 + O(\Delta x^6) \right]$$

$$\mathcal{DF}^{h,h} = -\frac{\sqrt{gH}}{2} \mathcal{D} [\mathcal{R}^+ - \mathcal{R}^-]$$

$$\mathcal{DF}^{h,h} = -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{4}(k\Delta x)^4 + \frac{1}{24}(k\Delta x)^6 + O(\Delta x^8) \right]$$

$$F_2^h = H \left[ 2ik - \frac{i}{3}k^3(\Delta x)^2 + \frac{i}{60}k^5(\Delta x)^4 + O(\Delta x^5) \right] u_j \\ + -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{4}k^4(\Delta x)^3 + \frac{1}{24}k^6(\Delta x)^5 + O(\Delta x^7) \right] h_j \quad (10)$$

$$\mathcal{DF}^{u,u} = -\frac{\sqrt{gH}}{2} \mathcal{DG} [\mathcal{R}^+ - \mathcal{R}^-]$$

$$\mathcal{DF}^{u,u} = -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{12}H(H^2 + 3)k^4\Delta x^4 + \frac{1}{48}H(H^2 + 2)k^6\Delta x^6 + O(\Delta x^7) \right]$$

$$\mathcal{DF}^{u,h} = \frac{gH}{2} \mathcal{D}[\mathcal{R}^- + \mathcal{R}^+]$$

$$\mathcal{DF}^{u,h} = \frac{gH}{2} \left[ 2ik\Delta x + \frac{i}{6}k^3\Delta x^3 - \frac{13i}{120}k^5\Delta x^5 + O(\Delta x^7) \right]$$

$$\begin{aligned} F_2^u = & -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{12}H(H^2 + 3)k^4\Delta x^3 + \frac{1}{48}H(H^2 + 2)k^6\Delta x^5 + O(\Delta x^6) \right] u_j \\ & + \frac{gH}{2} \left[ 2ik + \frac{i}{6}k^3\Delta x^2 - \frac{13i}{120}k^5\Delta x^4 + O(\Delta x^6) \right] h_j \quad (11) \end{aligned}$$

So everyting is the correct order for the second order scheme as well.

$$\mathcal{DF}_3^{h,u} = H\mathcal{DR}_3^u$$

$$\mathcal{DF}_3^{h,u} = H \left[ ik\Delta x - \frac{i}{24}k^3\Delta x^3 - \frac{11i}{480}k^5\Delta x^5 + O(\Delta x^7) \right]$$

$$\mathcal{DF}_3^{h,h} = -\frac{\sqrt{gH}}{2} \mathcal{D} [\mathcal{R}_3^+ - \mathcal{R}_3^-]$$

$$\mathcal{DF}_3^{h,h} = -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{6}k^4\Delta x^4 + \frac{1}{48}k^6\Delta x^6 - \frac{1}{2880}k^8\Delta x^8 + O(\Delta x^9) \right]$$

$$\begin{aligned} F_3^h = & H \left[ ik - \frac{i}{24}k^3\Delta x^2 - \frac{11i}{480}k^5\Delta x^4 + O(\Delta x^6) \right] u_j \\ & + -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{6}k^4\Delta x^3 + \frac{1}{48}k^6\Delta x^5 - \frac{1}{2880}k^8\Delta x^7 + O(\Delta x^8) \right] h_j \quad (12) \end{aligned}$$

$$\mathcal{DF}_3^{u,u} = -\frac{\sqrt{gH}}{2} \mathcal{DG}_3 [\mathcal{R}_3^+ - \mathcal{R}_3^-]$$

$$\begin{aligned} \mathcal{DF}_3^{u,u} = & -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{6}k^4\Delta x^4 + \frac{1}{48}k^6\Delta x^6 - \frac{1}{2880}k^8\Delta x^8 + O(\Delta x^9) \right] \\ & \left[ \left( \frac{H^3}{3} + H \right) - \frac{H^3k^4\Delta x^4}{270} + \frac{H^3k^6\Delta x^6}{3024} - \frac{H^3k^8\Delta x^8}{64800} + O(\Delta x^9) \right] \quad (13) \end{aligned}$$

$$\mathcal{DF}_3^{u,u} = -\frac{\sqrt{gH}}{2} \left[ -\left(\frac{H^3}{3} + H\right) \frac{1}{6} k^4 \Delta x^4 + \frac{1}{48} \left(\frac{H^3}{3} + H\right) k^6 \Delta x^6 + O(\Delta x^8) \right] \quad (14)$$

$$\mathcal{DF}_3^{u,h} = \frac{gH}{2} \mathcal{D}[\mathcal{R}_3^- + \mathcal{R}_3^+]$$

$$\mathcal{DF}_3^{u,h} = \frac{gH}{2} \left[ 2ik\Delta x + \frac{i}{12} k^3 \Delta x^3 - \frac{53i\Delta x^5}{720} + O(\Delta x^7) \right]$$

$$F_3^u = \frac{\sqrt{gH}}{2} \left[ -\left(\frac{H^3}{3} + H\right) \frac{1}{6} k^4 \Delta x^3 + \frac{1}{48} \left(\frac{H^3}{3} + H\right) k^6 \Delta x^5 + O(\Delta x^7) \right] u_j \\ + \frac{gH}{2} \left[ 2ik + \frac{i}{12} k^3 \Delta x^2 - \frac{53ik^5 \Delta x^4}{720} + O(\Delta x^6) \right] h_j \quad (15)$$

So yes our second order error in the flux continues over, it does however occur only in the imaginary direction, does that matter?

### 3 3rd

M was defined wrong, so we have an error throughout, only effects third order.

$$R_3^- = 1 + \frac{i}{2} k \Delta x - \frac{1}{8} (k \Delta x)^2 - \frac{5}{48} (k \Delta x)^3 + O(\Delta x^4)$$

$$R_3^+ = 1 + \frac{i}{2} k \Delta x - \frac{1}{8} (k \Delta x)^2 + \frac{1}{16} (k \Delta x)^3 + O(\Delta x^4)$$

$$\mathcal{DF}_3^{h,u} = H \mathcal{DR}_3^u$$

$$\mathcal{DF}_3^{h,u} = H \left[ ik\Delta x - \frac{i}{24} k^3 \Delta x^3 - \frac{11i}{480} k^5 \Delta x^5 + O(\Delta x^7) \right]$$

$$\mathcal{DF}_3^{h,h} = -\frac{\sqrt{gH}}{2} \mathcal{D}[\mathcal{R}_3^+ - \mathcal{R}_3^-]$$

$$\mathcal{DF}_3^{h,h} = -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{6}k^4\Delta x^4 + \frac{5}{144}k^6\Delta x^6 - \frac{71}{17280}k^8\Delta x^8 + O(\Delta x^9) \right]$$

$$\mathcal{DF}_3^{u,u} = -\frac{\sqrt{gH}}{2} \mathcal{DG}_3 [\mathcal{R}_3^+ - \mathcal{R}_3^-]$$

$$\begin{aligned} \mathcal{DF}_3^{u,u} = & -\frac{\sqrt{gH}}{2} \left[ -\frac{1}{6}k^4\Delta x^4 + \frac{5}{144}k^6\Delta x^6 - \frac{71}{17280}k^8\Delta x^8 + O(\Delta x^9) \right] \\ & \left[ \left( \frac{H^3}{3} + H \right) - \frac{H^3k^4\Delta x^4}{270} + \frac{H^3k^6\Delta x^6}{3024} - \frac{H^3k^8\Delta x^8}{64800} + O(\Delta x^9) \right] \end{aligned} \quad (16)$$

$$\mathcal{DF}_3^{u,u} = -\frac{\sqrt{gH}}{2} \left[ -\left( \frac{H^3}{3} + H \right) \frac{1}{6}k^4\Delta x^4 + \frac{5}{144} \left( \frac{H^3}{3} + H \right) k^6\Delta x^6 + O(\Delta x^8) \right] \quad (17)$$

$$\mathcal{DF}_3^{u,h} = \frac{gH}{2} \mathcal{D}[\mathcal{R}_3^- + \mathcal{R}_3^+]$$

$$\mathcal{DF}_3^{u,h} = \frac{gH}{2} \left[ 2ik\Delta x - \frac{i}{12}k^3\Delta x^3 - \frac{9i\Delta x^5}{160} + O(\Delta x^7) \right]$$