

Generalised Serre-Green-Naghdi Model

June 4, 2020

0.1 Numerical Experiments

0.1.1 Smooth Dambreak

$$h(x, 0) = h_0 + \frac{h_1 - h_0}{2} \left(1 + \tanh \left(\frac{x}{\alpha} \right) \right) \quad (1)$$

$$u(x, 0) = 0 \quad (2)$$


$$G(x, 0) = 0 \quad (3)$$

$$\alpha = 2$$


$$\alpha = 0.1$$

References


- [1] Clamond, D. and D. Dutykh, Non-dispersive conservative regularisation of nonlinear shallow water (and isentropic Euler equations), *Communications in Nonlinear Science and Numerical Simulation*, 55(42), 237-247.
- [2] Dutykh, D., M. Hoefer and D. Mitsotakis, Solitary wave solutions and their interactions for fully nonlinear water waves with surface tension in the generalized Serre equations, *Theoretical and Computational Fluid Dynamics*, 32(3), 371-397.
- [3] L.C. Evans, *Partial Differential Equations*, Graduate Studies in Mathematics, Volume 19, American Mathematical Society, New York, (1997).
- [4] A. Harten, High resolution schemes for hyperbolic conservation laws, *Journal of Computational Physics*, 49 (3) (1983) 357-393.
- [5] A. Kurganov, S. Noelle, G. Petrova, Semidiscrete central-upwind schemes for hyperbolic conservation laws and Hamilton-Jacobi equations, *Journal of Scientific Computing*, Society for Industrial and Applied Mathematics, 23 (3) (2002) 707-740.
- [6] B. van Leer, Towards the ultimate conservative difference scheme, V. A second-order sequel to Godunov's method, *Journal of Computational Physics*, 32 (1) (1979) 101-136.
- [7] C.W. Shu, S. Osher, Efficient implementation of essentially non-oscillatory shock-capturing schemes, *Journal of Computational Physics*, 77 (2) (1988) 439-471.
- [8] F. Serre, Contribution à l'étude des écoulements permanents et variables dans les canaux, *La Houille Blanche*, 6 (1953) 830-872.
- [9] C. Zoppou, Numerical solution of the one-dimensional and cylindrical Serre equations for rapidly varying free surface flows, Ph.D., Mathematical Sciences Institute, College of Physical and Mathematical Sciences, The Australian National University, (2014).
- [10] C. Zoppou, J. Pitt and S.G. Roberts, Numerical solution of the fully non-linear weakly dispersive Serre equations for steep gradient flows, *Applied Mathematical Modelling*, 48, 70-95.




(a) h



(b) h



(c) u



(d) G

Figure 1: Initial Conditions $\alpha = 2$




Figure 2: Improve Dispersion Serre Family $\beta_1 = \beta_2$ for smooth dam-break $\alpha = 2$ at $t = 15s$. $\beta_1 = \beta_2 = 0.1$ to $\beta_1 = \beta_2 = 1$.




Figure 3: Regularised SWWE Family $\beta_1 = \beta_2 - \frac{2}{3}$ for smooth dam-break $\alpha = 2$ at $t = 15s$. $\beta_1 = \beta_2 - \frac{2}{3}$ with $\beta_2 = 0$ to $\beta_2 = 5$.




Figure 4: Region 2 Family $\beta_1 = \beta_2 - \frac{2}{3} - k$ and $\beta_2 = 2k$ (should find something better so we get expanding wave train), so $\alpha = 2$ for smooth dam-break $\alpha = 2$ at $t = 15s$.




Figure 5: Initial Conditions $\alpha = 0.1$




Figure 6: Improve Dispersion Serre Family $\beta_1 = \beta_2$ for smooth dam-break $\alpha = 0.1$ at $t = 15s$. $\beta_1 = \beta_2 = 0.1$ to $\beta_1 = \beta_2 = 1$.




Figure 7: Improve Dispersion Serre Family $\beta_1 = \beta_2$ for smooth dam-break $\alpha = 2$ at $t = 15s$. $\beta_1 = \beta_2 - \frac{2}{3}$ with $\beta_2 = 0$ to $\beta_2 = 5$.