

# Vascular Model Repository

## Specifications Document



0075\_1001

<b>Species</b>	Human
<b>Anatomy</b>	Aorta
<b>Disease</b>	Congenital Heart Disease Single Ventricle Defect
<b>Procedure</b>	Fontan

# Clinical Significance and Background

## Aorta

The largest blood vessel and the human body's primary artery, the aorta is responsible for carrying oxygenated blood pumped from the heart to rest of the body. The aorta is divided into four sections: the ascending aorta, the aortic arch, the thoracic aorta, and the abdominal aorta.

The ascending aorta starts at the left ventricle of the heart where at the root, it supplies blood to the heart muscle through the coronary arteries. From the aortic root, the ascending aorta continues to rise up until it reaches the aortic arch.

The aortic arch loops over the bifurcation of the pulmonary trunk and has three major artery branches leaving through the top: the brachiocephalic trunk, the left common carotid artery, and the left subclavian artery. The brachiocephalic trunk sends blood to the right side of the brain and right arm/neck/chest while the left common carotid artery sends blood to the left side of the brain and the left subclavian artery sends blood to the left arm/neck/chest.

After the aortic arch, the aorta begins to descend all the way to the abdomen. The section of the descending aorta that starts after the aortic arch and ends at the diaphragm is called the thoracic aorta, and it supplies blood to the chest and spinal cord.

The last section of the aorta, the abdominal aorta, starts at the diaphragm and ends just above the pelvis. This section is responsible for supplying blood to the stomach, kidneys, liver, and intestines. Past the abdominal aorta, the artery branches into two separate iliac arteries, one for each leg, which are responsible for supplying oxygenated blood to the legs and lower half of the body.

## Congenital Heart Disease

Congenital heart disease is one or more problems with the heart's structure that exist since birth. Congenital means that you're born with the condition. Congenital heart disease in adults and children can change the way blood flows through the heart. Some types of congenital heart disease may be mild. But complex defects may cause life-threatening complications. However, advances in diagnosis and treatment continue

to improve survival for those with congenital heart disease. People with congenital heart disease need lifelong medical care. Treatment may include regular checkups (watchful waiting), medications or surgery.

## Single Ventricle Defect

A single ventricle defect (SVD) is a type of heart defect that a child is born with. It occurs when one of the two pumping chambers in the heart, called ventricles, isn't large enough or strong enough to work correctly. In some cases, the chamber might be missing a valve. Single ventricle defects are rare, affecting only about five out of 100,000 newborns. They are also one of the most complex heart problems, usually requiring at least one surgery. There are several types of single ventricle defects which include but are not limited to: tricuspid atresia, hypoplastic left heart syndrome (HLHS), mitral valve atresia (usually associated with HLHS), single left ventricle, double inlet left ventricle (DILV), double outlet right ventricle (DORV), pulmonary atresia with intact ventricular septum (PA/IVS), Ebstein's anomaly, and atrioventricular canal defect (AV Canal).

## Fontan

The Fontan procedure is a type of open-heart surgery. Children who need this surgery usually have it when they're 18 - 36 months old. The Fontan procedure is done for children who are born with heart problems like hypoplastic left heart syndrome (HLHS), tricuspid atresia, and double outlet right ventricle, and depending on the heart problem, children may need the Norwood procedure and Glenn procedure before the Fontan surgery.

After the Fontan procedure, the blood from the lower body goes directly to the lungs. The blood with high oxygen goes into the heart. This way the single ventricle only pumps blood to the body and only pumps blood with high oxygen to the body. There is no more mixing of oxygen-rich blood and oxygen-poor blood.

During the Fontan procedure, the surgeon first disconnects the inferior vena cava (IVC) from the heart and connects it to the pulmonary artery using a conduit (tube). Then, the surgeon makes a small hole between the conduit and the right atrium. This hole (or fenestration) lets some blood still flow back to the heart. It prevents too much blood from flowing to the lungs right away so they have time to adjust. Doctors can close the fenestration later by doing a cardiac catheterization procedure.

# Clinical Data

## General Patient Data

Age (yrs)	17
Sex	Female

## Specific Patient Data

BSA (m <sup>2</sup> )	1.55
CI (L/min/m <sup>2</sup> )	2.3
P IVC MP cath	18
P SVC MP cath	18
P LPA MP cath	17
P RPA MP cath	17
P aorta SP cath	102
P aorta DP cath	67
P aorta MP cath	78

## Notes

Model of aorta on patients who have undergone the Fontan procedure without aortic reconstruction. See [DOI](#) for more details. See below for information on the image data and boundary conditions associated with the model.

**Image Modality:** MR

**Image Type:** DICOM

**Image Source:** TLAB

**Image Manufacturer:** GE MEDICAL SYSTEMS

**Boundary Conditions:** Refer to boundary conditions in the SimVascular file.

## Publications

See the following publications which include the featured model for more details:

Marsden, A. L., Reddy, V. M., Shadden, S. C., Chan, F. P., Taylor, C. A., & Feinstein, J. A. (2010). A new multiparameter approach to computational simulation for Fontan

assessment and redesign. *Congenital Heart Disease*, 5(2), 104-117.  
<http://www.doi.org/10.1111/j.1747-0803.2010.00383.x>

# License

Copyright (c) Stanford University, the Regents of the University of California, Open Source Medical Software Corporation, and other parties.

All Rights Reserved.

Permission is hereby granted, free of charge, to any person obtaining a copy of this data to use the data for research and development purposes subject to the following conditions:

The above copyright notice and the README-COPYRIGHT file shall be included in all copies of any portion of this data. Whenever reasonable and possible in publications and presentations when this data is used in whole or part, please include an acknowledgement similar to the following:

"The data used herein was provided in whole or in part with Federal funds from the National Library of Medicine under Grant No. R01LM013120, and the National Heart, Lung, and Blood Institute, National Institutes of Health, Department of Health and Human Services, under Contract No. HHSN268201100035C"

AND/OR

N.M. Wilson, A.K. Ortiz, and A.B. Johnson, "The Vascular Model Repository: A Public Resource of Medical Imaging Data and Blood Flow Simulation Results," J. Med. Devices 7(4), 040923 (Dec 05, 2013) doi:10.1115/1.4025983.

AND/OR

Reference the official website for this data: [www.vascularmodel.com](http://www.vascularmodel.com)

THE DATA IS PROVIDED "AS IS", WITHOUT WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE AND NONINFRINGEMENT. IN NO EVENT SHALL THE AUTHORS OR COPYRIGHT HOLDERS BE LIABLE FOR ANY CLAIM, DAMAGES OR OTHER LIABILITY, WHETHER IN AN ACTION OF CONTRACT, TORT OR OTHERWISE, ARISING FROM, OUT OF OR IN CONNECTION WITH THE DATA OR THE USE OR OTHER DEALINGS IN THE DATA.