

Drug Facts	
Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg.....	Pain reliever aid
*nonsteroidal anti-inflammatory drug	
Use	
<ul style="list-style-type: none"> treats migraine 	
Warnings	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> hives facial swelling asthma (wheezing) shock <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 2 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed <p>Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.</p>	
Do not use	
<ul style="list-style-type: none"> if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist 	
Ask a doctor before use if	
<ul style="list-style-type: none"> you have never had migraines diagnosed by a health professional you have a headache that is different from your usual migraines you have the worst headache of your life you have fever and stiff neck you have headaches beginning after or caused by head injury, exertion, coughing or bending you experienced your first headache after the age of 50 you have daily headaches you have a migraine so severe as to require bed rest you have liver disease stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic you have asthma you have problems or serious side effects from taking pain relievers or fever reducers you have vomiting with your migraine headache 	
Ask a doctor or pharmacist before use if you are	
<ul style="list-style-type: none"> taking a prescription drug for: <ul style="list-style-type: none"> anticoagulation (thinning of the blood) diabetes gout arthritis under a doctor's care for any serious condition taking any other drug taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> an allergic reaction occurs. Seek medical help right away. you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> feel faint vomit blood have bloody or black stools have stomach pain that does not get better your migraine is not relieved or worsens after first dose new or unexpected symptoms occur ringing in the ears or loss of hearing occurs 	
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.	
Keep out of reach of children.	
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
<ul style="list-style-type: none"> do not use more than directed adults: take 2 caplets with a glass of water if symptoms persist or worsen, ask your doctor do not take more than 2 caplets in 24 hours, unless directed by a doctor under 18 years of age: ask a doctor 	
Other information	
<ul style="list-style-type: none"> store at 20°-25°C (68°-77°F) read all product information before using. Keep this box for important information. 	
Inactive ingredients	
carnauba wax, crospovidone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide	
Questions or comments? 1-800-468-7746	