

ACCEPTED MANUSCRIPT

Three-component soliton states in spinor $F = 1$ Bose-Einstein condensates with PT-symmetric generalized Scarf-II potentials

To cite this article before publication: Zhang-Lei Han *et al* 2024 *Commun. Theor. Phys.* in press <https://doi.org/10.1088/1572-9494/ad8c27>

Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2024 Institute of Theoretical Physics CAS, Chinese Physical Society and IOP Publishing. All rights, including for text and data mining, AI training, and similar technologies, are reserved. Printed in China.



During the embargo period (the 12 month period from the publication of the Version of Record of this article), the Accepted Manuscript is fully protected by copyright and cannot be reused or reposted elsewhere.

As the Version of Record of this article is going to be / has been published on a subscription basis, this Accepted Manuscript will be available for reuse under a CC BY-NC-ND 3.0 licence after the 12 month embargo period.

After the embargo period, everyone is permitted to use copy and redistribute this article for non-commercial purposes only, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by-nc-nd/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected, unless specifically stated otherwise in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

Three-component soliton states in spinor $F = 1$ Bose-Einstein condensates with PT-symmetric generalized Scarf-II potentials

Journal:	<i>Communications in Theoretical Physics</i>
Manuscript ID	CTP-240350.R2
Manuscript Type:	Paper
Date Submitted by the Author:	22-Sep-2024
Complete List of Authors:	Han, Zhang-Lei; China Agricultural University, College of Science Lao, Jun-Yi; China Agricultural University, College of Science Zhang, Jia-Rui; China Agricultural University, College of Science Shen, Yu-Jia; China Agricultural University, College of Science
Keywords:	Three-component soliton states, PT-symmetric generalized Scarf-II potentials, Bose-Einstein condensates, Three-coupled nonlinear Gross-Pitaevskii equations

SCHOLARONE™
Manuscripts

Accepted Manuscript

Three-component soliton states in spinor $F = 1$ Bose-Einstein condensates with \mathcal{PT} -symmetric generalized Scarf-II potentials

Zhang-Lei Han¹, Jun-Yi Lao¹, Jia-Rui Zhang¹, and Yu-Jia Shen^{1,*}

¹College of Science, China Agricultural University, Beijing 100083, China

Introducing \mathcal{PT} -symmetric generalized Scarf-II potentials into the three-coupled nonlinear Gross-Pitaevskii equations offers a new way to seeking stable soliton states in quasi-one-dimensional spin-1 Bose-Einstein condensates. In scenarios where the spin-independent parameter c_0 and the spin-dependent parameter c_2 vary, we use both analytical and numerical methods to investigate the three-coupled nonlinear Gross-Pitaevskii equations with \mathcal{PT} -symmetric generalized Scarf-II potentials. We obtain analytical soliton states and find that simply modulating c_2 may change the analytical soliton states from unstable to stable. Additionally, we obtain numerically stable double-hump soliton states propagating in the form of periodic oscillations, exhibiting distinct behavior in energy exchange. For further investigation, we discuss the interaction of numerical double-hump solitons with Gaussian solitons and observe the transfer of energy among the three components. These findings may contribute to a deeper understanding of solitons in Bose-Einstein condensates with \mathcal{PT} -symmetric potentials and may hold significance for both theoretical understanding and experimental design in related physics experiments.

Keywords: Three-component soliton states; \mathcal{PT} -symmetric generalized Scarf-II potentials; Bose-Einstein condensates; Three-coupled nonlinear Gross-Pitaevskii equations.

I. INTRODUCTION

The phenomenon of a marked increase in wave-like behavior occurs in atomic gases as temperatures decrease, leading to special physical properties such as superfluidity [1–3], superconductivity [4, 5], and quantum entanglement [6–8], is referred to as Bose-Einstein condensates (BECs). With the observation of spinor BECs in the gas of spin-1 Na²³ atoms [9], interests in related theories are sparked, which drive the exploration of various new phenomena, including spin configurations [10, 11], quantum magnetization [12, 13], and the second-order Zeeman effect [14–16]. In recent years, methods involving time-dependent nonlinearity manipulated by Feshbach-resonance techniques have sparked research interest [17–20]. For instance, Feshbach-resonance is utilized to control scattering lengths, and numerical methods are employed to establish the complete stability of solitons [17]. Simultaneously, physical techniques enable control over the collapse and expansion in BECs [18]. Furthermore, methods for modulation of BEC systems involving external potentials with spatiotemporal dependencies have also been investigated [21–23]. Specifically, numerous studies focus on solitons in BECs or spinor BECs, with a primary focus on scenarios that only involve real external potential while a fraction further introduce gain-and-loss distribution manifested as imaginary

* Corresponding author: yjshen2018@cau.edu.cn

2

component [24–26]. Moreover, there has been relatively scant research on modulating stable solitons in spinor BECs based on the concept of \mathcal{PT} -symmetry [27–30], i.e. external potential and gain-and-loss distribution must satisfy the parity conditions. \mathcal{PT} -symmetry was first proposed by Bender et al. in 1998, and it has found significant applications in various fields including optics [31–34], matter waves [35, 36], magnetics [37, 38], acoustics [39, 40], and electronics [41, 42]. In dissipative systems, the \mathcal{PT} -symmetry of complex potentials is a necessary condition for the existence of soliton families [43]. Despite the presence of gain and loss, the linear spectrum of the system can still be entirely real [69], allowing all linear modes to exhibit regular wave behavior as in conservative systems. Various \mathcal{PT} -symmetric potentials have been applied to BEC research since Klaiman *et al.* proposed considering BECs in double-well potentials in 2008 [44], leading to an increasing number of studies on solitons in BECs [45–47].

The spinor BECs has $2F + 1$ components. Governed by the variational principle, the evolution of the wave function in quasi-one-dimensional spin-1 BECs can be expressed using the three-coupled nonlinear Gross-Pitaevskii equations (GPEs) in the following form [48–52]:

$$\begin{aligned} i\hbar\partial_\tau\Psi_+ &= -\frac{\hbar^2}{2m}\partial_\zeta^2\Psi_+ + \hat{V}_+(\zeta)\Psi_+ + (b_0 + b_2)(|\Psi_+|^2 + |\Psi_0|^2)\Psi_+ + (b_0 - b_2)|\Psi_-|^2\Psi_+ + b_2\Psi_0^2\Psi_*, \\ i\hbar\partial_\tau\Psi_0 &= -\frac{\hbar^2}{2m}\partial_\zeta^2\Psi_0 + \hat{V}_0(\zeta)\Psi_0 + (b_0 + b_2)(|\Psi_+|^2 + |\Psi_-|^2)\Psi_0 + b_0|\Psi_0|^2\Psi_0 + 2b_2\Psi_0^*\Psi_+\Psi_-, \\ i\hbar\partial_\tau\Psi_- &= -\frac{\hbar^2}{2m}\partial_\zeta^2\Psi_- + \hat{V}_-(\zeta)\Psi_- + (b_0 + b_2)(|\Psi_0|^2 + |\Psi_-|^2)\Psi_- + (b_0 - b_2)|\Psi_+|^2\Psi_- + b_2\Psi_0^2\Psi_+, \end{aligned} \quad (1)$$

where $\Psi = (\Psi_+, \Psi_0, \Psi_-)^T$, $\Psi_j(\zeta, \tau)$ ($j = +, 0, -$) is the wave function of ζ, τ which describe the transverse coordinate and propagation time respectively. And the asterisk stands for the complex conjugate, \hbar is the reduced Planck constant, m is the atomic mass and $\hat{V}_j(\zeta)$ ($j = +, 0, -$) stands for the real external trap potential. Here, the mean-field coupling constants b_0 and b_2 , which denote the interactions among the three components $\Psi_j(\zeta, \tau)$ ($j = +, 0, -$) responsible for spin-independent and spin-dependent collisions between identical spin-1 bosons, taking the form $b_0 = \pi\hbar^2(a_0 + 2a_2)/3m$ and $b_2 = \pi\hbar^2(a_2 - a_0)/3m$ where a_0 and a_2 represent the s-wave scattering lengths within the symmetric channels characterized by the total spin of the colliding atoms, $F = 0$ and $F = 2$, respectively [53–55].

In this paper, considering the transformation $(\Psi_+, \Psi_0, \Psi_-) \rightarrow (\psi_+, \sqrt{2}\psi_0, \psi_-)$, $x = \zeta\sqrt{2m}/\hbar$, $t = \tau/\hbar$, the dimensionless three-component GPEs are formed as:

$$\begin{aligned} i\partial_t\psi_+ &= -\partial_x^2\psi_+ + U_+(x)\psi_+ + (c_0 + c_2)(|\psi_+|^2 + 2|\psi_0|^2)\psi_+ + (c_0 - c_2)|\psi_-|^2\psi_+ + 2c_2\psi_0^2\psi_*, \\ i\partial_t\psi_0 &= -\partial_x^2\psi_0 + U_0(x)\psi_0 + (c_0 + c_2)(|\psi_+|^2 + |\psi_-|^2)\psi_0 + 2c_0|\psi_0|^2\psi_0 + 2c_2\psi_0^*\psi_+\psi_-, \\ i\partial_t\psi_- &= -\partial_x^2\psi_- + U_-(x)\psi_- + (c_0 + c_2)(|\psi_-|^2 + 2|\psi_0|^2)\psi_- + (c_0 - c_2)|\psi_+|^2\psi_- + 2c_2\psi_0^2\psi_+, \end{aligned} \quad (2)$$

where $U_j(x) = V_j(x) + iW_j(x)$ ($j = +, 0, -$) is the \mathcal{PT} -symmetric generalized Scarf-II potential [56, 57] which can modulate the degree of the energy distribution of matter-wave solitons presented under the external potential well and $W_j(x)$ ($j = +, 0, -$) represents the gain-and-loss distribution. Since the introduction of the concept of \mathcal{PT} -symmetry, achieving a balance between gain and loss in quantum systems has been a long-standing goal in quantum

mechanics, and it has only recently been realized experimentally [58]. Such \mathcal{PT} -symmetric Scarf-II potentials in a spinor BEC can possibly be realized by creating arbitrary optical potentials in time average [59] and also the experimental realization of localized loss [60–62] and of localized gain [63]. The concrete forms of the \mathcal{PT} -symmetric generalized Scarf-II potential are shown as:

$$V_j(x) = v_j \operatorname{sech}^2(x) + \alpha_j \operatorname{sech}^{2n}(x), \quad (3)$$

$$W_j(x) = w_j \operatorname{sech}(x) \tanh(x), \quad j = +, 0, -,$$

with v_j, α_j, w_j ($j = +, 0, -$), and $n > 0$ being all real-valued potential parameters, which can modulate the structure of \mathcal{PT} -symmetric potential. Clearly, when $\alpha_j = 0$ or $n = 1$ ($v_j + \alpha_j \neq 0, j = +, 0, -$), the complex potential (3) transforms into the classical Scarf-II potential [64–66]. As the case of $c_0 = c_2 < 0$ has been investigated in the references [67, 68], yielding analytical solutions of Eqs. (1) under \mathcal{PT} -symmetric Scarf-II potential and \mathcal{PT} -symmetric harmonic-Hermitian-Gaussian potential, along with results on different types of soliton collisions, this paper primarily focuses on the scenarios where $c_2 = 0.1$ (antiferromagnetic) and $c_2 = -0.5$ (ferromagnetic) under \mathcal{PT} -symmetric generalized Scarf-II potentials.

The subsequent sections of this paper are organized as follows. In Section II, we explore the soliton states of Eqs. (2), obtaining its families of analytical soliton states under \mathcal{PT} -symmetric generalized Scarf-II potentials. We conduct linear stability analysis, distinguishing stable and unstable regions. The results are validated through dynamic evolution with 5% perturbations. In Section III, we utilize the power-conserving squared-operator methods [69] to generate families of numerical soliton states for Eqs. (2). A detailed investigation of points on the curves is carried out, obtaining the evolution of corresponding stable and unstable numerical solitons. Some analytical soliton states, numerical single-hump soliton states, and numerical double-hump soliton states are chosen to be interacted with exotic Gaussian solitons. Finally, our conclusions are presented in Section IV.

II. ANALYTICAL SOLITON STATES

With the stationary states transformation $\psi_j(x, t) = \phi_j(x)e^{i\mu_j t}$ ($j = +, 0, -$), the three-component GPEs can be derived as

$$\begin{aligned} -\mu_+ \phi_+ + \phi_+'' - [V_+(x) + iW_+(x)]\phi_+ - (c_0 + c_2)(|\phi_+|^2 + 2|\phi_0|^2)\phi_+ - (c_0 - c_2)|\phi_-|^2\phi_+ - 2c_2\phi_0^2\phi_-^* &= 0, \\ -\mu_0 \phi_0 + \phi_0'' - [V_0(x) + iW_0(x)]\phi_0 - (c_0 + c_2)(|\phi_+|^2 + |\phi_-|^2)\phi_0 - 2c_0|\phi_0|^2\phi_0 - 2c_2\phi_0^*\phi_+\phi_- &= 0, \\ -\mu_- \phi_- + \phi_-'' - [V_-(x) + iW_-(x)]\phi_- - (c_0 + c_2)(|\phi_-|^2 + 2|\phi_0|^2)\phi_- - (c_0 - c_2)|\phi_+|^2\phi_- - 2c_2\phi_0^2\phi_+^* &= 0, \end{aligned} \quad (4)$$

where μ_j ($j = +, 0, -$) represents the real **chemical potential**. For obtaining the soliton states of Eqs. (4), we utilize the constraints $w_+ + w_- = 2w_0$, $\mu_+ = \mu_0 = \mu_- = n^2$, $A_0^2 = A_+A_-$, $\alpha_+ = \alpha_0 = \alpha_-$,

$$v_j = -\left(\frac{w_j}{2n+1}\right)^2 - n^2 - n, \quad j = +, 0, -,$$

1
2
3
4
5
and derive

$$\begin{aligned} \phi_j(x) &= A_j \operatorname{sech}^n(x) e^{i\theta_j(x)}, \\ \theta_j(x) &= -\frac{2w_j}{2n+1} \arctan(\tanh \frac{x}{2}), \quad j = +, 0, -, \\ A_+ + A_- &= \sqrt{\frac{\alpha_+}{-(c_0 + c_2)}}. \end{aligned} \quad (5)$$

15 To demonstrate the stability of soliton states, we perform the linear stability analysis on the obtained soliton
16 solutions $\psi_j(x, t)$ ($j = +, 0, -$) of Eqs (2). Substituting the perturbed solutions ($\epsilon \ll 1$)
17

$$\psi_j(x, t) = \phi_j(x) e^{i\mu_j t} + \epsilon [f_j(x) e^{i\delta t} + g_j^*(x) e^{-i\delta^* t}] e^{i\mu_j t}, \quad j = +, 0, -, \quad (6)$$

21 into Eqs. (2) yields an eigenvalue problem $\mathbf{L}\mathbf{Y} = \delta\mathbf{Y}$, where δ stands for the eigenvalue, $f_j(x)$, $g_j(x)$ ($j = +, 0, -$)
22 are components of the corresponding eigenfunction, $\mathbf{Y} = (f_1, g_1, f_2, g_2, f_3, g_3)^T$, and the specific form of \mathbf{L} is shown
23 in Appendix.

26 For the given eigenvalue problem, we discretize the differential operator into a matrix using spectral methods [70,
27 71] and obtain the numerical values for δ . Considering the inherent errors in numerical methods, when the maximum
28 of $|\operatorname{Im}(\delta)|$ is sufficiently small ($|\operatorname{Im}(\delta)| < 10^{-5}$), δ can be approximated as a real number. In this case, the solutions
29 $\psi_j(x, t)$ ($j = +, 0, -$) can be considered linearly stable. This theoretical assumption is further validated through
30 numerical evolution with 5% perturbations.
31

34 As a result, when we choose the system parameters as $c_0 = 1, c_2 = -0.5$, the curve of analytical soliton family for
35 n concerning the atomic densities of soliton can be obtained, as illustrated in Fig. 1(a1). The atomic densities is
36 defined as $P_j = \int |\psi_j(x)|^2 dx$ ($j = +, 0, -$), and the energy flux can be described by introducing the Poynting vector
37 $S_j = i(\phi_j \phi_{j,x}^* - \phi_j^* \phi_{j,x})/2$ ($j = +, 0, -$) [72]. Due to constraints on the system parameters for analytical soliton
38 states, only scenarios under the special condition where $P_+ = P_0 = P_- = P$ are studied in this paper. Through
39 linear stability analysis, we distinguish the stable and unstable portions of the curve. The unstable regions are
40 indicated by the red dashed lines, while the stable regions are represented by the blue solid lines. We select one
41 point from each of the stable and unstable sections on the curve for further investigation. Point A in Fig. 1(a1)
42 is selected from the unstable region, corresponding to $P = 1.6000$ and $n = 0.5$. In Fig. 1(b1)-(b3), we present
43 its evolutionary results after undergoing 5% perturbations as initial conditions and the corresponding eigenvalue
44 distribution is reflected in Fig. 1(b4). Point B in Fig. 1(a1) is chosen from the stable region, with $P = 0.8000$ and
45 $n = 1.5$. Similarly, in Fig. 1(c1)-(c4), we illustrate its evolutionary results with 5% perturbations, along with the
46 corresponding eigenvalue distribution. Clearly, the soliton states corresponding to point A is unstable, while that
47 of point B is stable, confirming the accuracy of linear stable analysis. Moreover, we observe that as the value of n
48 increases from 0.5 to 1.5, the atomic density decreases by just half, with α_j remaining constant and $|v_j|$ ($j = +, 0, -$)
49 increasing. This suggests that higher values of $|v_j|$ ($j = +, 0, -$) may correspond to more stable soliton states.
50
51
52
53
54
55
56
57
58
59
60




Fig. 1: Families of analytical soliton states, the progression of stable/unstable analytical solitons and their eigenvalues for linear stability, associated with $c_2 = -0.5$ and $c_2 = 0.1$ respectively. The parameters are chosen as: $c_0 = 1$, $w_j = 0.6$, $A_j = 0.5$ ($j = +, 0, -$). (b1)-(b4) correspond to point A in (a1) where $v_j = -0.8400$, $\alpha_j = -0.5$ ($j = +, 0, -$) and $P = 1.6000$, $n = 0.5$. (c1)-(c4) correspond to point B in (a1) where $v_j = -3.7725$, $\alpha_j = -0.5$ and $P = 0.8000$, $n = 1.5$. (d1)-(d4) correspond to point C in (a2) where $v_j = -0.8400$, $\alpha_j = -1.1$ ($j = +, 0, -$) and $P = 1.6000$, $n = 0.5$.

When the value of c_2 is changed from -0.5 to 0.1 , the curve of the obtained analytical solution family exhibit the same shape and position, but the stable region is significantly enlarged, as shown in Fig. 1(a2). From the curve in Fig. 1(a2), we select point C with the same coordinates as point A for further investigation. In Fig. 1(d1)-(d4), we present its evolutionary results after undergoing 5% perturbations as initial conditions and the corresponding eigenvalue distribution. However, the evolution corresponding to point C is stable, indicating that increasing c_2 from -0.5 to 0.1 can decrease the value of α_j , and smaller α_j ($j = +, 0, -$) values may correspond to stable solitons.

III. NUMERICAL SOLITON STATES

Obtaining the analytical solution of Eqs. (2) demands stringent system parameter specifications, rendering the implementation of BECs experiments challenging. Consequently, to surmount these limitations and manipulate the propagation coefficient, numerical methods are employed in our research endeavors.

In our numerical investigation, we first focus on the case when $c_0 = 1$ and $c_2 = -0.5$. With the system parameters chosen as $w_j = 2.5$, $v_j = -3.8$, and $\alpha_j = -0.4$ ($j = +, 0, -$), we present families of single-hump numerical soliton states in Fig. 2(a1)-(a3). Clearly, as n grows, the entire curve relocates towards smaller μ values. We further investigate points D and E selected from the curves shown in Fig. 2(a1)-(a3). Point D corresponds to $P_+ = 3.6600$, $P_0 = 4.6321$, $P_- = 5.3079$, $\mu_+ = 1.8563$, $\mu_0 = 1.8555$, $\mu_- = 1.8571$; point E corresponds to $P_+ = 4.0000$, $P_0 = 5.0624$, $P_- = 4.5376$, $\mu_+ = 1.8724$, $\mu_0 = 1.7685$, $\mu_- = 2.0890$. By performing linear stability analysis on the soliton states corresponding to these three points, we ascertain that the soliton states corresponding to point D is stable, and the stability of the soliton states corresponding to point E is unstable. Similarly, we validate these findings by conducting numerical evolution with 5% perturbations.

Next, we investigate the numerical families of double-hump soliton states when $c_0 = 1$ and $c_2 = 0.1$. Given the constancy of other conditions, the numerical double-hump solution families across various n values exhibit significant overlap, hence in Fig. 3, we only show the scenario for $n = 2$. We change the values of w_j from 2.5 to 0.8 , v_j from -3.8 to -6 , while keeping the values of α_j ($j = +, 0, -$) unchanged. Similarly, we further investigate points F and G selected from the curves shown in Fig. 3(a1)-(a3). Fig. 3(b1)-(b4) and Fig. 3(d1)-(d4) depict the progression of stable soliton states respectively, along with their associated eigenvalue distributions. Of note, as shown in Fig. 3(b1)-(b3), the double-hump soliton states corresponding to point F exhibit periodic oscillations while conserving the energy during propagation, which can be seen as a form of energy exchange between soliton states. Furthermore, compared to previous studies where all three soliton states exhibited periodic oscillations, the phenomenon of energy exchange only existing between ψ_+ and ψ_0 , with ψ_- remaining constant, might be relatively infrequent. To further investigate the oscillatory behavior of the double-hump soliton states, the total number energy flux from the numerical solutions is investigated. Point G from Fig. 3(a1)-(a3) is selected for study, compared to point F, the corresponding oscillation period is longer, and ψ_- also begins to exhibit slight oscillations, as shown in Fig. 3(c1)-(e4). This suggests that the oscillation of the double-hump soliton states may be related to the magnitude of the energy and indicates the dynamic stability of the double-hump soliton against small perturbations.




Fig. 2: Families of numerical single-hump soliton states, the progression of stable/unstable numerical single-hump solitons and their eigenvalues for linear stability, associated with different n . The parameters are chosen as: $c_0 = 1$, $c_2 = -0.5$, $w_+ = w_0 = w_- = 2.5$, $v_+ = v_0 = v_- = -3.8$, $\alpha_+ = \alpha_0 = \alpha_- = -0.4$.

Next, we turn our attention to studying the interaction between Gaussian solitons and the solitons depicted in Fig. 1 to Fig. 3. We consider the following initial conditions

$$\overline{\psi_j}(x, 0) = \psi_j(x, 0) + q_j e^{-0.05(x+23)^2 + 3ix} + q_j e^{-0.05(x-23)^2 - 3ix}, \quad j = +, 0, -, \quad (7)$$

and

$$\overline{\psi_j}(x, 0) = \psi_j(x, 0) + q_j e^{-0.05(x+23)^2 + 4ix} + q_j e^{-0.05(x-23)^2 - 4ix}, \quad j = +, 0, -, \quad (8)$$

where q_j is the maximum amplitude of the exotic Gaussian soliton. Fig. 4(a1)-(a3) correspond to the case of analytical solitons with $c_0 = 1$, $c_2 = -0.5$, $n = 0.5$, $w_j = 0.6$, $A_j = 0.5$, $P = 1.6000$, $v_j = -3.7725$, and $\alpha_j = -0.5$ ($j = +, 0, -$), satisfying the initial condition (7); Fig. 4(b1)-(b3) correspond to the case of numerical single-hump

solitons (point D) with $P_+ = 3.6600$, $P_0 = 4.6321$, $P_- = 5.3079$, $\mu_+ = 1.8563$, $\mu_0 = 1.8555$, $\mu_- = 1.8571$, $\mu_0 = 1.7329$, $\mu_- = 1.7262$, and $n = 0.5$, satisfying the initial condition (7); Fig. 4(c1)-(c3) correspond to the case of numerical double-hump solitons (point F) with $P_+ = 0.0400$, $P_0 = 0.0506$, $P_- = 13.5547$, $\mu_+ = 0.7273$, $\mu_0 = 0.6493$, $\mu_- = 0.6556$, and $n = 2$, satisfying the initial condition (7); Fig. 4(d1)-(d3) correspond to the case of numerical double-hump solitons (point G) with $P_+ = 0.3200$, $P_0 = 0.4050$, $P_- = 12.8750$, $\mu_+ = 0.7083$, $\mu_0 = 0.6319$, $\mu_- = 0.6497$, and $n = 2$, satisfying the initial condition (8). For Fig. 4(a1)-(a3), we choose $q_+ = q_0 = q_- = 0.2$, for Fig. 4(b1)-(b3), we choose $q_+ = q_0 = 0.05$, $q_- = 0.3$, while for Fig. 4(d1)-(d3), we choose $q_+ = q_0 = 0.1$, $q_- = 0.5$. We notice that after the interaction between the stable soliton states and the exotic Gaussian solitons, both of them maintain their original shapes and continue to propagate forward, as shown in Fig. 4(a1)-(c3). This indicates that an elastic collision occurs between the soliton states and the exotic Gaussian solitons. Moreover, some studies suggest that interactions between solitons may trigger energy flow among the three components. However as shown in Fig. 4(d1)-(d2), we observe a specific scenario where the energy of ψ_+ decreases, and the energy of ψ_0 increases, while the energy of ψ_- remains unchanged after the interaction between the exotic Gaussian solitons and the double-hump solitons. This can be regarded as a special form of energy flow involving only two components. In order to explore more properties about the interaction, the corresponding energy flux are checked. As illustrated in Fig. 5, the Poynting vectors of $\psi_j(x, t)$ all keep invariant. Moreover, the energy flux of the soliton states has the same characteristics as that $\psi_j(x, t)$ propagate, which further verifies the correctness of Fig. 4.

IV. CONCLUSIONS

In conclusion, we introduce \mathcal{PT} -symmetric generalized Scarf-II potentials into three-coupled Gross-Pitaevskii equations with two different scenarios of inter-component interaction parameter values c_0 and c_2 , and obtain corresponding analytical/numerical soliton states and soliton families. In both scenarios of parameter configurations, the analytical soliton families exhibit identical curve shapes, but the stable region of the analytical soliton family curves significantly increases under the condition of $c_2 = 0.1$. With the condition of $c_2 = -0.5$ (ferromagnetic), the single-hump solution families obtained through iterative algorithms shift towards smaller values of μ as n increases. Besides, in the case of $c_2 = 0.1$ (antiferromagnetic), we obtain point F whose corresponding double-hump soliton states exhibit periodic oscillations while conserving the energy during propagation, within the double-hump soliton families. Notably, the phenomenon of energy exchange only existing between ψ_+ and ψ_0 , with ψ_- remaining constant, is observed. Moreover, after the interaction between the double-hump soliton states and the exotic Gaussian solitons, the energy of ψ_+ decreases and the energy of ψ_0 increases, while the energy of ψ_- unchanged. This phenomenon can be seen as collisions triggering the flow of energy among the three components. These findings could contribute to a deeper understanding of solitons in Bose-Einstein condensates with \mathcal{PT} -symmetric potentials and potentially aid in designing relevant physics experiments.




Fig. 3: Families of numerical double-hump soliton states, the progression of stable numerical double-hump solitons and their energy flux associated with $\text{https://m3t2.manuscriptcentral.com/ctphyss}$. The parameters are chosen as $\mu_1 = 1$, $c_2 = 0.1$, $w_+ = w_0 = w_- = 0.8$, $v_+ = v_0 = v_- = -6$, $\alpha_+ = \alpha_0 = \alpha_- = -0.4$. And their maximum of $|\text{Im}(\delta)|$ for linear stability are all 10^{-7} .




Fig. 4: Interactions of single-hump and double-hump soliton states with exotic Gaussian solitons. (a1)-(a3) correspond to analytical soliton states at $c_0 = 1$, $c_2 = -0.5$, $n = 0.5$, $w_j = 0.6$, $A_j = 0.5$, $P = 1.6000$, $v_j = -3.7725$ and $\alpha_j = -0.5$ ($j = +, 0, -$). (b1)-(b3) correspond to numerical single-hump soliton states at $n = 0.5$. (c1)-(d3) correspond to numerical double-hump soliton states at $n = 2$.




Fig. 5: The energy flux of the interactions of single-hump and double-hump soliton states with exotic Gaussian solitons in Fig. 4. The parameters are chose the same as Fig. 4.

12

ACKNOWLEDGMENTS

We express our sincere thanks to the editor, referees, and all the members of our discussion group for their valuable comments. This work was supported by NSFC under Grant number 12272403, and Beijing Training Program of Innovation (Grant number S202410019024). The funding body plays an important role in the design of the study, in analysis, calculation, and in writing of the manuscript.

DECLARATIONS OF INTEREST

The authors declare that they have no conflict of interest.

DATA AVAILABILITY

The authors declare that the manuscript has no associated data.

FUNDING

This work was supported by NSFC under Grant number 12272403, and Beijing Training Program of Innovation (Grant number S202410019024). The funding body plays an important role in the design of the study, in analysis, calculation, and in writing of the manuscript.

APPENDIX

The matrix \mathbf{L} of linear eigenvalue problem, which is derived during linear stability analysis, has specific form as follows:

$$\mathbf{L} = \begin{pmatrix} L_1 & L_{12} & L_{13} & L_{14} & L_{15} & L_{16} \\ L_{21} & -L_1^* & L_{23} & L_{24} & L_{25} & L_{26} \\ L_{31} & L_{32} & L_2 & L_{34} & L_{35} & L_{36} \\ L_{41} & L_{42} & L_{43} & -L_2^* & L_{45} & L_{46} \\ L_{51} & L_{52} & L_{53} & L_{54} & L_3 & L_{56} \\ L_{61} & L_{62} & L_{63} & L_{64} & L_{65} & -L_3^* \end{pmatrix}, \quad (9)$$

1
2
3
4
5 where
6

$$\begin{aligned}
 L_1 &= \partial_x^2 - 2(c_0 + c_2)\phi_+\phi_+^* - 2(c_0 + c_2)\phi_0\phi_0^* - (c_0 - c_2)\phi_-\phi_-^* - U_+(x) - \mu_+, \\
 L_2 &= \partial_x^2 - (c_0 + c_2)\phi_+\phi_+^* - 4c_0\phi_0\phi_0^* - (c_0 + c_2)\phi_-\phi_-^* - U_0(x) - \mu_0, \\
 L_3 &= \partial_x^2 - (c_0 - c_2)\phi_+\phi_+^* - 2(c_0 + c_2)\phi_0\phi_0^* - 2(c_0 + c_2)\phi_-\phi_-^* - U_-(x) - \mu_-, \\
 L_{12} &= -(c_0 + c_2)\phi_+^2, & L_{14} &= -2(c_0 + c_2)\phi_+\phi_0, \\
 L_{15} &= -(c_0 - c_2)\phi_+\phi_-^*, & L_{21} &= (c_0 + c_2)\phi_+^{*2}, \\
 L_{23} &= 2(c_0 + c_2)\phi_+^*\phi_0^*, & L_{26} &= (c_0 - c_2)\phi_+^*\phi_-, \\
 L_{32} &= -(c_0 + c_2)\phi_+\phi_0, & L_{36} &= -(c_0 + c_2)\phi_0\phi_-, \\
 L_{41} &= (c_0 + c_2)\phi_+^*\phi_0^*, & L_{45} &= (c_0 + c_2)\phi_0^*\phi_-^*, \\
 L_{51} &= -(c_0 - c_2)\phi_+^*\phi_-, & L_{54} &= -2(c_0 + c_2)\phi_0\phi_-, \\
 L_{56} &= -(c_0 + c_2)\phi_-^2, & L_{62} &= (c_0 + c_2)\phi_+\phi_-^*, \\
 L_{63} &= 2(c_0 + c_2)\phi_0^*\phi_-^*, & L_{65} &= (c_0 + c_2)\phi_-^{*2}, \\
 L_{13} &= -2(c_0 + c_2)\phi_+\phi_0^* + 4c_2\phi_0\phi_-^*, & L_{16} &= -(c_0 - c_2)\phi_+\phi_- - 2c_2\phi_0^2, \\
 L_{24} &= 2(c_0 + c_2)\phi_+^*\phi_0 + 4c_2\phi_0^*\phi_-, & L_{25} &= (c_0 - c_2)\phi_+^*\phi_-^* + 2c_2\phi_0^{*2}, \\
 L_{31} &= -(c_0 + c_2)\phi_+^*\phi_0 - 2c_2\phi_0^*\phi_-, & L_{34} &= -2c_0\phi_0^2 - 2c_2\phi_+\phi_-, \\
 L_{35} &= -(c_0 + c_2)\phi_0\phi_-^* - 2c_2\phi_+\phi_0^*, & L_{42} &= (c_0 + c_2)\phi_+\phi_0^* + 4c_2\phi_0\phi_-^*, \\
 L_{43} &= 2c_0\phi_0^{*2} + 2c_2\phi_+^*\phi_-^*, & L_{46} &= (c_0 + c_2)\phi_0^*\phi_- + 2c_2\phi_+^*\phi_0, \\
 L_{52} &= -(c_0 - c_2)\phi_+\phi_- - 2c_2\phi_0^2, & L_{53} &= -2(c_0 + c_2)\phi_0^*\phi_- - 4c_2\phi_+^*\phi_0, \\
 L_{61} &= (c_0 - c_2)\phi_+^*\phi_-^* + 2c_2\phi_0^{*2}, & L_{64} &= 2(c_0 + c_2)\phi_0\phi_-^* + 4c_2\phi_+\phi_0^*.
 \end{aligned}$$

- [1] G. Lerario, A. Fieramosca, F. Barachati, *et al.*, Room-temperature superfluidity in a polariton condensate, *Nat. Phys.* **13** (2017) 837-841.
- [2] D. M. Stamper-Kurn and M. Ueda, Spinor Bose gases: Symmetries, magnetism, and quantum dynamics, *Rev. Mod. Phys.* **85** (2013) 1191.
- [3] A. V. Yulin, Y. V. Bludov and V. V. Konotop, *et al.*, Superfluidity of Bose-Einstein condensates in toroidal traps with nonlinear lattices, *Phys. Rev. A* **84** (2011) 063638.
- [4] A. S. Alexandrov, *d*-wave Bose-Einstein condensate and tunnelling in superconducting cuprates, *Physica C* **305** (1998) 46-56.
- [5] M. Sun, A. V. Parafilo, K. H. A. Villegas, *et al.*, Bose-Einstein condensate-mediated superconductivity in graphene, *2D Mater.* **8** (2021) 031004.
- [6] Y. X. Huang, N. J. Hui and W. F. Liu, *et al.*, Quantum Entanglement in a Spin-Orbit Coupled Bose-Einstein Condensate, *Commun. Theor. Phys.* **64** (2015) 644.
- [7] Z. X. Yu, Z. Y. Jiao and J. Z. Sun, Quantum Effects of Bose-Einstein Condensates, *Commun. Theor. Phys.* **41** (2004) 593.
- [8] H. G. Zhu, J. Shu, G. Q. Huang GQ and C. L. Luo, Coherence and Entanglement in Spinor Bose-Einstein Condensate, *Int. J. Theor. Phys.* **53** (2014) 2803-2809.
- [9] D. M. Stamper-Kurn, M. R. Andrews, A. P. Chikkatur, *et al.*, Optical confinement of a Bose-Einstein condensate, *Phys. Rev. Lett.* **80** (1998) 2027.
- [10] N. V. Hung, P. Szankowski, V. V. Konotop and M. Trippenbach, Four-wave mixing in spin-orbit coupled Bose-Einstein condensates, *New. J. Phys.* **22** (2020) 053019.
- [11] T. Liu and T. C. H. Liew, Spontaneous spin bifurcations in a Bose-Einstein condensate of indirect excitons, *Superlattice. Microst.* **108** (2017) 57-63.
- [12] S. Hoshi and H. Saito, Magnetization of a half-quantum vortex in a spinor Bose-Einstein condensate, *Phys. Rev. A* **78** (2008) 053618.
- [13] J. M. Zhang, S. Cui, H. Jing, *et al.*, Probing the quantum ground state of a spin-1 Bose-Einstein condensate with cavity transmission spectra, *Phys. Rev. A* **80** (2009) 043623.
- [14] T. Isoshima and S. Yip, Effect of quadratic Zeeman energy on the vortex of spinor Bose-Einstein condensates, *J. Phys. Soc. Jpn.* **75** (2006) 074605.
- [15] J. C. Sandoval-Santana, R. Zamora-Zamora, R. Paredes and V. Romero-Rochín, Intrinsic decoherence and recurrences in a large ferromagnetic $F = 1$ spinor Bose-Einstein condensate, *Symmetry* **13** (2021) 67.
- [16] D. Zhao, S. W. Song, L. Wen, *et al.*, Topological defects and inhomogeneous spin patterns induced by the quadratic Zeeman effect in spin-1 Bose-Einstein condensates, *Phys. Rev. A* **91** (2015) 013619.
- [17] P. G. Kevrekidis, G. Theocharis, D. J. Frantzeskakis and B. A. Malomed, Feshbach resonance management for Bose-Einstein condensates, *Phys. Rev. Lett.* **90** (2003) 230401.
- [18] M. Yan, B. J. DeSalvo, B. Ramachandhran, *et al.*, Controlling condensate collapse and expansion with an optical Feshbach resonance, *Phys. Rev. Lett.* **110** (2013) 123201.
- [19] K. Enomoto, K. Kasa, M. Kitagawa and Y. Takahashi, Optical Feshbach resonance using the intercombination transition, *Phys. Rev. Lett.* **101** (2008) 203201.

- [20] Y. J. Shen, Y. T. Gao, D. W. Zuo, *et al.*, Nonautonomous matter waves in a spin-1 Bose-Einstein condensate, *Phys. Rev. E* **89** (2014) 062915.
- [21] X. F. Zhang, X. H. Hu, X. X. Liu and W. M. Liu, Vector solitons in two-component Bose-Einstein condensates with tunable interactions and harmonic potential, *Phys. Rev. A* **79** (2009) 033630.
- [22] S. Rajendran, P. Muruganandam and M. Lakshmanan, Bright and dark solitons in a quasi-1D Bose-Einstein condensates modelled by 1D Gross-Pitaevskii equation with time-dependent parameters, *Physica D* **239** (2010) 366.
- [23] Y. Q. Yao, W. Han, J. Li and W. M. Liu, Localized nonlinear waves and dynamical stability in spinor Bose-Einstein condensates with time-space modulation, *J. Phys. B* **51** (2018) 105001
- [24] D. Dast, D. Haag, H. Cartarius and G. Wunner, Purity oscillations in Bose-Einstein condensates with balanced gain and loss, *Phys. Rev. A* **93** (2016) 033617.
- [25] D. Dast, D. Haag, H. Cartarius and G. Wunner, Quantum master equation with balanced gain and loss, *Phys. Rev. A* **90** (2014) 052120.
- [26] P. Lunt, D. Haag, D. Dast, H. Cartarius and G. Wunner, Balanced gain and loss in Bose-Einstein condensates without \mathcal{PT} -symmetry, *Phys. Rev. A* **96** (2017) 023614.
- [27] C. M. Bender, Making sense of non-Hermitian Hamiltonians, *Rep. Prog. Phys.* **70** (2007) 947.
- [28] C. M. Bender, D. C. Brody and H. F. Jones, Must a hamiltonian be hermitian?, *Am. J. Phys.* **71** (2003) 1095-1102.
- [29] C. M. Bender and S. Boettcher, Real spectra in non-Hermitian Hamiltonians having \mathcal{PT} -symmetry, *Phys. Rev. Lett.* **80** (1998) 5243.
- [30] J. Y. Lao, Z. Y. Qin, J. R. Zhang and Y. J. Shen, Peakons in spinor $F = 1$ Bose-Einstein condensates with \mathcal{PT} -symmetric δ -function potentials, *Chaos Solitons Fractals* **180** (2024) 114497.
- [31] K. G. Makris, R. El-Ganainy, D. N. Christodoulides and Z. H. Musslimani, Beam dynamics in \mathcal{PT} -symmetric optical lattices, *Phys. Rev. Lett.* **100** (2008) 103904.
- [32] S. Longhi, Bloch oscillations in complex crystals with \mathcal{PT} -symmetry, *Phys. Rev. Lett.* **103** (2009) 123601.
- [33] L. Feng, M. Ayache, J. Q. Huang, *et al.*, Nonreciprocal light propagation in a silicon photonic circuit, *Science* **333** (2011) 729-733.
- [34] A. Regensburger, C. Bersch and M. A. Miri, *et al.*, Parity-time synthetic photonic lattices, *Nature* **488** (2012) 167-171.
- [35] H. Cartarius and G. Wunner, Model of a \mathcal{PT} -symmetric Bose-Einstein condensate in a δ -function double-well potential, *Phys. Rev. A* **86** (2012) 013612.
- [36] H. Moritz, K. Tsampikos and O. Alexander, Bifurcations in resonance widths of an open Bose-Hubbard dimer, *Phys. Rev. A* **73** (2006) 063625.
- [37] J. M. Lee, T. Kottos and B. Shapiro, Macroscopic magnetic structures with balanced gain and loss, *Phys. Rev. B* **91** (2015) 094416.
- [38] A. Galda and V. M. Vinokur, Parity-time symmetry breaking in magnetic systems, *Phys. Rev. B* **94** (2016) 020408.
- [39] R. Fleury, D. Sounas and A. Alù, An invisible acoustic sensor based on parity-time symmetry, *Nat. Commun.* **6** (2015) 5905.
- [40] C. Shi, M. Dubois, Y. Chen, *et al.*, Accessing the exceptional points of parity-time symmetric acoustics, *Nat. Commun.* **7** (2016) 11110.
- [41] N. Bender, S. Factor, J. D. Bodyfelt, *et al.*, Observation of asymmetric transport in structures with active nonlinearities, *Phys. Rev. Lett.* **110** (2013) 234101.
- [42] F. Bagarello, From self-adjoint to non-self-adjoint harmonic oscillators: Physical consequences and mathematical pitfalls, *Phys. Rev. A* **88** (2013) 032120.

- [43] J. K. Yang, Necessity of \mathcal{PT} -symmetry for soliton families in one-dimensional complex potentials, *Phys. Lett. A* **378** (2014) 367-373.
- [44] S. Klaiman, G. Uwe and N. Moiseyev, Visualization of branch points in \mathcal{PT} -symmetric waveguides, *Phys. Rev. Lett.* **101** (2008) 080402.
- [45] Y. X. Xu, Y. Zhao, Q. H. Huang, et al., Rydberg-dressed solitons in Bose-Einstein condensates with parity-time symmetry, *Results in Physics* **49** (2023) 106534.
- [46] Y. V. Kartashov, V. V. Konotop and F. Kh. Abdullaev, Gap solitons in a spin-orbit-coupled Bose-Einstein condensate, *Phys. Rev. Lett.* **111** (2013) 060402.
- [47] B. Sanjana, G. Amit, J. Soumendu and C. N. Kumar, Stationary hypergeometric solitons and their stability in a Bose-Einstein condensate with \mathcal{PT} -symmetric potential, *Phys. Lett. A* **469** (2023) 128751.
- [48] T. L. Ho, Spinor Bose condensates in optical traps, *Phys. Rev. Lett.* **81** (1998) 742.
- [49] Y. Kawaguchi and M. Ueda, Spinor Bose-Einstein condensates, *Phys. Rep.* **520** (2012) 253-381.
- [50] T. Ohmi and K. Machida, Bose-Einstein condensation with internal degrees of freedom in alkali atom gases, *J. Phys. Soc. Jpn.* **67** (1998) 1822-1825.
- [51] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*. Cambridge University Press, Cambridge (2008).
- [52] Z. H. Zhang, C. Zhang, S. J. Yang and S. Feng, Analytical solutions to the spin-1 Bose-Einstein condensates, *J. Phys. B* **45** (2012) 215302.
- [53] B. J. Dabrowska-Wüster, E. A. Ostrovskaya, T. J. Alexander and Y. S. Kivshar, Multicomponent gap solitons in spinor Bose-Einstein condensates, *Phys. Rev. A* **75** (2007) 023617.
- [54] J. P. Martikainen, A. Collin and K. A. Suominen, Coreless vortex ground state of the rotating spinor condensate, *Phys. Rev. A* **66** (2002) 053604.
- [55] H. E. Nistazakis, D. J. Frantzeskakis and P. G. Kevrekidis, Bright-dark soliton complexes in spinor Bose-Einstein condensates, *Phys. Rev. A* **77** (2008) 033612.
- [56] Y. Chen, Z. Y. Yan and B. A. Malomed, Higher-dimensional soliton generation, stability and excitations of the \mathcal{PT} -symmetric nonlinear Schrödinger equations, *Physica D* **430** (2022) 133099.
- [57] Y. Chen, Z. Y. Yan and Mihalache D, Soliton formation and stability under the interplay between parity-time-symmetric generalized Scarf-II potentials and Kerr nonlinearity, *Phys. Rev. E* **102** (2020) 012216.
- [58] S. Altinisik, D. Dizdarevic, and J. Main, Balanced gain and loss in spatially extended non- \mathcal{PT} -symmetric multiwell potentials, *Phys. Rev. A* **100** (2019) 063639.
- [59] K. Henderson1, C. Ryu1, C. MacCormick and M. G. Boshier, Experimental demonstration of painting arbitrary and dynamic potentials for Bose-Einstein condensates, *New J. Phys.* **11** (2009) 043030.
- [60] T. Gericke, P. Würtz, D. Reitz, et al., High-resolution scanning electron microscopy of an ultracold quantum gas, *Nature Phys* **4** (2008) 949-953.
- [61] P. Würtz, T. Langen, T. Gericke, et al., Experimental Demonstration of Single-Site Addressability in a Two-Dimensional Optical Lattice, *Phys. Rev. Lett.* **103** (2009) 080404.
- [62] G. Barontini, R. Labouvie, F. Stubenrauch, et al., Controlling the Dynamics of an Open Many-Body Quantum System with Localized Dissipation, *Phys. Rev. Lett.* **110** (2013) 035302.
- [63] D. Döring, G. R. Dennis, N. P. Robins, et al., Pulsed pumping of a Bose-Einstein condensate, *Phys. Rev. A* **79** (2009) 063630.
- [64] Y. J. Shen, Z. C. Wen, Z. Y. Yan and C. Hang, Effect of \mathcal{PT} -symmetry on nonlinear waves for three-wave interaction models in the quadratic nonlinear media, *Chaos* **28** (2018) 043104.

- [65] Z. Ahmed, Real and complex discrete eigenvalues in an exactly solvable one-dimensional complex \mathcal{PT} -invariant potential, Phys. Lett. A **282** (2001) 343.
- [66] Z. H. Musslimani, K. G. Makris, R. El-Ganainy and D. N. Christodoulides, Optical solitons in \mathcal{PT} periodic potentials, Phys. Rev. Lett. **100** (2008) 030402.
- [67] J. J. Qi, D. Zhao and W. M. Liu, Soliton collisions in spin-orbit coupled spin-1 Bose-Einstein condensates, J. Phys. A: Math. Theor. **56** (2023) 255702.
- [68] M. Zhong, Z. Y. Yan and S. F. Tian, Stable matter-wave solitons, interactions, and excitations in the spinor $F = 1$ Bose-Einstein condensates with \mathcal{PT} -and non- \mathcal{PT} -symmetric potentials, Commun. Nonlinear Sci. Numer. Simul. **118** (2023) 107061.
- [69] J. Yang, *Nonlinear Waves in Integrable and Nonintegrable Systems*, SIAM, Philadelphia (2010).
- [70] J. Shen and T. Tang, *Spectral and High-Order Methods with Applications*, Science Press, Beijing (2006).
- [71] L. N. Trefethen, *Spectral Methods in MATLAB*, SIAM, Philadelphia (2000).
- [72] I. V. Barashenkov, D. A. Zezyulin and V. V. Konotop, Jamming anomaly in-symmetric systems, New J. Phys. **18** (2016) 075015.