

# Top upregulated GO terms enriched in microarray or RNAseq

| GO term                                                        | Enriched in | Median difference (microarray - RNAseq) | Raw p-value           | Adjusted p-value      |
|----------------------------------------------------------------|-------------|-----------------------------------------|-----------------------|-----------------------|
| regulation of hematopoietic stem cell proliferation            | Microarray  | 1.67                                    | $5.73 \times 10^{-4}$ | $3.11 \times 10^{-2}$ |
| aflatoxin metabolic process                                    | Microarray  | 1.51                                    | $2.39 \times 10^{-4}$ | $2.40 \times 10^{-2}$ |
| alkaloid metabolic process                                     | Microarray  | 1.51                                    | $1.59 \times 10^{-3}$ | $4.52 \times 10^{-2}$ |
| mycotoxin metabolic process                                    | Microarray  | 1.51                                    | $2.39 \times 10^{-4}$ | $2.40 \times 10^{-2}$ |
| protein oxidation                                              | Microarray  | 1.51                                    | $8.92 \times 10^{-4}$ | $3.55 \times 10^{-2}$ |
| lipid hydroxylation                                            | Microarray  | 1.45                                    | $9.66 \times 10^{-4}$ | $3.69 \times 10^{-2}$ |
| positive regulation of cilium movement                         | Microarray  | 1.33                                    | $6.11 \times 10^{-4}$ | $3.12 \times 10^{-2}$ |
| positive regulation of cilium-dependent cell motility          | Microarray  | 1.33                                    | $6.11 \times 10^{-4}$ | $3.12 \times 10^{-2}$ |
| positive regulation of flagellated sperm motility              | Microarray  | 1.33                                    | $6.11 \times 10^{-4}$ | $3.12 \times 10^{-2}$ |
| olefinic compound biosynthetic process                         | Microarray  | 1.31                                    | $1.23 \times 10^{-3}$ | $3.75 \times 10^{-2}$ |
| positive regulation of protein deubiquitination                | RNAseq      | -1.93                                   | $4.46 \times 10^{-4}$ | $2.70 \times 10^{-2}$ |
| negative regulation of PERK-mediated unfolded protein response | RNAseq      | -1.78                                   | $1.11 \times 10^{-3}$ | $3.69 \times 10^{-2}$ |
| vascular endothelial growth factor production                  | RNAseq      | -1.78                                   | $3.72 \times 10^{-5}$ | $1.51 \times 10^{-2}$ |
| positive regulation of T-helper 1 type immune response         | RNAseq      | -1.67                                   | $1.19 \times 10^{-3}$ | $3.69 \times 10^{-2}$ |
| positive regulation of glial cell proliferation                | RNAseq      | -1.65                                   | $1.19 \times 10^{-3}$ | $3.69 \times 10^{-2}$ |
| positive regulation of hormone biosynthetic process            | RNAseq      | -1.56                                   | $1.49 \times 10^{-4}$ | $2.00 \times 10^{-2}$ |
| regulation of T-helper 1 type immune response                  | RNAseq      | -1.55                                   | $2.02 \times 10^{-4}$ | $2.40 \times 10^{-2}$ |
| interleukin-6-mediated signaling pathway                       | RNAseq      | -1.50                                   | $5.31 \times 10^{-5}$ | $1.73 \times 10^{-2}$ |
| cytokine production                                            | RNAseq      | -1.48                                   | $7.11 \times 10^{-4}$ | $3.37 \times 10^{-2}$ |
| regulation of miRNA catabolic process                          | RNAseq      | -1.46                                   | $1.43 \times 10^{-4}$ | $2.00 \times 10^{-2}$ |