

Funciones de prueba, definidas en el intervalo [-10,10]

Nombre de la función	Ref.	Fórmula	Punto mínimo	Valor mínimo
Alpine 1 Function	[1]	$f_1(x) = \sum_{i=1}^D  x_i \sin(x_i) + 0.1x_i $	$x^* = f(0,0)$	$f(x^*)=0$
Dixon & Price Function	[2]	$f_2(x) = (x_1 - 1)^2 + \sum_{i=2}^D i(2 \sin(x_i) - x_{i-1})^2$	$x^* = f(2(\frac{2^i-2}{2^i}))$	$f(x^*)=0$
Quintic Function	[3]	$f_3(x) = \sum_{i=1}^D  x_i^5 - 3x_i^4 + 4x_i^3 - 2x_i^2 - 10x_i - 4 $	$x^* = f(-1 \text{ or } 2)$	$f(x^*)=0$
Schwefel 2.23 Function	[4]	$f_4(x) = \sum_{i=1}^D x_i^{10}$	$x^* = f(0,0)$	$f(x^*)=0$
Stretched V Sine Wave Function	[5]	$f_5(x) = \sum_{i=1}^{D-1} (x_{i+1}^2 + x_i^2)^{0.25} \left[ \sin^2 \left\{ 50(x_{i+1}^2 + x_i^2)^{0.1} \right\} + 0.1 \right]$	$x^* = f(0,0)$	$f(x^*)=0$
Sum Squares Function	[6]	$f_6(x) = \sum_{i=1}^D ix_i^2$	$x^* = f(0,0)$	$f(x^*)=0$

## References

1. Rahnamayan, S., H.R. Tizhoosh, and M.M. Salama, *A novel population initialization method for accelerating evolutionary algorithms*. Computers & Mathematics with Applications, 2007. **53**(10): p. 1605-1614.
2. Dixon, L.C.W. and R. Price, *Truncated Newton method for sparse unconstrained optimization using automatic differentiation*. Journal of Optimization Theory and Applications, 1989. **60**(2): p. 261-275.
3. Mishra, S.K., *Global optimization by differential evolution and particle swarm methods: Evaluation on some benchmark functions*. 2006.
4. Schwefel, H.P., *Numerical Optimization for Computer Models*. 1981: John Wiley & Sons.
5. Schaffer, J.D., et al., *A study of control parameters affecting online performance of genetic algorithms for function optimization, in 3rd International Conference on Genetic Algorithms*. 1989: San Mateo, California. p. 51-60.
6. Jamil, M. and X.-S. Yang, *A literature survey of benchmark functions for global optimisation problems*. International Journal of Mathematical Modelling and Numerical Optimisation, 2013. **4**(2): p. 150-194.