

Inverse CDF method

Suppose that we have a continuous random variable X having Cumulative Density Function (CDF) F_X . Then, the random variable

$$U = F_X(X)$$

has a Uniform distribution $U \sim U_{[0,1]}$. So to generate a random variate x from the distribution of X , we can use the following transformation

$$F_X^{-1}(U) = x$$

Weibull example

Example We want to generate a sample of 10,000 random variates from a Weibull distribution $W(5, 2)$ using the Inverse CDF method.

If $X \sim W(\alpha, \beta)$ having PDF and CDF defined respectively as

$$\text{PDF} \quad f_X(x) = \frac{\alpha}{\beta} \left(\frac{x}{\beta} \right)^{\alpha-1} \exp\left(- \frac{x}{\beta} \right)^{\alpha} \mathbf{1}_{\mathbb{R}^+}(x)$$

$$\text{CDF} \quad F_X(x) = 1 - \exp\left(- \frac{x}{\beta} \right)^{\alpha}$$

Weibull example solved

Then by the Inverse CDF, we can generate realizations of X by posing $U = F_X(x)$ and solving for x . We have

$$U = 1 - \exp\left(-\frac{x}{2}\right)^5$$

$$1 - U = \exp\left(-\frac{x}{2}\right)^5$$

$$-\ln(1 - U) = \left(\frac{x}{2}\right)^5$$


$$(-\ln(1 - U))^{1/5} = \frac{x}{2}$$

$$2(-\ln(1 - U))^{1/5} = x$$

Code for Weibull realizations using the inverse CDF in R

```
1 # Inverse CDF function
2 Inverse.CDF.Weibull <-function(n, alpha, beta) {
3   u <- runif(n)                                # generate uniform numbers
4   data <- beta*((-log(1-u))^(1/alpha))          # formula derived
5   return(data.frame(data))                     # return the data
6 }
7
8 # realizations and plot
9 set.seed(2023)
10 dataset <- Inverse.CDF.Weibull(n = 1000, alpha = 5, beta = 2)
11 head(dataset)
12 #      data
13 # 1 1.8226044
14 # 2 1.6719235
15 # 3 1.4157089
16 # 4 1.7441408
17 # 5 0.9975117
18 # 6 1.3275159
```


Histogram in R



Code for Weibull realizations using the inverse CDF in Python

```
1 # Inverse CDF function
2 def Inverse_CDF_Weibull(n, alpha, beta) :
3     u = np.random.uniform(low=0.0, high=1.0, size=n)           # generate
4     data = beta*((-np.log(1-u))**(1/alpha))                     # formula
5     derived
6     return pd.DataFrame(data = data, columns = ['data'])       # return a
7     data frame instead of an array
8 # realizations and plot
9 np.random.seed(2023)
10 dataset = Inverse_CDF_Weibull(n = 1000, alpha = 5, beta = 2)
11 dataset
12 #    data
13 #0  1.655497
14 #1  2.343973
15 #2  1.952544
16 #3  1.340683
17 #4  1.372833
18 #... ...
19 #995  2.069098
20 #996  2.056594
21 #997  1.996904
22 #998  2.001000
23 #999  1.925175
```

Histogram in Python



Further reading and code

Rizzo, M.L. (2019). Statistical Computing with R, Second Edition (2nd ed.). Chapman and Hall/CRC.

<https://doi.org/10.1201/9780429192760>

The R Project for Statistical Computing:

<https://www.r-project.org/>

Python:

<https://www.python.org/>