

10.007: Modeling the Systems World

Lecture 3

Unconstrained Optimization (Convex Objective Function)

Term 3, 2017



Overview



- Convexity and global optimality.
- “Easy” problems.



An example

Find the global minimum of the following problem:

$$\left. \begin{array}{ll} \min & x^2 + 3x + y^2 - 2y \\ \text{s.t.:} & \end{array} \right\} \begin{array}{ll} x^2 + y^2 & \leq 4 \\ x - y & \leq 0 \end{array}$$



An example

Find the global minimum of the following problem:

$$\left. \begin{array}{ll} \min & x^2 + 3x + y^2 - 2y \\ \text{s.t.:} & \\ & x^2 + y^2 \leq 4 \\ & x - y \leq 0 \end{array} \right\}$$

How do we
proceed?




An example

Find the global minimum of the following problem:

$$\left. \begin{array}{ll} \min & x^2 + 3x + y^2 - 2y \\ \text{s.t.:} & \end{array} \right\} \begin{array}{ll} x^2 + y^2 \leq 4 \\ x - y \leq 0 \end{array}$$

How do we
proceed?





Relaxation of the problem

We can first solve the unconstrained problem:

$$\min \quad x^2 + 3x + y^2 - 2y$$

Relaxation of the problem



We can first solve the unconstrained problem:

$$\min \quad x^2 + 3x + y^2 - 2y$$

How are these
problems related?



Relaxation of the problem

We can first solve the unconstrained problem:

$$\min \quad x^2 + 3x + y^2 - 2y$$

How are these
problems related?

The relaxed (unconstrained) problem must have an objective function value at least as good as the constrained problem!
(Can you see why?)



Solving the relaxation

Let $f(x, y)$ be the objective function, and set $\nabla f = 0$:

$$\begin{aligned}\frac{\partial f}{\partial x} &= 2x + 3 = 0 \\ \frac{\partial f}{\partial y} &= 2y - 2 = 0.\end{aligned}$$

Solution: $x = -3/2, y = 1$. Compute the Hessian at $(-3/2, 1)$:

$$\det \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} = 4 > 0.$$


Local minimum at $(-3/2, 1)$, it is **inside** the feasible region.

Therefore, it is a local minimum to the constrained problem too!

Is this a **global** minimum?



An example



Go global!




Solving an optimization problem typically requires finding a **global optimum**.

However, it is **very difficult** to write algorithms that find global optima. In most cases, we can only guarantee **local** optimality.

There are problems for which global optimality follows from local optimality: these problems are considered **“easy”**. This is the “ideal” situation.



Convexity: intuition



A function is convex if the line between $(y, f(y))$, $(z, f(z))$ lies above the graph of the function over the segment \overline{yz} .



Convexity

Definition

A function $f(x) : \mathbb{R}^n \rightarrow \mathbb{R}$ is called **convex** if for every $y, z \in \mathbb{R}^n$ and every scalar $0 \leq \lambda \leq 1$ we have:

$$f(\lambda y + (1 - \lambda)z) \leq \lambda f(y) + (1 - \lambda)f(z).$$

It is called **strictly convex** if for every $y \neq z \in \mathbb{R}^n$ and every scalar $0 < \lambda < 1$ we have:

$$f(\lambda y + (1 - \lambda)z) < \lambda f(y) + (1 - \lambda)f(z).$$


The expression $\lambda y + (1 - \lambda)z$ is simply the line segment between y and z when λ ranges from 0 to 1.



Concavity

Definition

A function f is **concave** if $-f$ is convex.



Necessary and Sufficient Conditions for Minimizing Convex Functions



Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **convex** function, and assume that f is continuously differentiable at point x^* . Then x^* is a global minimum of f **if and only if** $\nabla f(x^*) = \vec{0}$.



Necessary and Sufficient Conditions for Minimizing Convex Functions



Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **convex** function, and assume that f is continuously differentiable at point x^* . Then x^* is a global minimum of f **if and only if** $\nabla f(x^*) = \vec{0}$.

Sketch of Proof: Think about how a single variable convex function looks, the tangent line is always 'below' the function. When the tangent line is horizontal at a point $(x^*, f(x^*))$, all the function values are greater than or equal to $f(x^*)$.





Necessary and Sufficient Conditions for Minimizing Convex Functions



Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **convex** function, and assume that f is continuously differentiable at point x^* . Then x^* is a global minimum of f **if and only if** $\nabla f(x^*) = \vec{0}$.

Sketch of Proof: Think about how a single variable convex function looks, the tangent line is always ‘below’ the function. When the tangent line is horizontal at a point $(x^*, f(x^*))$, all the function values are greater than or equal to $f(x^*)$.



This means that we have a very simple procedure to find the global minimum of a convex and differentiable function.



Optimality conditions for convex functions

Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a twice continuously differentiable **convex** function. We want to solve: $\min_{x \in \mathbb{R}^n} f(x)$.

Because the function is convex, the approach becomes simpler:

1. Find all x^* such that $\nabla f(x^*) = \vec{0}$. These are **global minima**.
2. (Optional unless requested) Check if $H_f(x^*)$ is pd. If yes, x^* is a **strict** global minimum.

If we are not interested in checking if global minima are strict, solving Step 1 is sufficient. As before, Step 1 is often **very hard** even for the convex case.



Back to the example

Recall our initial problem:

$$\left. \begin{array}{ll} \min & x^2 + 3x + y^2 - 2y \\ \text{s.t.:} & \end{array} \right\} \begin{array}{ll} x^2 + y^2 & \leq 4 \\ x - y & \leq 0 \end{array}$$

The objective function is **convex** (sum of convex functions – see previous cohort). By the Theorem, a local minimum is a **global** minimum.

The point $(-3/2, 1)$ is the **global optimum** of the problem unconstrained problem. It is also feasible for the constrained problem, therefore, it is a global minimum of that as well. We are done!

Summary



- Local/global optima.
- Convex functions .
- Implications of convexity on **global optimality**.